

G9101

**STUDIES ON THE ANTIULCEROGENIC AND
ANTIHEPATOTOXIC POTENTIAL OF CHITOSAN AND
GLUCOSAMINE IN EXPERIMENTAL RATS**

Thesis Submitted to

COCHIN UNIVERSITY OF SCIENCE AND TECHNOLOGY

In partial fulfillment of the requirements for the degree of

**DOCTOR OF PHILOSOPHY
IN BIOCHEMISTRY**

By

SANTHOSH. S., M.Sc



**FISH PROCESSING DIVISION
CENTRAL INSTITUTE OF FISHERIES TECHNOLOGY
(INDIAN COUNCIL OF AGRICULTURAL RESEARCH)
COCHIN-682029**

**FACULTY OF MARINE SCIENCES
COCHIN UNIVERSITY OF SCIENCE AND TECHNOLOGY
COCHIN**

OCTOBER 2006

Telephone } 2666845, 266846, 266847, 266848, 266849
2666763, 2666764, 2666765, 2666766
2668576, 2668577, 2668578, 2668579, 2668580

www.cift.igss.in
Fax : 0091-484-2668212
E-mail : enk_ciftaris@sancharnet.in
cift@ciftmail.org



केन्द्रीय मात्स्यकी प्रौद्योगिकी संस्थान
CENTRAL INSTITUTE OF FISHERIES TECHNOLOGY

भारतीय कृषि अनुसंधान परिषद
(Indian Council of Agricultural Research)
विलिंगडन आईलैंड, मत्स्यपुरी पी.ओ., कोच्चिन - 682 029
Willingdon Island, Matsyapuri P.O., Cochin - 682 029



CERTIFICATE

This is to certify that this thesis entitled **“STUDIES ON THE ANTIULCEROGENIC AND ANTIHEPATOTOXIC POTENTIAL OF CHITOSAN AND GLUCOSAMINE IN EXPERIMENTAL RATS”** embodies the original research work conducted by **Mr. Santhosh. S.** under my guidance from 16.10.2002 to 30.07.2006. I further certify that no part of this thesis has previously been formed the basis of award of any degree, diploma, associateship, fellowship or any other similar titles of this or in any other university or institution. He has also passed the Ph.D qualifying examination of the COCHIN UNIVERSITY OF SCIENCE AND TECHNOLOGY, Cochin held in November 2004.

Cochin-29,
26-10-2006

MS Mathew

Dr. P.T. Mathew
Principal Scientist
Fish Processing Division
Central Institute Of Fisheries Technology
Cochin-682029.

ABSTRACT

The present study examined the antiulcer effect of glucosamine on mucosal antioxidant defense system in ibuprofen-induced peptic ulcer in male albino rats. The number of lesions in the gastric mucosa, volume of gastric juice, acid output, pepsin activity, lipid peroxides, reduced glutathione content, the activities of glutathione dependent antioxidant enzymes (glutathione peroxidase and glutathione-S-transferase) and antiperoxidative enzymes (catalase and superoxide dismutase), activities of membrane bound ATPases, protein content, glycoprotein components, minerals status, SDS-PAGE electrophoretic pattern of mucosal protein, amino acid composition and histopathology were determined. Prior oral administration of chitosan and glucosamine significantly prevented the ibuprofen-induced increases in the number of lesions in the gastric mucosa, volume of gastric juice and acidity. They also maintained the activity of pepsin at near normal level. Oral pretreatment of chitosan and glucosamine exerted a significant antioxidant effect by preventing ibuprofen-induced lipid peroxidation and by maintaining the level of reduced glutathione and the activities of mucosal antioxidant enzymes at near normalcy. Prior oral administration of glucosamine significantly prevented the ibuprofen-induced depletion of protein and glycoprotein components and maintained the activities of membrane bound ATPases as compared to untreated ulcer induced group of rats. Amino acid composition and mineral status were also brought to the near normal level in the chitosan and glucosamine pretreated groups. The antiulcerogenic activity of glucosamine might be ascribable to its ability to neutralize the hydrochloric acid secreted into the stomach and to its capability to strengthen the mucosal barrier by increasing mucosal glycoprotein synthesis and to its free radical scavenging property. Histopathological investigations and amino acid observations of the mucosal tissue were also supported the antiulcerogenic effect of chitosan and glucosamine. The antioxidant capacity of chitosan and glucosamine to inhibit ibuprofen-induced lipid peroxidation is also supportive to the antiulcer effect.

Tuberculosis is a dangerous disease and its death toll is increasing year by year. Intake of isoniazid and rifampicin, the most common antitubercular drugs, lead to fatal hepatotoxic condition. I have studied the protective effect of chitosan and glucosamine supplementations against the hepatotoxicity induced by antitubercular drugs with respect to the changes in the levels of protein, SDS PAGE-electrophoretic pattern protein, levels of diagnostic marker enzymes, lipid components, lipid peroxidation, albumin, globulin, albumin/globulin ratio, glucose, creatinine and urea in the serum and diagnostic marker enzymes, hepatic thiols, lipid components, lipid peroxidation, reduced glutathione, the activities of glutathione dependent antioxidant enzymes and antiperoxidative enzymes, activities of membrane bound ATPases, protein content, glycoprotein components, minerals status, glycoprotein conjugates, electrophoretic pattern of protein, fatty acid composition and histopathology in the liver tissue of normal and experimental groups of rats. Co-administration of chitosan and glucosamine was found to prevent significantly the antitubercular drugs-induced changes in the levels of diagnostic marker enzymes and albumin/globulin ratio in experimental groups of rats. Isoniazid and rifampicin-induced lipid peroxidation was also found to be prevented by the administration of chitosan and glucosamine. The oral administration of antitubercular drugs caused a significant elevation in the levels of diagnostic marker enzymes and cholesterol, triglycerides, free fatty acids and lipid peroxidation in serum and liver of experimental rats. There was a slight decline in the level of phospholipids in liver tissue also observed. Chitosan and glucosamine exerted a significant antilipidemic effect against isoniazid and rifampicin-induced hepatitis by maintaining the levels cholesterol, triglycerides, free fatty acids and phospholipids in serum and liver at near normalcy. Further, chitosan administration maintained the levels of lipid peroxides, glucose, creatinine, albumin/globulin, urea, hepatic thiols and protein to normalcy in experimental groups compared to hepatotoxicity-induced group of rats. Levels of glycoconjugates were also maintained to near normal level by chitosan and glucosamine co-administration. From the results obtained, it can be concluded

that co-chitosan and glucosamine are beneficial against antitubercular drugs-induced hepatotoxicity. SDS page electrophoretic pattern, fatty acid composition and histopathological observations also support this finding.

LIST OF TABLES

Table	Title
Table 2.1	Chitin content in various sources
Table 4.1.1	Number of lesions, volume of gastric juice, and acid output and pepsin activity of the gastric mucosa of normal and experimental groups of rats
Table 4.1.2	Activities of membrane bound ATPases of the gastric mucosa of normal and experimental groups of rats
Table 4.1.3	Levels of reduced glutathione and lipid peroxides in the gastric mucosa of normal and experimental groups of rats
Table 4.1.4	Activities of antioxidant enzymes in the gastric mucosa of normal and experimental groups of rats
Table 4.1.5	Mineral status of the stomach of normal and experimental groups of rats
Table 4.1.6	Levels of various amino acids in the stomach of normal and experimental groups of rats
Table 4.2.1	Levels of diagnostic marker enzymes in the serum of normal and experimental groups of rats
Table 4.2.2	Levels of urea, albumin, globulin, albumin/globulin ratio in the serum of normal and experimental groups of rats
Table 4.2.3	Levels of creatinine and glucose in the serum of normal and experimental
Table 4.2.4	Levels of glycoconjugates in the liver of normal and experimental groups of rats
Table 4.2.5	Levels of lipid peroxides in the serum and liver of normal and experimental groups of rats
Table-4.2.6	Levels of antioxidant enzymes in the liver of normal and experimental groups of rats
Table 4.2.7	Activities of membrane bound ATPases in the liver of normal and experimental groups of rats
Table 4.2.8	Levels of total thiols and free thiol in the liver of normal and experimental groups of rats
Table 4.2.9	Levels of lipid components in the serum or normal and experimental groups of rats
Table 4.2.10	Levels of lipid components in the liver normal and experimental groups of rats
Table 4.2.11	Fatty acid composition of the liver of normal and experimental groups of rats
Table 4.2.12	Levels of calcium, magnesium and zinc in the liver of normal and experimental groups of rats

LIST OF FIGURES

Figure	Title
Fig.2.1.	Structure of chitosan
Fig.2.1.	Structure of β -D-Glucosamine
Fig.2.3.	Structure of stomach
Fig.2.4.	<i>Helicobacter pylori</i>
Fig.2.5.	Structure of ibuprofen
Fig.2.6.	Fig. 2.6. The Cyclooxygenase Pathway
Fig.2.7.	Structure of isoniazid
Fig.2.8.	Metabolic pathway of Isoniazid
Fig.2.9.	Structure of rifampicin
Fig.4.1.1.	Levels of mucosal protein in experimental and normal rats.
Fig.4.1.2.	Levels of hexose and hexosamine in the stomach tissue of control and treated animals.
Fig.4.1.3.	Sialic acid levels in the stomach of normal and experimental groups of rats.
Fig.4.1.4.	Levels of sodium and potassium in the stomach of normal and experimental groups of rats.
Fig.4.2.1.	Levels of ALT and AST in the liver of normal and experimental groups of rats.
Fig.4.2.2.	Levels of protein in the serum of normal and experimental groups of rats.
Fig.4.2.3.	Protein levels in the liver of normal and experimental rats.
Fig.4.2.4.	Levels of HDL-cholesterol and LDL-cholesterol in the serum of normal and experimental groups of rats.
Fig.4.2.5.	Levels of sodium and potassium in the liver of normal and experimental groups of rats.

LIST OF PLATES

Plate	Title
Plate 4.1.1	Gross pathology of normal stomach in control rat
Plate 4.1.2	Gross pathology of ulcerated stomach of rat
Plate 4.1.3.	Gross pathology of chitosan pre-treated stomach
Plate 4.1.4.	Gross pathological view of glucosamine pre-treated stomach.
Plate 4.1.5.	Histopathological architecture of the stomach tissue of normal rat
Plate 4.1.6.	The architecture of stomach tissue of chitosan control rat
Plate 4.1.7.	The normal stomach architecture of glucosamine administered rat
Plate 4.1.8.	Ibuprofen induced ulcerated stomach tissue
Plate 4.1.9.	Near normal architecture of stomach in chitosan pre treated animal
Plate 4.1.10.	Architecture of stomach tissue of glucosamine treated rats.
Plate 4.1.11.	The SDS-polyacrylamide gel electrophoretic pattern of stomach proteins
Plate 4.2.1.	Histopathological view of normal liver tissue of normal and experimental groups of rat
Plate 4.2.2.	The histopathological view of the liver tissue of chitosan control animal
Plate 4.2.3.	The histopathological architecture of the liver tissue of glucosamine alone fed rat
Plate 4.2.4.	The histopathology of the liver tissue of antitubercular drugs-administered rats.
Plate 4.2.5.	Histopathology of liver tissue of Chitosan co-administered rats
Plate 4.2.6.	Architecture of the liver tissue of glucosamine co-administered rat
Plate 4.2.7.	SDS-page electrophoretic pattern of serum protein
Plate 4.2.8.	Electrophoretic pattern of liver protein in normal and experimental groups of rat

ABBREVIATIONS

ANSA	- 1-amino-2-naphthol-4-sulphonic acid
AR	- Analar
ATP	- Adenosine triphosphate
BSA	- Bovine serum albumin
°C	- Degree celsius
CAT	- Catalase
CDNB	- 1-chloro-2,4-dinitro benzene
dl	- Decilitre
DNPH	- 2,4-dinitrophenyl hydrazine
DTNB	- 5,5-dithio-bis-(2-nitro benzoic acid)
EDTA	- Ethylene diamine tetra acetic acid
Fig.	- Figure
FFA	- Free fatty acid
g	- grams
GPx	- Glutathione peroxidase
GSH	- Glutathione
HCl	- Hydrochloric acid
H ₂ O ₂	- Hydrogen peroxide
H ₂ SO ₄	- Sulphuric acid
h	- Hour
HDL	- High density lipoprotein
i.p.	- Intra peritoneal
Kg	- Kilogram
KOH	- Potassium hydroxide
KI	- Potassium iodide.
LDH	- Lactate dehydrogenase.
LDL	- Low density lipoprotein
M	- Molar
mM	- Millimolar.

mg	- Milligram
Mm	- Millil-molar.
mm	- Millimeter
N	- Normal
μ moles	- Micro molesmin
Min	- Minute
ml	- Militer
NADH	- Reduced nicotinamide adenine dicutedldiacid
NADP	- Nictonamine
Pi	- Inorganic phohpshris.
OD	- Optical denstiry
SD	- Standard division
SOD	- Superoxide dismutase
TBA	- Thio barbituric acid.
v/v	- Volume/volume
VLDL	- Very low density lipoprotein
w/v	- Weight/volume
w/w	- Weight/weight

CONTENTS

Sr. No	Titles	Page No.
1.	INTRODUCTION	1
2.	REVIEW OF LITERATURE	4
2.1.	Chitosan	4
2.2.	Glucosamine	8
2.3.	Peptic Ulcer	11
2.3.1.	Ibuprofen	16
2.4.	Hepatotoxicity	20
2.5.	Antitubercular drugs	25
2.5.1.	Tuberculosis	25
2.5.2.	Isoniazid	27
2.5.3.	Rifampicin	30
3.	MATERIALS AND METHODS	34
4.	RESULTS AND DISCUSSION	75
4.1.	ANTI ULCEROGENIC POTENTIAL OF CHITOSAN AND GLUCOSAMINE	75
4.2.	ANTI HEPATOTOXIC POTENTIAL OF CHITOSAN AND GLUCOSAMINE	84
5.	SUMMARY AND CONCLUSION	98
6.	REFERENCES	102

1. INTRODUCTION

Disease and sickness have been a part of mankind's progress through the ages. It comes as no surprise that a substantial portion of the intellectual efforts of every historical age has been directed at understanding disease and finding cures. It is but natural that each of the formally organized and structured attempts has to depend on the knowledge base and paradigms that were available at the particular period of time when the system evolved and matured. As our view of the world changed, largely due to the efforts of science, newer systems of medicine emerged and older ones fell out of favour or even discarded. Over the last century or less, keeping pace with the advances made by modern science, the Western or allopathic medical system has been the dominant influence on the health care scene. There is no denying its contribution to our well-being. Yet, there remains the lingering reality that no single system of health care delivery is able to provide all the answers to health and disease.

Traditional medicines have been the starting point for the discovery of many important modern drugs (Patwardhan, 2005). This fact has led to chemical and pharmacological investigations and general biological screening programs for natural products all over the world. Ocean, the biggest wonder in the universe, covers almost three quarters of the surface of earth. It is not only a great source of a lot of beautiful living and non-living things, but also a treasure of plenty of substances, which are beneficial in all faces of our lifestyle and in our needs of day-to-day life. Marine organisms produce many bioactive substances, which are having a lot of potential biomedical and pharmaceutical applications.

Marine food has become an inevitable ingredient in our daily menu. Prawn shell, a main waste material of marine food industry, can be efficiently utilized by converting it into various useful marine polysaccharides byproducts such as chitin, chitosan, glucosamine etc. (Synowiecki and Al-Khateeb, 2003). The mucopolysaccharide, chitin is produced by the demineralization and deproteinisation of shrimp shell waste. It is considered as the second most

abundant biopolymer in the world next to cellulose (Sini *et al.*, 2005). Chitin can be converted into two value-added products, chitosan and glucosamine. Chitosan and glucosamine are having immense applications in the fields of biomedical, pharmaceutical and nutraceutical industries.

Chitosan [poly- (β -1-4)-D-glucosamine] is prepared by the deacetylation of chitin using concentrated alkali (Krisana *et al.*, 2004; Shepherd *et al.*, 1997). It has profound applications in the fields of clarification and purification, chromatography, paper and textiles, photography, food and nutrition, agriculture, pharmaceutical and medical, cosmetics, biodegradable membranes and biotechnology. It has been reported to possess immunological (Nishimura *et al.*, 1984; Mori *et al.*, 1997) antibacterial (Tokura *et al.*, 1997; Tanigawa *et al.*, 1992) and wound healing activities (Okamoto *et al.*, 1993; Kweon *et al.*, 2003; Khnor and Lim, 2003). Chitosan is an effective and adequate haemostatic agent even under the most severe conditions of anticoagulation. Chitosan solution is found beneficial for healing "athletes foot" conditions (Allan *et al.*, 1984). Both hard and soft contact lenses can be made from chitosan. Sapelli *et al.* (1986) studied the application of chitosan in dentistry.

Han *et al.*, (1999) has reported that supplementation of chitosan can reduce high fat diet-induced lipidemia by its antilipidemic property. The free radical quenching property of this marine polysaccharide has also been studied in detail (Xing *et al.*, 2005). Reports by Filipovic-Grcic *et al.*, (2001) indicates the membrane stabilizing property of chitosan. In our laboratory, Anandan *et al.* observed the antiulcerogenic potential of chitosan against HCl-ethanol mixture induced peptic ulcer in rats. Since chitosan can wrap solid particles in liquids to bring them together and agglomerate, it is effectively utilized for the controlled release of drug (Krisana *et al.*, 2004). Dissolution properties and bioavailability of poorly soluble drugs can be improved by grinding them with chitosan. Though the beneficial effects of chitosan have been extensively studied, the antihepatotoxic potential of chitosan on lipid metabolism in antitubercular drugs-induced hepatotoxicity has not yet been explored. Muzzarelli (1996) has reported that

supplementation of chitosan can prevent experimentally induced diabetic complications by its antioxidant and membrane stabilizing properties.

Glucosamine is another value added by-product prepared by the hydrolysis of chitin with conc. HCl. This is an amino monosaccharide having a lot of potential biomedical, pharmaceutical and industrial applications. It is an important constituent of cartilage and it has been used for the treatment of osteoarthritis (Timothy *et al.*, 2004). In combination with chondroitin sulfate, it can build blocks for cartilage, up-regulate chondrocyte and reduce the extent of cartilage degradation (Yu Shao *et al.*, 2004). Sal'nikova *et al.* (1990) suggests the antioxidant nature of glucosamine, by virtue of which it can trap and dismutate free radicals. Glucosamine is an essential component required for glycoprotein synthesis in living beings (Wu *et al.*, 2004). Mutoh *et al.* (1995) reported the cytoprotectivity of glucosamine rich mucus.

In traditional medicine, the shrimp and cuttlefish exoskeleton powder has been used to cure arthritis, diabetes, stomach disorders, epilepsy and various liver disorders. Yet there is little documentary evidence regarding the exact mechanism of action involved in the cytoprotective effect of these marine polysaccharides. Glucosamine is the basic unit of chitin and chitosan and it is an essential component required for glycoprotein synthesis in living beings (Zhang *et al.*, 2004). The beneficial actions of chitin and chitosan may be ascribable to their basic unit glucosamine. Hence, it thought to be important to study the protective effect chitosan and glucosamine on experimentally induced ulcer and hepatotoxic conditions to derive a conclusion regarding the exact biochemical mechanism involved in the beneficial effects of shrimp and cuttle fish exoskeleton powder in alleviating the ulcerative and liver disorders by virtue of their antilipidemic, antioxidant and membrane stabilizing properties.

2. REVIEW OF LITERATURE

2.1. Chitosan

Chitosan is a natural mucopolysaccharide (Van der Lubben *et al.*, 2001) of marine origin. While it has been in existence for millennia, its current form has just recently been prepared. Chitosan is one of the most available polysaccharides with positive charges found in nature (Xing *et al.*, 2005). Technically speaking, Chitosan is a naturally occurring substance that is chemically similar to cellulose (Barbara, 2005), which is a plant fiber. Chitosan is shown to have superior characteristics and especially flexibility in its use.

2.1.1. Source

Chitosan is obtained from chitin (Shepherd *et al.*, 1997; Krisana *et al.*, 2004), which is believed to be the second most abundant biomaterial after cellulose (Sini *et al.*, 2005). Chitin is widely distributed in the nature. Major sources of chitin are shellfish, clams, krill, oysters, squid, fungi and insects. Chitin content in various sources is given in Table 1 (Allan *et al.*, 1978; Kong, 1975). Amongst several sources, the exoskeleton of crustaceans consist of 15-20 % chitin by dry weight. Chitin found in nature is a renewable bio-resource. It is present not only in animals, but also in plants as cell wall material. Chitin resembles cellulose both in chemical structure and in biological function as a structural polymer. The crystalline structure of chitin has been shown to be similar to cellulose in the arrangements of inter- and intra chain hydrogen bonding. The annual biosynthesis of chitin has been estimated to 10^9 to 10^{11} tons. Chitosan is found in nature, to a lesser extent than chitin, in the cell walls of fungi. Zygomycetes contain both chitin and chitosan (Austin *et al.*, 1981; Rudall, 1969).

Chitosan is prepared by alkaline N-deacetylation of chitin (Kittur *et al.*, 2002). Deacetylation of chitin is done by using concentrated NaOH solution (40-50%).

Source	Quantity harvested 10 ³ tonnes	Chitinous waste, 10 ³ tonnes	Chitin potential, 10 ³ tonnes
Shell fish	1700	468	39
Krill (potential landing)	18200	3640	56
Clam/ Oysters	1390	521	22
Squid	660	99	1
Fungi	790	790	32
Total	22,740	5,118	150

Table 2.1. Chitin content in various sources

2.1.2. Properties

Chitosan occurs as odourless substance. It is an amorphous solid and off-white in colour. The properties of chitosan vary considerably depending on the source and production process. Average molecular weight of chitosan is around 1.2×10^5 Daltons. Most of the commercial polysaccharides like cellulose, dextran, pectin, alginic acid etc. are neutral or acidic. But chitosan is an abundant basic polysaccharide. Its pH comes around 8 and this basic nature makes it unique for different applications. Chitosan has special properties like polyelectrolyte behaviour, polyoxysalt formation, film formation and chelation of metal ions (Austin *et al.*, 1981). The main advantage of Chitosan for different biological applications is that it is nontoxic (Arai *et al.*, 1968) in nature. Chitosan is insoluble in water but soluble in dilute acids. Pure chitosan is not hydrolysed by lysozyme


while chitin is hydrolysed. Chitosan undergoes the typical reactions of amines, of which N-acetylation and Schiff reaction are the most important. Although β (1-4) anhydroglucosidic bond of chitosan is also present in cellulose, the characteristic properties of chitosan are not shared by cellulose (Muzarelli, 1978). Chitosan forms aldimines and ketimines respectively with aldehydes and ketones at room temperature. When reacted with ketoacids followed by reduction with sodium borohydride, chitosan produces glucans. N-carboxymethyl chitosan and N-carboxyl benzyl chitosan are examples of glucans derived from chitosan (Muzarelli, *et al.*, 1982 a,b; Muzarelli and Tanfani, 1982). On hydrogenation with aldehydes, chitosan produces N-alkyl chitosans. In spite of the hydrophobicity of the alkyl chains, N-alkyl chitosans swell in water, but its film forming capacity is retained (Muzarelli *et al.*, 1983). In acidic medium, chitosan develops a violet colour with iodine (Van Wisselingh, 1898; Campbell, 1929; Krishnan and Sundara Rajulu, 1964). This test is used as the detection test for chitosan. An enzymatic method was also put forwarded by Jeuniaux (1963 & 1965) for the detection of chitosan. As per the reports of Sundara Rajulu *et al.* (1982), enzymatic test was found more reliable amongst these two tests. Chitosan can be hydrolysed in the presence of chitosanase (Monaghan *et al.*, 1972). According to Fenton and Eveleigh (1981) purified chitosanase from microbes hydrolyses specifically chitosan.

2.1.3. Structure

Chitin, source of chitosan, is a polymer of β (1-4)-N-acetyl-D-glucosamine (Fig. 2.1). Chitin occurs in three polymorphic forms, which differ in the arrangement of molecular chains (Muzarelli, 1977 a). In chitosan, the N-acetyl group of chitin is removed. Chitosan is a polymer of β (1-4)-D-glucosamine (Yasser and Ahmed, 2002)

2.1.4. Applications of chitosan

Since chitosan is nontoxic (Arai *et al.*, 1968), it is suitable for different applications. Chitosan is one of the most abundant natural amino polysaccharides with variety of applications (Majeti and Kumar, 2000) It is having



Chitosan

Fig.2.1. Structure of Chitosan

immunological (Nishimura *et al.*, 1984; Mori *et al.*, 1997) antibacterial (Tokura *et al.*, 1997; Tanigawa *et al.* (1992) and wound healing activities (Okamoto *et al.*, 1993; Kweon *et al.*, 2003; Khnor and Lim, 2003). As chitosan is having antibacterial, haemostatic, fungistatic, antitumoral and anticholesteremic properties (Barbara, 2005), it has wide pharmaceutical applications. Muzarelli *et al.* (1988) studied the application of chitosan in wound dressing since chitosan wound healing property. Chitosan has profound applications in the fields of clarification and purification, chromatography, paper and textiles, photography, food and nutrition, agriculture, cosmetics, biodegradable membranes and biotechnology.

Since chitosan is a long chain polymer, it can wrap solid particles in liquids to bring them together and agglomerate. Thus, chitosan can be used as a coagulant aid. Commercially, chitosan is used for waste water clarification. Free amino group and hydroxyl groups present in chitosan make it suitable for chromatographic applications. Muzarelli (1977) and Allan *et al.* (1972,1975 & 1977) made studies on the application of chitosan on improving the properties of paper. Chitosan can be used for the production of fibers and films (Kunike, 1926). Since chitosan has resistance to abrasion, optical characteristics and behaviour with silver complexes, it has immense applications in photography. Animal feed studies proved that the utilization of whey might be improved if the diet contain small amount of chitinous material. Chitosan feeding studies conducted in chicken, pig, mice, rabbits etc. improved their weights significantly. A thin coating of chitosan in seeds enhances the seed growth. Chitosan coating can be used for the ripening of fruits also. Chitosan can be effectively utilized for the controlled release of drug (Agnihotri and Aminabhavi, 2004). Chitosan is used as a substituent for microcrystalline cellulose (MCC) as a diluent in tablet making by direct compression. Dissolution properties and bioavailability of poorly soluble drugs can be improved by grinding them with chitosan. Enzyme immobilization is another important use of chitosan. Many methods are adopted for enzyme immobilization like entrapment and adaption, fixing by cross-linking etc. There are many advantages when immobilized enzymes are used; enzymic reaction can be stopped at any desired time and small amount of enzyme is

sufficient for large amount of substrate. Since, chitosan is physiologically harmless it can be used in hair cosmetic products instead of synthetic resins (Gross *et al.*, 1982). Chitosan films have good water absorbing property. Chitosan membranes can be used for desalination, ultrafiltration and wastewater treatment (Rutherford and Dunson, 1984). Chitosan is an effective and adequate haemostatic agent even under the most severe conditions of anticoagulation. Chitosan solution is found beneficial for healing “athletes foot” conditions (Allan *et al.*, 1984). Chitosan has good ophthalmologic applications. Both hard and soft contact lenses can be made from chitosan. The effect of chitosan have been investigated in patients with chronic renal failure undergoing long term stable haemodialysis treatment (Jing *et al.*, 1997) Ingestion of chitosan effectively reduced serum cholesterol level and increased serum haemoglobin levels. Significant reduction in urea and creatinine levels in the serum were observed after 4 weeks of chitosan ingestion. The data suggests that chitosan might be effective in the treatment for renal failure patients. Sapelli *et al.* (1986) studied the application of chitosan in dentistry. Knapczyk *et al.* (1989) proved that chitosan is applicable in immunology. Like some plant fibers, chitosan is not digestible; therefore it has no caloric value. No matter how much Chitosan ingest, its caloric count remains at zero. This is a very important property for any weight loss product.

2.2. Glucosamine

Glucosamine is an amino monosaccharide found in most of the tissues in our body. Unlike other forms of sugar in the body, amino sugars are components of carbohydrates that are incorporated into the structure of body tissues, rather than being used as a source of energy. Glucosamine is involved in the formation of the nails, tendons, skin, eyes, bones, ligaments, and heart valves. It also plays a role in the mucous secretions of the digestive, respiratory, and urinary tracts. It is incorporated in the biosynthesis of glycosaminoglycans and proteoglycans, essential for the extracellular matrix of connective tissues. Glucosamine is a nutritional supplement and therefore the authority of the Food and Drug Administration to regulate it is now severely limited (Angell and Kassirer, 1998).

There have been relatively few trials of glucosamine use in humans, and those that have been published are studies that examined the efficacy of glucosamine in the relief of joint pain (Mc Alindon *et al.*, 2000). There have been no reported trials of long-term glucosamine use, and its long-term safety is unknown. Due to favorable, but scientifically inconclusive, reports of glucosamine therapy for arthritis symptoms are less (Loes *et al.*, 1998). Glucosamine is available as an over-the-counter oral supplement, has few adverse side effects, and is relatively inexpensive. Several clinical studies have reported that glucosamine works better in reducing the symptoms of osteoarthritis (Reginster *et al.*, 2001). However, the mechanisms underlying this effect are not yet known.

2.2.1. Source

Exoskeleton of marine organisms are the main source of glucosamine. Chitin is prepared from the shells of crab, shrimp etc. From chitin, glucosamine is prepared by hydrolysis with concentrated acids under drastic conditions (Madhavan, 1992). Glucosamine is found in hyaluronic acid, a compound responsible for the lubricating and shock-absorbing properties of synovial fluid. Glucosamine is synthesized in chondrocytes in the body from glucose and glutamine.

2.2.2. Structure

Glucosamine is an aminomonosacharide. i.e., amino group is attached to a glucose ring (Fig.2.2). Thus comes the name glucose-amine. Chemically glucosamine is 2-amino-2-deoxy-alpha-D-glucose.

2.2.3. Properties

Glucosamine is a colourless and odourless substance with sweet taste. It is crystalline in nature. It is having acidic pH, ranging from 3.5-4.5 and has density of 0.75 g/ml. Specific rotation of glucosamine is 70° and is readily soluble in water. When heated above 80 °C, it undergoes Maillard reaction and its colour




Fig. 2. 2. Structure of β -D-Glucosamine

will be faded. On further heating, glucosamine will be charred. Glucosamine has anti-inflammatory and antioxidant actions. Glucosamine blocks the generation of superoxide free radicals by macrophages. Glucosamine is well tolerated. No severe side effects were reported for glucosamine (Reginster *et al.*, 2001). Glucosamine can increase proteoglycan synthesis (Corinne *et al.*, 1998), which is a fundamental component found in articular cartilage. Proteoglycans synthesized in the presence of glucosamine had smaller glycosaminoglycan chains with a corresponding decrease in lipoprotein retention. This property makes it useful against atherosclerosis (Tannock *et al.*, 2002). Glucosamine inhibits the cartilage-destructive enzyme collagenase. Glucosamine helps in the synthesis of cartilage by increasing key components of cartilage such as glycosaminoglycans. Various reports confirmed that diabetes patients can also consume Glucosamine, which will not increase blood glucose level (Daren *et al.*, 2003). Glucosamine appears to undergo a significant first-pass effect in the liver, which metabolizes a significant proportion of the dose to CO₂, water, and urea (Setnikar *et al.*, 1993)

2.2.4. Applications

Setnikar and colleagues (1984, 1986, 1993) have published several reports on the pharmacokinetics of glucosamine in rats, dogs, and humans. In both humans and rats, oral administration of radio labeled glucosamine results in absorption of more than 90% of the ingested dose (Tannock *et al.*, 2002).

Glucosamine is found largely in cartilage and plays an important role in its health and resiliency. As we age, we lose some of the glucosamine and other substances in cartilage. This can lead to thinning of cartilage and the onset and progression of osteoarthritis. According to Braham (2003), glucosamine supplementation can provide some degree of pain relief and improved function in persons who experience regular knee pain, which may be caused by prior cartilage injury and/or osteoarthritis. Since, glucosamine can increase the synthesis of proteoglycans and glycosaminoglycans, which are necessary ingredients of connective tissue, it is proven to be effective against osteoarthritis

(Timothy *et al.*, 2004; Reginster *et al.*, 2001). Canapp *et al.* (1999) reported that glucosamine was found beneficial against acute synovitis.

Every year about twenty thousand people die from using NSAIDs. The combination of glucosamine and NSAIDs may reduce the doses needed for anti-inflammatory activity as well as the side effects associated with these NSAID drugs (Zupanets *et al.*, 1991). Dettmer (1979) examined the effects of simultaneous administration of glucosamine and chondroitin sulfate on osteoarthritis. This study demonstrated greater effectiveness of this combination therapy. In combination with chondroitin sulfate, it can build blocks for cartilage, up-regulate chondrocyte and reduce the extent of cartilage degradation (Yu Shao *et al.*, 2004).

Fabio *et al.* (2005) synthesized D-glucosaminic acid on a multigram scale by air oxidation of D-glucosamine, catalyzed by the enzyme glucose oxidase. D-Glucosaminic acid is a component of bacterial lipopolysaccharides and a member of the 'chiral pool' and has been used as a starting material for the synthesis of various amino acids.

2.3. Peptic ulcer

Peptic ulcer disease is a very common ailment, affecting one out of eight persons in the United States and commonly seen in adults of India. It is a sore on the lining of the stomach (Fig. 2.3.) or duodenum, which is the beginning of the small intestine. The causes of peptic ulcer have gradually become clear. With this understanding have come new and better ways to treat ulcers and even cure them. These ulcers can occur in the stomach, where they are called gastric ulcers. Or they can occur in the first portion of the intestine. These are called duodenal ulcers. "Peptic Ulcer" is the term used to describe either or both of these two types of ulcers.




Fig. 2.3. Structure of Stomach

2.3.1. Anatomy and function of the Stomach

The stomach produces a very strong acid. This acid helps to digest and break down food before it enters the small intestine (duodenum). The lining of the stomach is covered by a thick protective mucous layer, which prevents the acid from injuring the wall of the stomach.

2.3.2. Causes of Ulcer

An ulcer is an open wound in the lining of the stomach or intestine, much like mouth or skin ulcers. Peptic ulcers are eventually caused by acid and pepsin, a digestive stomach enzyme.

In the end, it is the production of large amounts of acid in the stomach, passing under the term 'hyperchlorhydria', that causes the injury to the stomach or bowel lining. In a few cases, cancerous tumors in the stomach or pancreas can cause ulcers. Peptic ulcers are not caused by stress or eating spicy food, but these can make ulcers worse. However, a revolutionary and startling recent discovery is that most peptic ulcers result from a stomach infection caused by the bacteria, *Helicobacter pylori* (Fig.2.4). Use of non-steroidal anti-inflammatory drugs (NSAIDs) will also cause ulcers.

2.3.2.1. Helicobacter pylori (*H. pylori*)

Complicating our understanding of stomach ulcers is Barry Marshall and Robin Warren's discovery in 1982 that bacteria are the primary cause of stomach and duodenal ulcers excluding those caused by aspirin or arthritis drugs. This bacterium has a twisted spiral shape and infects the mucous layer lining of the stomach. This infection produces an inflammation in the stomach wall called gastritis. The body even develops a protein antibody in the blood against it. The bacterium is probably acquired from contaminated food or from a drinking glass. It is only after *H. pylori* bacteria injure the protective mucous layer of the stomach, allowing damage by stomach acid, that an ulcer develops. It takes





Fig. 2.4. Helicobacter pylori

advantage of the stomach's own mucus for protection. Any acid that does reach the bacteria is converted by *H. pylori*'s urease enzyme in the following reaction



The products of this reaction, bicarbonate and ammonia, are strong bases that further protect the bacteria because of their acid-neutralizing capability. The body's immune system responds to the presence of *H. pylori* and sends infection-fighting cells to the area. However, the neutrophils cannot reach the *Helicobacter pylori* infection because they cannot easily get through the stomach lining. Inflammation in the stomach tissue occurs as the neutrophils die and release superoxide radicals on the stomach wall, damaging tissue. The immune system sends in more nutrients to help the neutrophils, and the *H. pylori* can feed on these nutrients. It may not be the *H. pylori* itself that causes a stomach ulcer, but inflammation in the stomach lining as part of the immune response. Age is also a factor in *H. pylori* infection. In Western countries, children are unlikely to be infected. *H. pylori* infections occur in about 20% of persons below the age of 40 years, and 50% of persons above the age of 60 years.

2.3.2.2. Non Steroidal Anti-inflammatory Drugs (NSAIDs)

NSAIDs include ibuprofen, fenoprofen, aspirin, diclofenac, sulindac, diflusal, naproxen, tolmetin and many others. They can damage the mucous layer of the stomach, after which the stomach acid causes the final injury.

There are two components to NSAID-induced ulceration. First, there is a local acid effect of the dissolved drug. Most NSAIDs are weakly acidic, lipid-soluble compounds. Since the cell membranes on the stomach wall contain lipids for protection against strong acids, they offer little resistance to the lipid-soluble NSAID. The NSAID acts against the cell membrane, increasing its permeability. This results in cell swelling and death. The local acid effect of NSAIDs has been reduced by enteric-coating the drug, delaying dissolution until later in the

digestive process. However, not all NSAIDs are enteric-coated as it increases cost. In addition, enteric-coating does little more than improving the symptoms of upset stomach. Patients must be informed that enteric-coated NSAIDs are still just as likely to cause stomach ulcers as regular NSAIDs. The second and much more significant component to NSAID-induced ulceration is the systemic effect after being absorbed into the bloodstream. NSAIDs inhibit COX-1, reducing prostaglandin production. Normal COX-1 present in stomach tissue produces prostaglandins which:

- increase mucous and bicarbonate production,
- inhibit stomach acid secretion,
- increase blood flow within the stomach wall.

By acting on COX-1, NSAIDs restrict these self-protection mechanisms, allowing stomach ulcers to develop. It is primarily through this mechanism, not a local acid effect, that NSAIDs cause stomach ulcers.

So, *H. pylori* and certain drugs are the two major factors that cause ulcers. In rare cases, a patient will produce very large amounts of acid and develop ulcers. This condition is called Zollinger -Ellison syndrome. Finally, some people get ulcers for unknown reasons.

2.3.3. Other chemicals inducing Ulcer

Other than NSAIDS, certain chemicals can also cause gastrointestinal damage. In a study conducted by Silvana *et al.* (2003) ulcer was induced with indomethacin. Tan *et al.* (2002) induced ulcer by the administration of HCl-ethanol. Reserpine, an alkaloid can also be used for the induction of ulcer (Salvatore *et al.*, 2003). Jianfeng *et al.* (2002) induced ulcer in rat models by using acetic acid.

2.3.4. Symptoms

Ulcers cause gnawing, burning pain in the upper abdomen. These symptoms frequently occur several hours following a meal, after the food leaves the stomach but while acid production is still high. The burning sensation can occur during the night and be so extreme as to wake the patient. Instead of pain, some patients experience intense hunger or bloating. Antacids and milk usually give temporary relief. Other patients have no pain but have black stools, indicating that the ulcer is bleeding. Bleeding is a very serious complication of ulcers.

2.3.5. Diagnosis

A diagnosis of peptic ulcers can be suspected from the patient's medical history. However, the diagnosis should always be confirmed either by an upper intestinal endoscopy, which allows direct examination of the ulcer or by a barium x-ray of the stomach. Rarely an ulcer can be malignant. With endoscopy, a biopsy specimen can be obtained to determine if this is so.

2.3.6. Treatment

Therapy of peptic ulcer disease has undergone profound changes. There are now available very effective medications to suppress and almost eliminate the outpouring of stomach acid. These acid-suppressing drugs have been dramatically effective in relieving symptoms and allowing ulcers to heal. If an ulcer has been caused by aspirin or an arthritis drug, then no subsequent treatment is usually needed. Avoiding these latter drugs, should prevent ulcer recurrence.

The second major change in peptic ulcer disease treatment has been the discovery of the *H. pylori* infection. When this infection is treated with antibiotics, the infection, and the ulcer, do not come back. Increasingly, physicians are not just suppressing the ulcer with acid-reducing drugs, but they are also curing the underlying ulcer problem by getting rid of the bacterial infection. If this infection is not treated, the ulcers invariably recur.

There are a number of antibiotic programs available to treat *H. pylori* and cure ulcers. Working with the patient, the physician will select the best treatment program available. Antacid, H₂ receptor antagonists (ranitidine, cimetidine, nizatidine, famotidine), gastric acid pump inhibitor (omeprazole), barrier agent (sucralfate) etc. were developed to heal ulcers, but they fail to prevent the occurrence of NSAID-induced ulceration (Agrawal, 1995; Blower, 1996). These treatments are often effective in alleviating ulceration symptoms. They are also used in combination with antibiotic treatment for *H. pylori*-induced ulcers. However, the treatments listed earlier do not prevent NSAID-induced ulcers. Only misoprostol, a synthetic prostaglandin, has been shown to prevent NSAID-induced ulcers.

Some plant extracts were also found effective in the treatment of ulcer. Tan *et al.* (2002) proved the gastric cytoprotective anti-ulcer effects of the leaf methanol extract of *Ocimum suave* (Lamiaceae) in rats. Jianfeng *et al.* (2002) found that sea buckthorn (*Hippophae rhamnoides*) seed and pulp oils have both preventive and curative effects against experimental gastric ulcers in rats. Sea buckthorn (*Hippophae rhamnoides*) is a Euro-Asian wild, newly cultivated, edible berry with exceptionally high contents of nutrients and phytochemicals such as lipids, water and fat soluble vitamins, and flavonoids. Silvana *et al.* (2003) reports that *Tanacetum larvatum* has very good gastroprotective effects. Yeel *et al.* (2003) says that crude extract from *Angelica sinensis* (ASCE), which mainly consisted of polysaccharides, has a direct wound healing effect on gastric mucosa.

2.3.7. Ibuprofen

Ibuprofen is in a class of drugs called Non steroidal Antiinflammatory Drugs (NSAIDs) (Tyagi *et al.*, 2005). Ibuprofen is reported to have analgesic properties also (Polat and Karaman, 2005). Ibuprofen works by reducing hormones that cause inflammation and pain in the body. Ibuprofen inhibits cyclooxygenase (COX); thus inhibits prostaglandin synthesis. Ibuprofen appears to have gastrointestinal adverse drug reactions of all NSAIDs.

2.3.7.1. Structure

Ibuprofen is chemically 2-(p-isobutylphenyl) propionic acid (Fig. 2. 5). It has an empirical formula of $C_{13}H_{18}O_2$.

Ibuprofen like other 2-arylpropionate derivatives (including ketoprofen, flurbiprofen, naproxen etc.) contains a chiral carbon in the β -position of the propionate moiety. As such there are two possible enantiomers of ibuprofen with the potential for different biological effects and metabolism for each enantiomer.

Indeed, it was found that S-ibuprofen (sinisteribuprofen) was the active form both invitro and invivo. Logically then, there was the potential for improving the selectivity and potency of ibuprofen formulations by marketing ibuprofen as a single-enantiomer product (as occurs with naproxen, another NSAID).

Further, invivo testing, however, revealed the existence of an isomerase, which converted R-ibuprofen to the active S-enantiomer. Thus, due to the expense and futility that might be involved in marketing the single enantiomer, all ibuprofen formulations currently marketed are a racemic mixture of both enantiomers.

2.3.7.2. Properties

Ibuprofen has a molecular weight of 206.3 g. It has hepatic metabolism and has renal excretion.

Common adverse effects include nausea, dyspepsia, gastrointestinal ulceration/bleeding, raised liver enzymes, diarrhoea, headache, dizziness, salt and fluid retention, hypertension (Rozzi, 2004). Infrequent adverse effects include oesophageal ulceration, heart failure, hyperkalaemia, renal impairment, confusion, bronchospasm, rash etc. (Rozzi, 2004).




Fig. 2 .5. Structure of Ibuprofen

As with other NSAIDs, ibuprofen has been reported to be a photosensitizing agent (Castell *et al.*, 1987). Ibuprofen however has a very weak absorption spectrum which does not reach into the solar spectrum. The molecule contains only a single phenyl moiety and no bond conjugation, resulting in a very weak chromophore system. Ibuprofen, therefore, is only a very weak photosensitizing agent when compared with other members of the 2-arylpropionic acids.

2.3.7.3. Mechanism of action

Nonsteroidal anti-inflammatory drugs work by interfering with the cyclooxygenase pathway (Fig. 2.6). The normal process begins with arachidonic acid a dietary unsaturated fatty acid obtained from animal fats. This acid is converted by the enzyme cyclooxygenase to synthesize different prostaglandins. The prostaglandins go on to stimulate many other regulatory functions and reactionary responses in the body. Earlier research (Tannenbaum, 1996; Vane, 1996; Emery, 1996) have shown that there are two types of cyclooxygenase, denoted COX-1 and COX-2. Each type of cyclooxygenase lends itself to producing different types of prostaglandins.

Different mechanisms stimulate the two types of cyclooxygenase. COX-1 is stimulated continuously by normal body physiology. The COX-1 enzyme is constitutive meaning that its concentration in the body remains stable. It is present in most tissues and converts arachidonic acid into prostaglandins. These prostaglandins in turn stimulate normal body functions, such as stomach mucus production and kidney water excretion, as well as platelet formation. The location of the COX-1 enzyme dictates the function of the prostaglandins it releases (Vane, 1996). For example, COX-1 in the stomach wall produces prostaglandins that stimulate mucous production. In contrast, the COX-2 enzyme is induced. It is not normally present in cells but its expression can be increased dramatically by the action of macrophages the scavenger cells of the immune system (Tannenbaum, 1996) COX-2's most important role is in inflammation. COX-2 is involved in producing prostaglandins for an inflammatory response.




Fig. 2.6. The Cyclooxygenase Pathway

COX-1 is stimulated continually, and COX-2 is stimulated only as a part of an immune response.

2.3.7.4. Biological applications

Ibuprofen is used to reduce the fever, pain, inflammation, and stiffness caused by many conditions, such as osteoarthritis, rheumatoid arthritis, and abdominal cramps associated with menstruation. Ibuprofen is used widely in the community for the relief of headache including migraine. It is also widely marketed as an analgesic agent rather than as an anti-inflammatory and is often used for general pain conditions including those arise from various injuries such as sporting injuries, illness such as influenza, shingles, gout and post operative pain. As with other NSAIDs, ibuprofen inhibits platelet aggregation, but is not used therapeutically for this action since it is a minor and reversible effect. Litkowski *et al.* (2005) studied the analgesic effect of ibuprofen with oxycodone in combination therapy. According to Rostom *et al.* (2005), ibuprofen can cause hepatotoxicity. An earlier study (Hollenz and Labenz, 2004) reported that administration of ibuprofen could lead to gastrointestinal injury like ulcer. Long-term use of ibuprofen will lead to Nephrogenic adenoma, which is an infrequent benign lesion of the urinary system (Scelzi *et al.*, 2004). Dokmeci (2004) proved that Ibuprofen might be a promising new therapeutic avenue for the treatment of neurodegenerative diseases such as Alzheimer's disease (AD).

2.4. Hepatotoxicity

Liver is one of the most important organs in our body. It is the largest gland, which is placed behind the lower right part of ribs. Ribs help in keeping the liver from being injured. Liver is very important to human health and has a lot of functions in the body:

- Stores vitamins, sugars, fats and other nutrients from the food that we eat
- Builds chemicals that the body needs to stay healthy.

- Breaks down harmful substances, like alcohol and other toxic (poisonous) chemicals.
- Removes waste products from the blood.
- Makes sure that the body has just the right amount of other chemicals that it needs.

Many diseases can affect the liver. If anyone has one of these diseases, the liver may not work. Some of the most common diseases that affect the liver are:

1. **Viral Hepatitis:** Hepatitis is a medical term that means "inflammation (swelling) of the liver." Viruses that attack the liver cause some of the most common forms of hepatitis. Virus attack can be identified by a blood test. Three of the most common viruses that attack the liver are: Hepatitis A virus (HAV), Hepatitis B virus (HBV) and Hepatitis C virus (HCV).
2. **Cirrhosis:** Cirrhosis is a medical term that means "scarring of the liver." When one has cirrhosis, large parts of the liver are damaged. Because it has been damaged, the liver may not work as well as it should. Cirrhosis of the liver is often the result of drinking too much alcohol. Other common causes of cirrhosis include hepatitis; especially hepatitis C. Cirrhosis can be very dangerous if it is not treated properly.
3. **Liver Cancer:** Like many other body organs, liver also can get cancer. Liver cancer is a disease in which some of the cells in the liver begin to reproduce faster than they should. These cells form growths called tumors. Having hepatitis B or hepatitis C can increase the chances of getting liver cancer. Liver cancer can be deadly.
4. **Hepatotoxicity:** Toxicity of liver can be obtained by certain chemicals. Prolonged use of some drugs can also cause hepatotoxicity. Millions of people are suffering from hepatic failure. It may lead to death also. Drug induced liver injury was high in patients aged ≥ 35 years (Tasduq *et al.*,

2005). A recent study by Jasmer *et al.* (2002) found that patients >35 years old had a higher risk of grade 3 or 4 hepatotoxicity.

Fulminant hepatitis carries a very high mortality, resulting from acute hepatitis caused by virus infection, alcohol or drugs. Conventional medical therapies can rescue only about 10% of patients with fulminant hepatitis (Hiroaki *et al.*, 2003). Although liver transplantation has improved their mortality, about 40% of these patients die while waiting for liver transplantation (Shakile *et al.*, 2000)

2.4.1. Chemicals inducing hepatotoxicity

Administration of some chemicals can induce hepatotoxicity. Sodium diethyl dithiocarbamate, diclofenac and ketoconazol are three important chemotherapeutic agents that are commonly associated with hepatotoxicity (Amr and Alaa, 2005). One of the most hazardous pollutants called hexachlorobenzene can also cause hepatotoxicity (Billi de Catabbi *et al.*, 2005). Kaufmann (2005) reported that treatment with some drugs causes liver injury. A lot of drugs are devastated from the market due to their induction of hepatotoxicity. Factors affecting susceptibility to drug-induced injury include age, sex, concomitant use of other drugs, and genetic polymorphism in metabolic pathways involved in activation or disposition of therapeutic drugs (Maddrey, 2005). Older persons who are receiving more drugs simultaneously are more susceptible to liver injuries. Guzman *et al.* (2005) studied about the hepatotoxic effect of paroxetine. Galactosamine is capable of inducing wild hepatic failure (Abul *et al.*, 2005). It resembles viral hepatitis. In a study conducted by Senthilkumar and Nalini (2004) hepatotoxicity was induced by the intake of alcohol. Tomiyama *et al.* (2004) studied the effect of DDT in inducing toxicity to liver. Doxorubicin is an anthracycline antibiotic, popularly used in tumour therapy. Administration of doxorubicin was proved to cause hepatotoxicity (Kalender *et al.*, 2005).

Statins are among the most widely prescribed medications in the western world (Eidelman *et al.*, 2002). Their benefit in the primary and secondary prevention of

cardiovascular disease is unequivocal. According to Naga Chalasani (2005), intake of statin will lead to severe liver toxicity. As per the reports of Ishiyama *et al.* (1990) both hepatotoxicity and oxidative stress have been reported in rats following a single dose treatment with diethyl dithiocarbamate. In a study of Crupi *et al.* (2001), liver damage was induced by the use of carbon tetra chloride. Mitra *et al.* (1998) used paracetamol to induce hepatotoxicity in rats.

Herbal drugs have become increasingly popular among patients and physicians (Schuppan *et al.*, 1999) because of the perception that drugs derived from plants are normally safe. However, numerous reports have come forward, which showed hepatic adverse effects induced by plants containing pyrrolizidine alkaloids such as comfrey (Stickel and Seitz, 2000), preparations produced from germander (Larrey *et al.*, 1992), those containing chaparral extracts (Sheikh *et al.*, 1997), and drugs made of Chinese herbs (Yoshida *et al.*, 1996). These plants may contain alkaloids that are biotransformed to toxins. With most of the herbs, hepatotoxic reactions were reversible after withdrawal. Felix *et al.* (2003) studied the hepatotoxic effects of Kava (*Piper methysticum rhizoma*), which is used as a sleeping aid and for the treatment of anxiety disorders and depression.

2.4.2. Mechanism of drug induced liver injury

Mechanisms of drug-induced liver injury are many and varied. With many drugs, intermediary products produced during metabolism are highly reactive and toxic. In these situations, the balance between the rate of production of the metabolite and the effectiveness of the drug may determine whether or not hepatic injury occurs (Maddrey, 2005). Some earlier studies reported that diclofenac induced liver damage through various mechanisms such as mitochondrial permeability transition (Masubuchi *et al.*, 2002), activation of cytochrome P450 (Cantoni *et al.*, 2003) and generation of reactive oxygen species (ROS) (Cantoni *et al.*, 2003; Gomez-Lechon *et al.*, 2003). Ketoconazol has also been reported to mediate hepatotoxicity through chelating GSH with active toxic metabolites (Rodriguez and Buckholz, 2003).

Hepatic DNA fragmentation has been reported in many hepatotoxicity models following treatments with drugs such as acetaminophen (Ray *et al.*, 1992), carbon tetrachloride (Shi *et al.*, 1998) and diquat (Gupta *et al.*, 200). Amr and Alaa (2005) reported that in hepatotoxicity models induced by diethyl dithiocarbamate, diclofenac and ketoconazole, levels of serum ALT and AST have been significantly increased. These three toxic drugs decreased the levels of GSH and SOD. Diethyl dithiocarbamate induced hepatotoxicity in rats was associated with reactive oxygen species (ROS) and drug metabolism (Ishiyama *et al.*, 1990).

Other mechanisms of hepatotoxicity are summarized in the following Table 2 (Dominique, 2000)

Liver injuries	Mechanisms
Acute hepatitis	Metabolite-mediated toxicity Metabolite-mediated immunoallergy and/or autoimmunity
Acute cholestasis	Inhibition of biliary secretion
Macrovesicular steatosis	Decreased secretion of lipoproteins
Microvesicular steatosis	Inhibition of fatty acid mitochondrial beta-oxidation
Phospholipidosis	Inhibition of lysosomal phospholipases
Chronic hepatitis	Metabolite-mediated immune reaction
Vanishing bile duct syndrome	Autoimmune destruction of small bile ducts. Abnormal multidrug resistance protein system
Sclerosing cholangitis	Biliary ischemia caused by arterial lesions
Veno-occlusive disease	Metabolite-mediated endothelial lesions
Persinusoidal fibrosis	Activation of Ito cells

Table 2. Mechanisms of Hepatotoxicity

2.4.3. Treatment

A lot of treatment methods are found popular in the treatment of hepatotoxicity. The stem bark of the Betulaceae plant *Alnus japonica*, which is indigenous to Korea, has been used as a popular folk medicine for hepatitis induced by acetaminophen (Kim *et al.*, 2004). Tazduq *et al.* (2005) reported that Silymarin can reduce hepatic toxicity induced by rifampicin, isoniazid and pyrazinamide. Betaine supplementation blocked or significantly attenuated the induction of hepatotoxicity by alpha-naphthylisothiocyanate (ANIT) (Kim *et al.*, 2005). Jigrine was found to be effective against galactosamine induced hepatitis (Abul, 2005). Jigrine, a polypharmaceutical herbal hepatoprotective formulation containing aqueous extracts of 14 medicinal plants, is used in Indian system of medicine (Unani). Kalender *et al.* (2005) reported that Vitamin E and Catechin could reduce the hepatotoxic effects of antitumour drug doxorubicin. Porchezhian and Ansari (2005) studied the heparoprotective effect of *Abutilon indicum* against carbon tetrachloride and paracetamol-induced hepatotoxicity.

2.5. Antitubercular drugs

2.5.1. Tuberculosis

Tuberculosis (TB) is an infection caused by two species of *Mycobacteria*, "*Mycobacterium Tuberculosis* and *Mycobacterium Bovis*". Though it can cause disease involving every organ system in the body, it commonly affects the lungs. The disease was scraping the human kind even in the Neolithic period and till the early 20th century. During the 19th century, up to 25 per cent of deaths in Europe were caused by this disease. The death toll began to fall as living standards improved at the start of the 20th century and from the 1940s, effective medicines were developed. However, there are now more people in the world with TB than there were in 1950, and three million individuals may die this year from this disease - mainly in less developed countries. The disease is more common in areas of the world where poverty, malnutrition, poor general health and social disruption are present. Currently, around 1.7 billion people worldwide, a third of the world's population, are infected by *Mycobacterium tuberculosis* and 3 million

deaths a year is attributable to tuberculosis. Rapid spread of TB in humans is attributed to crowded living conditions that favour airborne transmission. There was a steady decline in the incidence of tuberculosis, especially in the developed countries till early 1980s; but, since then the trend has reversed and an increasing number of cases have been reported.

2.5.1.2. Causes and Pathogenesis

The term tubercle bacillus refers to two species, "*Mycobacterium Tuberculosis* and *Mycobacterium Bovis*". Other species are classified under a typical Mycobacterial Pathogens. Humans are the only reservoir for *Mycobacterium tuberculosis*. *Mycobacterium Bovis* was transmitted by contaminated milk once, but is no longer so. Though skin infection by inoculation is seen in pathologists and laboratory personnels, almost all infections are due to airborne transmission by inhalation of droplet nuclei. In general, 3 - 4% of infected individuals will develop active disease during the first year after exposure and a total of 5-15% thereafter. The likelihood of developing active disease varies with the intensity and duration of exposure. Malnutrition, alcoholism, renal failure and uncontrolled diabetes favour the progression of infection to active disease. HIV infection is the strongest risk factor today. The tubercle bacilli, once inhaled, reach the terminal air spaces in the lungs, escaping from the host defense mechanism. They multiply locally but are controlled and retained in the lungs by the body's white blood cells. Sometimes infected white cells carry the bacteria to the lymph nodes and from there to the blood stream. Seeding and unchecked proliferation in other organs can cause disease manifestations elsewhere in the body as well. Pulmonary tuberculosis is usually seen in the upper portions of the lung.

2.5.1.3. Symptoms and Signs

The clinical picture depends on the involved organ. Early tuberculosis of the lung is asymptomatic and may be discovered on a chest x-ray by chance. In case of tuberculosis of the lung, the patient will complain of fever, night sweats, fatigue,

cough, sputum production that is sometimes mixed with blood, weight loss and loss of appetite.

Tuberculosis can also appear as swelling of the glands in the neck with or without fever (lymph node TB), back pain, deformity of the spine and weakness in the lower limbs (TB of the spine), fever, headache, vomiting and drowsiness (TB meningitis), joint pain and swelling (TB arthritis), genitourinary symptoms like flank pain and infertility (genitourinary TB).

2.5.1.4. Treatment and Prognosis

With the advent of 'Combination Chemotherapy', successful treatment of tuberculosis is a reality, but problems of drug resistance, selection of an inappropriate regimen and non-compliance hinder effective therapy. WHO recommends a six-month short course therapy consisting of four drugs - rifampicin, isoniazid, ethambutol and pyrazinamide, for the first two months and two drugs - rifampicin and isoniazid, for the next four months.

2.5.2. Isoniazid

Isoniazid, an antibiotic, is used to treat and prevent tuberculosis since, it prevents Tuberculosis bacteria from multiplying in the body. Isoniazid consumption is associated with a lot of side effects like live difficulty breathing, closing of the throat, swelling of the lips, tongue, or face, unusual weakness or fatigue, abdominal pain, yellow skin or eyes, dark urine etc. Isoniazid toxicity is associated with a high mortality rate (Wason, 1981). If isoniazid is taken acutely, as little as 1.5 g can cause toxicity. Doses larger than 30 mg per kg often produce seizures. Ingestion of the drug in amounts greater than 80 to 150 mg per kg can rapidly lead to death (Shannon *et al.*, 1990). A 1-2 % risk of severe and potentially fatal hepatotoxicity (predominantly hepatic necrosis) associated with the use of isoniazid is problematic in the prophylaxis and treatment of tuberculosis. In addition, daily isoniazid administration is associated with mild elevations of liver enzyme activities in plasma up to 20 % of patients (Troy *et al.*, 1998).




Fig. 2.7. Structure of Isoniazid

2.5.2.1. Structure

Chemically isoniazid is Isonicotinic acid hydrazide. Its molecular formula is $C_6H_7N_3O$. The structure of isoniazid is depicted in Fig. 2. 7.

2.5.2.2. Properties

Isoniazid (INH) has a molecular mass of 137.14g. Isoniazid is a white crystalline powder in occurrence. The substance decomposes on heating and on burning produces toxic fumes including nitrogen oxides. Melting point of Isoniazid comes around 170-173°C. INH is a colourless, odourless, white crystalline powder slowly affected by exposure to air and to light. 1 g isoniazid is soluble in 8 g water. A 10% solution has a pH of 6.0 to 8.0. Isoniazid can induce hepatotoxicity when administered for a long time (Saraswathy and Shyamala Devi, 1999). Hepatitis is due to a toxic metabolite of monoacetyl hydrazine, which binds covalently to liver proteins (Black *et al.*, 1975). In some patients, an allergic mechanism has also been proposed: acetylation of hepatic macromolecules by acetyl hydrazine may lead to the release of antigenic macromolecules, which induce the formation of antibodies directed against the liver (Davies, 1981). Isoniazid interacts with dietary supplements. Isoniazid can interfere with the activity of vitamin B₆ (Goldman and Braman, 1972). Vitamin B₆ supplementation is recommended, especially in people with poor nutritional status, to prevent development of isoniazid-induced peripheral neuritis (inflamed nerves) (Mandell and Petri Jr., 1996). Isoniazid may kill friendly bacteria, in the large intestine, that produce vitamin K. (Holt, 1998). Isoniazid may interfere with the activity of other nutrients, including vitamin B₃ (niacin), vitamin B₁₂, vitamin D, and vitamin E, folic acid, calcium, and magnesium (Werbach, 1997; Holt, 1998).

2.5.2.3. Metabolic pathway of Isoniazid

The major route of isoniazid metabolism is hepatic acetylation by N-acetyl transferase, which produces acetylisoniazid. The rate of acetylation is




Fig. 2.8. Metabolic pathway of Isoniazid

genetically determined. Acetyisoniazid is further hydrolysed to isonicotinic acid and acetylhydrazine, both of which are excreted in the urine. Isonicotinic acid is conjugated with glycine. Acetylhydrazine is further metabolized to deacetylhydrazine and may be converted by the hepatic microsomal enzymes to the reactive metabolite (presumed to be hydrazine) that are thought responsible for INH-induced hepatotoxicity. Acid labile hydrazones of isoniazid are formed with α -ketoglutarate and pyruvate, but since these do not appear to any extent in the blood, they are thought to be produced in the pointed to cytolytic liver injury in rat studies, as a result of free radical generation, isoniazid bladder (Boxenbaum and Riegelman, 1974). Fig. 2. 8 shows the metabolic pathway of isoniazid.

2.5.2.4. Physiological changes

Some earlier studies have pointed to the cytolytic liver injury in rat studies, as a result free radical generation when isoniazid was used. These cytotoxic materials had been shown to come from lipid peroxidation and the suppression of the antioxidant system (Saraswathy *et al.*, 1998). Studies related to the suppression of the antioxidant system in antituberculosis drugs with rats have also been reported (Skakun and Slivka, 1992). Georgieva *et al.* (2004) reported that lipid peroxidation was found high in isoniazid treated animals. According to Tasduq *et al.* (2005) Alanine aminotransferase (ALT) and Aspartate aminotransferase (AST) and Alkaline phosphatase (ALP) levels were increased when isoniazid was orally administered in rats for induced hepatotoxicity.

2.5.3. Rifampicin

Rifampicin is an antibiotic drug of the rifamycin group. It is typically used to treat mycobacterium infection, including tuberculosis and leprosy and also has a role in the treatment of methicillin-resistant *Staphylococcus aureus* (MRSA) in combination with fusidic acid.

Rifampicin is a semisynthetic derivative of rifamycin antibiotics which are produced by the fermentation of a strain of *Streptomyces mediterranei*, a species which was first isolated in Italy in 1957 from a soil sample collected in France. The fermentation produces rifamycin B. Rifamycin B is transformed by a series of reactions into 3-formylrifamycin SV, which in turn is condensed with 1-amino-4-methylpiperazine in peroxide-free tetrahydrofuran to give rifampicin.

Rifampicin inhibits DNA-dependent RNA polymerase in bacterial cells by binding its beta subunit, thus preventing transcription of messenger RNA and subsequent translation to proteins.

2.5.3.1. Structure

Chemically rifampicin is 3-[[[(4-methyl-1-piperazinyl)imino]methyl]] rifamycin. Structure of rifampicin is given in Fig. 2. 9.

2.5.3.2. Properties

Molecular formula is $C_{43}H_{58}N_4O_{12}$. Molecular weight is 822.96g. Its colour is red to orange. Powder in nature. Odourless. Melting point is 138 to 188 °C. A 1% suspension in water has pH 4.5 to 6.5. Rifampicin is very soluble in water. Solubility in aqueous solution is increased at acidic pH. Freely soluble in chloroform.

Patients receiving antitubercular drugs frequently develop acute or chronic hepatitis (Hussain, 2003). Prolonged use of Rifampicin may affect gastrointestinal system also (Wada, 1998). When rifampicin is administered the levels of ALT was increased. Bilirubin serum levels were also found increasing (Lenaerts, 2005).




Fig. 2.9. Structure of rifampicin

This is an indication of drug-induced hepatotoxicity. Dawe *et al* (2004) reported acute renal failure associated with rifampicin administration.

2.5.3.3. Physiological changes

An earlier reported study (Chi *et al.*, 2003) proved that intake of rifampicin increased the level of alanine aminotransferase (ALT). In a study conducted by Prabakan *et al.* (2000) The level of liver mitochondrial protein and the activities of isocitrate dehydrogenase, α -ketoglutarate dehydrogenase, succinate dehydrogenase, malate dehydrogenase, NADH dehydrogenase and cytochrome c oxidase were found significantly decreased in rifampicin and isoniazid intoxicated rats.

3. MATERIALS AND METHODS

3.1. Drugs and chemicals

Ibuprofen was obtained from M/s. Cipla Pharmaceuticals Ltd., Mumbai; Bovine serum albumin, tetraethoxy propane, epinephrine and D-galactosamine from M/s Sigma Chemical Company, St. Louis, MO, USA. Glucosamine used in this study was prepared from chitin using a method developed in our laboratory.

Isoniazid and rifampicin were obtained by M/s. Lupin Pharmaceuticals Ltd., Mumbai. Bovine serum albumin, tetraethoxy propane and cholesterol were from M/s Sigma Chemical Company, St. Louis, MO, USA. Chitosan (MW 750000 Da; viscosity 8 cps; deacetylation rate 85-87%; purity 98.6%) used in this study was prepared from chitin in our laboratory (Madhavan, 1992).

All other chemicals used were of analytical grade.

3.2. Animals

Wistar strain male albino rats, weighing 120-150g, were selected for the study. The animals were housed in polyurethane cages under hygienic conditions and maintained at normal room temperature. The animals were allowed food and water *ad libitum*. The experiment was carried out according to the guidelines of the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA), New Delhi, India, and approved by the Institutional Animal Ethics Committee (IAEC).

3.2.1. Induction of ulcer (Santhosh *et al.*, 2005)

The ulcer was induced in experimental rats by oral administration of ibuprofen (50mg/kg, p.o.) twice in a day at an interval of 12 h. The animals were deprived of food for 24h prior to ulcer induction.

3.2.1.2. Experimental protocol

The experimental animals were divided into four groups of six rats each.

Group I, normal control rats received only the standard diet.

Group II, ulcer was induced by oral administration of ibuprofen (50mg/kg body weight/each rat, p.o) twice in a day at an interval of 12 h.

Group III, normal rats were treated with chitosan (100 mg/kg body weight/each rat/day, p.o for 20 days).

Group IV, pretreated with chitosan (100mg/kg body weight/each rat/day, p.o for 20 days) before induction of ulcer as described for Group II.

Group V, normal rats were treated with chitosan (100 mg/kg body weight/each rat/day, p.o for 20 days).

Group VI, pretreated with glucosamine (100mg/kg body weight/each rat /day, p.o for 20 days) before induction of ulcer as described for Group II.

At the end of the experiment, all six groups underwent surgery according to Takeuchi *et al.* (1976) and gastric juice was collected for 4 h. Rats were then killed with over dose of chloroform and the stomach was removed after the esophagus had been clamped. The gastric juice was centrifuged and the volume was noted. The stomach was inflated with normal saline and then incised through the greater curvature and examined for the number of lesions. The total acidity was determined by titration with 0.02 N NaOH with phenolphthalein used as indicator. One portion of the stomach tissue was kept in 10% formalin-saline for histopathological observations. The mucosal tissue was scraped from the stomach and the gastric mucosa was homogenized in ice-cold 0.1 M Tris-HCl, pH 7.2 and centrifuged. The supernatant was used for further biochemical analyses.

3.2.2. Induction of Hepatotoxicity (Santhosh *et al.*, 2006)

Hepatotoxicity was induced by the administration of rifampicin and isoniazid (200mg each /kg body weight/each rat/day).

3.2.2.1. Experimental Design

The experimental animals were divided into four groups of six rats each. Rats in **Group I** (normal control) were given only the standard diet. In **Group II**, normal rats were treated with chitosan (100 mg/kg body weight/day, intragastric intubation for 30 days). In **Group III**, normal rats were treated with glucosamine (100 mg/kg body weight/day, intragastric intubation for 30 days). In **Group IV**, hepatotoxicity was induced by oral administration of isoniazid and rifampicin (200mg each /kg body weight/day) for a period of 30 days. **Group V** animals were co-administered with chitosan (100mg/kg body weight/day, intragastric intubation for 30 days) along with antitubercular drugs described for Group IV. **Group VI** animals were co-administered with chitosan (100mg/kg body weight/day, intragastric intubation for 30 days) along with antitubercular drugs described for Group IV.

At the end of the experiment, rats were killed and blood was collected without any anticoagulant. The serum separated was used for different analyses. The liver and kidney tissues were excised immediately and washed with ice-cold saline. One portion of each tissue was fixed in 10% formalin-saline for histopathological observations.

Accurately weighed liver and kidney tissues were homogenized with 0.1M Tris HCl pH 7.4 and the homogenate was used for the estimation of enzyme activities and cellular macromolecules

3.3.1. Assay of alanine aminotransferase (EC 2.6.1.2)

The activity of alanine aminotransferase (ALT) was determined by the method of Mohur and Cook (1957).

Reagents

1. Buffered substrate solution (0.1 M phosphate buffer, pH 7.4, 0.2 M DL-alanine, 2.0 mM 2-oxoglutarate): 1.5g dipotassium hydrogen phosphate, 0.2g potassium dihydrogen phosphate, 0.03g 2-oxoglutaric acid and 1.78g DL-alanine were dissolved in distilled water. The pH was adjusted to 7.4 with 1N NaOH and made up to 100 ml.
2. 20mg 2,4 dinitrophenyl hydrazine (DNPH) in 100 ml of hot 1N hydrochloric acid.
3. 0.4N Sodium hydroxide.
4. Standard pyruvic acid: 12.5 mg of sodium pyruvate was dissolved in 10 ml of distilled water. 10 ml of this was diluted to 100 ml with distilled water and this was prepared freshly for the calibration curve.

Procedure

To 1.0 ml of the buffered substrate, 0.1 ml sample was added and incubated at 37°C for 30 min. The reaction was arrested by adding 1.0 ml of DNPH and left aside for 20 min at room temperature. Color developed by the addition of 10 ml of 0.4N NaOH was read at 540nm in a Shimadzu UV spectrophotometer against the reagent blank.

The enzyme activity was expressed as μ moles of pyruvate liberated /hr/l (serum); μ moles of pyruvate liberated /mg protein (liver)

3.3.2. Assay of aspartate aminotransferase (EC 2.6.1.1)

The aspartate aminotransferase (AST) activity was assayed by the method of Mohur and Cook (1957).

Reagents

1. Phosphate buffer: 0.15M, pH 7.5
2. Substrate: 300mg of L-aspartic acid and 50mg of α -ketoglutaric acid were dissolved in 20-30 ml of the phosphate buffer and added 10% sodium hydroxide to bring the pH to 7.5 and was made up to 100 ml with phosphate buffer.
3. 2,4-dinitro phenyl hydrazine (DNPH) reagent: Dissolved 200mg of DNPH in 85 ml of concentrated hydrochloric acid and made up to a litre with distilled water.
4. 0.4 N sodium hydroxide.
5. Standard pyruvic acid: 12.5mg of sodium pyruvate was dissolved in 10 ml of distilled water. 10 ml of this was diluted to 100 ml with distilled water and this was prepared freshly for the calibration curve.

Procedure

To 1.0 ml of the buffered substrate, 0.1 ml of the sample was added and incubated for one hour at 37°C. Then 1.0 ml of DNPH reagent was added and left for 20 min. At the end of the incubation, 10 ml of 0.4 N NaOH was added and the colour developed was estimated by reading OD at 540nm in a Shimadzu UV spectrophotometer after 10 min. The standards were also treated similarly.

The enzyme activity was expressed as μ moles of pyruvate liberated /hr/l (serum); μ moles of pyruvate liberated /mg protein (liver)

3.3.3. Assay of lactate dehydrogenase (EC 1.1.1.27)

The lactate dehydrogenase (LDH) activity was assayed according to the method of King (1965) with slight modification. The amount of pyruvate formed in the forward reaction was measured colorimetrically.

Reagents

1. 0.1M glycine buffer: 7.5 g of glycine and 5.88 g of sodium chloride were dissolved in one liter of distilled water.
2. Buffered substrate: 2.76 g of lithium lactate was dissolved in 125 ml of glycine buffer containing 75 ml of 0.1 N sodium hydroxide to adjust the pH to 10. This was prepared just prior to use.
3. 0.4 N Sodium hydroxide.
4. 5.0 mg of NAD^+ was dissolved in 1.0 ml of distilled water. This was prepared just before use.
5. 2,4-dinitrophenyl hydrazine (DNPH): 200 mg of DNPH was dissolved in one litre of 1 N HCl.
6. Standard pyruvate solution: 12.5 mg of sodium pyruvate was dissolved in 100 ml of buffered substrate.

Procedure

To 1.0 ml of the buffered substrate, 0.2 ml of the sample was added and the tubes were incubated at 37°C for 15 min. After adding 0.2 ml of NAD^+ solution, the incubation was continued for 30 minutes and then 1.0 ml of DNPH reagent was added. And the tubes were incubated at 37°C for 15 min. Then 7.0 ml of 0.4 N NaOH was added and the colour developed was measured at 540 nm in a

Shimadzu UV spectrophotometer against the reagent blank. Suitable aliquots of the standards were also treated in the same manner.

The enzyme activity was expressed as μ moles of pyruvate liberated /hr/l (serum); μ moles of pyruvate liberated /mg protein (liver)

3.3.4. Assay of Alkaline Phosphatase (EC 3.1.3.1)

Alkaline phosphatase was assayed by the method King (1965) using disodium phenyl phosphate as the substrate.

Reagents

1. M carbonate-bicarbonate buffer, pH 10.0
2. Substrate: 0.01M disodium phenyl phosphate solution.
3. Folin's phenol reagent: Into a 1500 ml round-bottomed flask, 100 g of sodium molybdate, 700 ml water, 50 ml of 85% O-phosphoric acid and 100 ml of concentrated hydrochloric acid were added. The mixture was refluxed gently for 10 hours. Then, 150 g of lithium sulfate, 50 ml of water and few drops of bromine were added. The mixture was boiled for 15 minutes to remove excess bromine. The contents were cooled and diluted to one litre and filtered. This was diluted 1:2 with double distilled water before use.
4. 15% sodium carbonate
5. 0.1 M magnesium chloride
6. Standard phenol solution: A solution of distilled crystalline phenol in water, containing 5.0 μ g in 0.1 ml was prepared.

Procedure

The incubation mixture contained the following components in a final volume of 3.0 ml. 1.5 ml of carbonate-bicarbonate buffer, 1.0 ml of substrate and 0.1 ml of magnesium chloride and requisite amount of the enzyme source (0.2 ml serum). The reaction mixture was incubated at 37°C for 15 minutes. The reaction was terminated by the addition of 1.0 ml of Folin's phenol reagent. If turbidity appeared, the tubes were centrifuged. Controls without enzyme sources were also incubated and the enzyme source was added after the addition of Folin's – phenol reagent. The 1.0 ml of 15% sodium carbonate solution was added and incubated for a further 10 minutes at 37°C. The blue colour developed was read at 640 nm against a blank. The standards were also treated similarly.

The activity of the enzyme is expressed as μ moles of phenol liberated/hr/l (serum); μ moles of phenol liberated/mg protein (liver)

3.3.5. Assay of Acid Phosphatase(EC 3.1.3.2)

The procedure adopted for the assay of acid phosphatase was the same as described above for alkaline phosphatase (King, 1965), excepting that a citrate buffer (0.1M,pH 4.9) was used and the magnesium ions were omitted from the incubation mixture. The enzyme activity is expressed as μ moles of phenol liberated/hr/l (serum); μ moles of phenol liberated/mg protein (liver)

3.3.6. Estimation of Glucose

Glucose was estimated by the method of Sasaki *et al.* (1972), using o-toluidine reagent.

Reagents

1. Trichloro acetic acid (TCA): 10%

2. O-Toluidine reagent : 12.5 g of thiourea and 12.0 g of boric acid were dissolved in 50 ml of distilled water heating over a mild flame. Exactly 75 ml of redistilled o-toluidine and 375 ml of acetic acid (A.R.) were mixed with thiourea-boric acid mixture and the total volume was made up to 500 ml with distilled water. The reagent was left in a refrigerator overnight and filtered.
3. Standard glucose solution : 10mg of pure glucose was dissolved in 100 ml of 0.2% boric acid in water.

Procedure

Mix 0.1 ml of blood with 1.9 ml of TCA solution to precipitate protein and then centrifuge. 1.0 ml of the supernatant was mixed with 4.0 ml of o-toluidine reagent was kept in a boiling water bath for 15 minutes. The green colour developed was read at 600 nm in a Spectronic Genesys spectrophotometer. A set of standard glucose solutions were also treated similarly. Blood sugar levels were arrived at by comparison with the standard curve.

The values are expressed as mg per dl.

3.3.7. Estimation of protein

The protein content was estimated by the method of Lowry *et al.* (1951)

Reagents

1. Alkaline copper reagent

Solution A: 2% sodium carbonate in 0.1N sodium hydroxide solution.

Solution B: 0.5 copper sulphate in water

Solution C: 1% sodium potassium tartrate in water. 50ml of solution A was mixed with 0.5ml of solution B and 1ml of solution C just before use.

2. Folin's phenol reagent: Into a 1500ml round bottom flask, 100g sodium molybdate, 700ml water, 50ml 85% O-phosphoric acid and 100ml concentrated hydrochloric acid were added. The mixture was refluxed gently for 10 hours. Then 150g of lithium sulfate, 50ml of water and few drops of bromine were added. The mixture was boiled for 15 minutes to remove excess bromine. The contents were cooled and diluted to one litre and filtered. This was diluted 1:2 with double distilled water before use.
3. Standard bovine serum albumin (BSA): Dissolved 100mg of BSA in 100ml water in a standard flask. Small quantities of alkali could be added to the effect complete dissolution of BSA. 10ml of the stock was diluted to 100ml to get a working standard containing 100µg/ml.

Procedure

Pipetted out 0.01ml of serum, 0.1 ml of tissue homogenate and standard BSA in the range of 20-100µg into test tubes and the total volume was made upto 1.0 ml with water. The blank contained 1.0 ml of water. Exactly 4.5ml of alkaline copper reagent was added to all the tubes and left at room temperature for 10 minutes after which was added 0.5 ml Folin's phenol reagent. The blue colour developed was read after 20 minutes at 640 nm against the reagent blank, in a Shimadzu UV spectrophotometer.

Protein values are expressed as g per dl (serum); mg per g (tissue).

3.3.8. Electrophoretic separation of serum and plasma proteins

Plasma and Serum proteins were separated by SDS-PAGE technique as described by Laemmli (1970).

It is based on the principle that, in the presence of 10% SDS and 2-mercaptoethanol, proteins dissociate into their sub units and bind large quantities of the detergent which mask the charge of the proteins and giving a constant charge to mass ratio. So that the proteins move according to their molecular

weight in an electric field. In this discontinuous buffer system, the separating (resolving gels) and stacking gels are made up in the electrode buffer, Tris-glycine. During electrophoresis, the leading ion is chloride while the trailing ion is glycine. In this experiment, 7.5% gel concentration is used for the effective separation.

Reagents

1. Tris-HCl : 0.5M, pH 6.8
2. Tris HCl : 1.5M, pH 8.8
3. SDS : 10%
4. Acrylamide/Bis : 30%T, 2.67 % C.
5. Sample buffer :

Distilled water: 3.8 ml

Tris-HCl 0.5 M, pH 6.8: 1 ml

Glycerol: 0.8 ml

10% SDS: 1.6 ml

2-mercapto ethanol: 0.4 ml.

1% Bromophenol blue: 0.4 ml.

6. Electrode Buffer :

Tris base : 9 g

Glycine : 43.2 g

SDS : 3 g

These reagents are dissolved in 600 ml of distilled water.

Working solution: Dilute 100 ml from stock to 500 ml with distilled water.

7. Separating gel (7.5%)

Distilled water: 4.85 ml

Tris-HCl 1.5M: 2.5 ml

10%SDS: 100 μ l

Acrylamide: 2.5 ml

APS 10%: 50 μ l

TEMED: 5 μ l

8. Stacking Gel (4%)

Distilled water: 6.1 ml

Tris-HCl 0.5M: 2.5 ml

10% SDS: 100 μ l

Acrylamide: 1.33 ml

APS 10%: 50 μ l

TEMED: 10 μ l

9. Ammonium per sulphate (APS): 10%

Procedure

Taken 0.1 ml of the suitably diluted plasma into a micro centrifuge tube and added 0.1 ml of the sample buffer, heated in a boiling water bath for 4-min, cooled and kept at 4^o C in a refrigerator.

The separating gel was prepared without TEMED and APS. Evacuated for 15 min to remove air bubbles. Added TEMED and APS with intermittent shaking after each addition immediately transferred the solution to the apparatus. Added a little water on the top of the gel to level it. Kept for 45 min. Prepared stacking gel in the same way. Kept the comb over the apparatus, tilted it to 45^o, poured the gel slowly, and pressed the comb slowly and evenly. Kept for 45 min. Marked the wells. After removing the comb, the whole apparatus was transferred to the sandwich clamp assembly in to the inner cooling core. Rinsed the apparatus and wells with electrode buffer and filled the inner chamber of the apparatus completely and the outer chamber to the optimum level. Injected 10 µl of the sample into the wells. The electrode lid was placed at proper position and connections were established. The power of 200V was supplied. Electrophoresis was carried out for 45 min approximately until the dye reaches the bottom. Subsequently, the gel was removed and is placed in a big petridish containing the stain, Coomassive blue. Kept for 30 min, and transferred the gel into 7% acetic acid for destaining. 7% acetic acid was changed intermittently till the gel got completely destained.

3.3.9. Extraction of Glycoconjugates

500mg of the tissue was weighed and the lipids were extracted using chloroform-methanol mixture by homogenization in a Potter-Elvehjem homogenizer with a Teflon pestle. The extraction was repeated thrice with fresh aliquots of the solvent mixture. The lipid extract was filtered through a fat free Whatmann No.41

filter paper into a separating funnel. The defatted tissue in the filter paper was dried and used for the estimation of hexose, hexosamine and sialic acid.

3.3.10. Estimation of Hexose

Hexose was estimated by the method of Niebes (1972).

Reagents

1. Sulphuric acid: water mixture (3:2, v/v).
2. 800mg of orcinol dissolved in 50 ml of 1 N H₂SO₄.
3. Orcinol-sulphuric acid mixture: 1.0 ml of reagent (2) was mixed with 7.5 ml of reagent (1). This mixture was prepared fresh at the time of assay.
4. Standard hexose: Equal quantities of galactose and mannose were dissolved in water to give a concentration of 100 µg/ml.

Procedure

An aliquot of the delipidised sample was treated with 1.0 ml of 0.1 N NaOH. Blank contained 1.0 ml of 0.1N NaOH. The tubes were cooled by placing in an ice-bath and 8.5 ml of Orcinol-sulphuric acid mixture was added slowly and mixed well. The tubes were stoppered, incubated at 80°C for 15 min in a water bath. Cooled and the color was allowed to develop in the dark for 25 min. The intensity was measured at 540 nm. Standard solutions containing 0.025 to 0.1 mg were treated similarly and hexose concentration was estimated.

The concentration of hexose was expressed as mg/g (tissue).

3.3.11. Estimation of hexosamine

Hexosamine was estimated by the method of Wagner (1979).

Reagents

1. Acetyl acetone reagent: 3.5% acetyl acetone in 1N Trisodium phosphate containing 0.5 N potassium tetra borate (98: 2 V/V).
2. Ehrlich's reagent: 3.2g of P-dimethyl aminobenzaldehyde was dissolved in 30 ml of 1 N HCl and diluted to 210 ml with isopropanol.
3. Standard Hexosamine: Galactosamine hydrochloride solution containing 10 mg/100 ml was prepared.

Procedure

An aliquot of the delipidised sample was hydrolyzed with 3 N HCl in a boiling water bath for 4 hrs and neutralized. 0.8 ml of the neutral hydrolysate was mixed with 0.6 ml of acetyl acetone reagent. The mixture was heated in a boiling water bath for 30 min, cooled and 2.0 ml of Ehrlich's reagent was added. The contents of the tubes were mixed and the absorbance was measured at 535 nm. Standard hexosamine solution containing 20 μ g to 80 μ g was used for the preparation of standard curve.

Hexosamine was expressed as mg/g (tissue).

3.3.12. Estimation of Sialic Acid

Sialic acid was estimated by the method of Warren (145)

Reagents

1. Sodium meta arsenite: A 10% solution was prepared in 0.5 M sodium sulphate in 0.1N H₂SO₄.
2. Sodium meta periodate : 0.2 M solution in 9 M phosphoric acid.

3. Thiobarbituric acid reagent : 0.6% solution was prepared in 0.5 M sodium sulphate.
4. Acidified butanol: 5 ml of concentration hydrochloric acid in 95 ml of n-butanol.
5. Standard sialic acid : 10mg of N-acetyl neuraminic acid was dissolved in 100 ml of water.

Procedure

An aliquot of the delipidised sample was hydrolysed with 0.1N H₂SO₄. at 80⁰C for 1 hour. 0.2 ml of the hydrolysate was mixed with 0.1 ml of metaperiodate and the solution was kept at room temperature for 20 min. 1.0 ml of sodium meta arsenite was added and shaken well so that the yellow brown colour disappered. 3.0 ml of thiobarbituric acid reagent was added and heated in a boiling water bath for 15 min. After cooling, 4.3 ml of acidified butanol was added, shaken well and the colour was extracted into butanol phase. The butanol phase was transferred to another set of tubes and the colour intensity was measured at 530 nm. The blank containing 0.2 ml of 0.1N H₂SO₄ and standard sialic acid solutions were treated similarly.

Sialic acid is expressed as mg/g (tissue)

3.3.13.Lipids

3.3.13.1.Extraction of total lipids

The total lipid content of the tissues was estimated by the method of Folch *et al.* (1957).

Reagents

Chloroform-methanol mixture (2:1 v/v)

Procedure

A weighed amount of the tissue was subjected to lipid extraction using chloroform-methanol mixture (2:1). The extraction was repeated twice with fresh aliquot of chloroform-methanol mixture. The lipid extracts were transferred to a separating funnel and added 20% of water into it and left overnight. Next day the lipid extracts were drained through filter paper containing anhydrous sodium sulphate and was collected in round bottom flask and was evaporated to dryness in a flash evaporator. The lipid in the round bottom flask was made up to 10 ml by using chloroform. From this 1.0 ml was taken into a pre-weighed vial and allowed to dry in warm temperature to constant weight and total lipid content were calculated from the difference in weight. Sample made up to 10 ml was used for the estimation of various lipid components viz., cholesterol (total and free), triglycerides, free fatty acids and phospholipids after evaporating the solvent in air at room temperature.

3.3.13.2. Estimation of total cholesterol

The total cholesterol present in plasma and heart was estimated according to method of Parekh and Jung (1970) with slight modifications.

Reagents

1. Standard cholesterol solution (stock): 1mg /ml in chloroform
2. Working standard: 1.0 ml of the stock was diluted to 10 ml with chloroform.
3. FeCl₃ stock solution: 10g FeCl₃ in 100 ml acetic acid.
4. FeCl₃ - H₂SO₄ reagent: 2.0 ml of FeCl₃ stock solution was diluted to 200 ml with conc. H₂SO₄.
5. 33% KOH (w / v): 10g of KOH was dissolved in 20 ml distilled water.

6. Alcoholic KOH solution: 6.0 ml of 33% KOH was made up to 100 ml with distilled ethanol. This solution is prepared fresh before use.

Procedure

1.0 ml of the lipid sample was taken into a 25 ml glass stoppered tube and evaporated off the chloroform. Added 5 ml of freshly prepared alcoholic KOH solution. The tubes were shaken well and incubated in a water bath at 37°C for 55 min. After cooling to room temperature, added 10 ml of petroleum ether and inverted the tubes once to mix the contents. Then added 5.0 ml of distilled water and shaken the tubes vigorously for 1 min. Took 0.5-2 ml aliquots from the supernatant (petroleum ether) into test tubes. Evaporated the petroleum ether extract under nitrogen. To each of the sample as well as the standard tubes including the blank, added 3.0 ml of glacial acetic acid followed by 0.1ml -distilled water. Mixed the tubes thoroughly and added 2 ml of the FeCl₃ - H₂SO₄ reagent to the sides of the test tubes. A brown ring was formed at the interface; tap the bottom of the tubes well to effect mixing and a light colour appeared which changed to an immense purple colour, which was measured in a Shimadzu-UV spectrophotometer at 560 nm.

The amount of total cholesterol was expressed as mg/dl (serum); mg/g (tissue).

3.3.13.3. Estimation of triglycerides

The level of triglycerides in plasma and heart were determined by the method of Rice (1970) with slight modifications.

Reagents

1. Activated silicic acid.
2. Saponification reagent: 5.0g of potassium hydroxide was dissolved in 60 ml distilled water and 4.0 ml isopropanol.

3. Sodium metaperiodate reagent: To 77g of anhydrous ammonium acetate in 700 ml distilled water, added 60 ml glacial acetic acid and 650 mg of sodium metaperiodate and was dissolved and diluted to 1 litre with distilled water.
4. Acetyl acetone reagent: To 0.75 ml of acetyl acetone, 20 ml of isopropanol was added and mixed well.
5. Stock solution: 400mg of triolein was dissolved in 100 ml chloroform.
6. Working standard: 1.0 ml of the stock solution was diluted to 10 ml.

Procedure

0.2 ml of the lipid sample was taken into a test tube and evaporated off the chloroform, added 4.0 ml isopropanol. It was mixed well and added 0.4g of activated silicic acid. It was shaken in a vortex mixer for 15 min and centrifuged at 4000rpm for 5 min. To 2.0 ml of the supernatant and standards ranging from 20-100mg made up 2.0 ml with isopropanol, 0.6 ml of saponifying reagent was added and incubated at 60-70°C for 15 min. After cooling, 1.0 ml sodium metaperiodate solution was added and mixed. To this, 5ml acetyl acetone was added, mixed and incubated at 50°C for 30 min. After cooling, the colour was estimated by measuring OD at 405nm in a Shimadzu-UV spectrophotometer.

The value of triglyceride in plasma was expressed as mg per dl and in tissue as mg per gm.

3.3.13.4. Estimation of free fatty acids

Free fatty acids in plasma and heart were estimated by the modified method of Horn and Menahan (1981) with colour reagent of Itaya (1977).

Reagents

1. Activated silicic acid

2. Chloroform, heptane, methanol (CHM) solvent mixture: It was prepared by mixing chloroform, heptane and methanol in the ratio of 200:150:7(v/v)
3. Copper-triethanolamine solution: 50 ml of 0.1M copper nitrate and 50 ml of 2M triethanolamine were mixed with 33 g of sodium chloride. The pH of the solution was adjusted exactly to 8.1.
4. Diethyldithiocarbamate (DDC) solution: 0.1% DDC in butanol was prepared.
5. Standard Stock: A solution containing 2 mg per ml of palmitic acid was prepared in CHM solvent. For working standard, the stock was diluted 1:10 in CHM to give a concentration of 200 μ g per ml.

Procedure

To 1.0 ml of the lipid sample, 6.0 ml of CHM solvent and 200mg of silicic acid were added. The mixture was shaken well, centrifuged at 4000 rpm for 5 min and 3.0 ml of the supernatant taken. Standard solution in the range of 25-100 μ g were taken and made up to 3.0 ml with CHM solvent. The blank contained 3.0 ml of CHM solvent. To all these samples, 2.0 ml of copper triethanolamine solution was added and then mixed on a mechanical shaker for 10 min. The tubes were centrifuged at 4000 rpm for 5 min. To the 2.0 ml of the supernatant taken, 1.0 ml of DDC solution was added and shaken well. The colour intensity was read immediately at 430 nm in a Shimadzu-UV spectrophotometer.

Values were expressed as mg/dl serum and mg/g wet tissue.

3.3.13.5. Estimation of phospholipids

Phospholipid content of plasma and heart was estimated by the method of Fiske and Subbarow (1925) as inorganic phosphorus liberated after Bartlette's perchloric acid digestion (1959).

Reagents

1. Ammonium molybdate reagent: 2.5 g of ammonium molybdate was dissolved in 100 ml of water.
2. Aminonaphthosulfonic acid (ANSA): 0.5 g of 1,2,4 aminonaphthosulfonic acid was dissolved in 195 ml of 15% sodium metabisulfite and 50 ml of 20% sodium sulfite was added for complete solubilisation. The solution was filtered and stored in a brown bottle.
3. Stock standard solution: 351 mg of potassium dihydrogen phosphate was accurately weighed, dissolved and made upto 100 ml with double distilled water to give a final concentration of 80 mg phosphorus per ml.
4. Working standard: 1ml of the stock was diluted to 10 ml to give a conc. of 80 μg phosphorus per ml.

Procedure

1 ml of the lipid sample was taken into a test tube and evaporated off chloroform. Added 0.5 ml of perchloric acid, the tubes were made up to 3.0 ml with double distilled water, and 1.0 ml of aliquot was taken. The tubes were made up to 4.0 ml with double distilled water. To all the tubes, 0.5 ml of ammonium molybdate reagent was added. After 10 min, added 0.5 ml of ANSA to all tubes. Aliquots of the standards and blank were carried through the same procedure. The blue colour developed was read after 20 min, at 620 nm in a Shimadzu-UV spectrophotometer.

The phospholipid content of plasma was expressed as mg/dl in serum and mg/g in tissue.

3.3.14. Lipid peroxidation and tissue antioxidant status

3.3.14.1. Estimation of lipid peroxides (LPO)

Lipid peroxides content was determined by thiobarbaturic acid reaction as described by Ohkawa *et al* (1979).

Reagents

1. Acetic acid 20%: 20 ml of glacial acetic acid dissolved in 100 ml distilled water.
2. Thiobarbaturic acid: 8% in 20% acetic acid.
3. Sodium dodecyl sulphate: 8%
4. Standard: 41.66mg of Tetraethoxy propane (TEP) dissolved in 100 ml distilled water. 1.0 ml of above was made up to 10 ml with distilled water.

Procedure

To 0.2 ml of sample, 1.5 ml of 20 % acetic acid, 0.2 ml of sodium dodecyl sulphate and 1.5 ml of TBA were added. The mixture was made upto 4.0 ml with distilled water and heated in a boiling water bath for one hr. After cooling the mixture was centrifuged at 3000 rpm for 10 min. Supernatant was taken and absorbance was read at 532nm in a Shimadzu-UV spectrophotometer.

The lipid peroxides content was expressed as n moles of malondialdehyde per mg protein.

3.3.14.2. Determination of total reduced glutathione (GSH)

The total reduced glutathione was determined by the method of Ellman (1959). The method is based on the reaction of reduced Glutathione with 5,5'-

dithiosbis(2-nitrobenzoic acid) (DTNB) to give a compound that has absorbance at 412 nm.

Reagents

1. DTNB: 0.6mM in 0.2M Phosphate buffer pH 8.0
2. Phosphate buffer: 0.2M, pH 8.0.
3. Trichloroacetic acid: 5%
4. Standard: 61.4mg of reduced glutathione was dissolved in 100 ml .02M EDTA 0.1 ml of this is made up to 10 ml with 0.02M EDTA.
5. Working standard: 2.0 ml of the above was made up to 10 ml.

Procedure

0.5 ml of the homogenate was precipitated with 5% of TCA. The contents were mixed well for complete precipitation of proteins and centrifuged at 4000 rpm for 5 min. To an aliquot of clear supernatant, 2.0 ml of DTNB reagent and 0.2 M Phosphate buffer were added to make a final volume of 4.0 ml. The absorbance was read at 412nm against a blank containing TCA instead of sample, series of standards treated in a similar way were also run to determine the reduced glutathione content.

The amount of glutathione was expressed as n moles/g wet tissue.

3.3.14.3. Estimation of glutathione peroxidase (EC 1.11.1.9)

The method of Paglia and Valentine (1967) was adopted for assay of glutathione peroxidase.

Reagents

1. 0.4 M Phosphate buffer, pH 7.
2. 0.4M Tris buffer, pH 8.9
3. 0.4mM EDTA
4. 2mM GSH
5. 10mM NaN₃
6. 10% TCA
7. DTNB: 99 mg in 25 ml of methanol.
8. H₂O₂ : 1mM was prepared freshly from commercial 30% solution.
9. GSH standard: 61.4mg GSH was dissolved in 100 ml distilled water. 1.0 ml of this solution was made up to 10 ml with distilled water.
10. Working standard: 2 ml of the stock was made up to 10 ml with distilled water.

Procedure

0.2 ml of tissue homogenate was added to a mixture containing 0.2 ml of buffer, 0.2 ml of EDTA, 0.1 ml of sodium azide, mixed well and added 0.1 ml reduced glutathione and 0.1 ml of hydrogen peroxide. Incubated in a water bath at 37°C for 10min. At the end of incubation period, 0.5ml of 10%TCA was added and centrifuged at 10000rpm for 5 min. 1.0 ml of the supernatant was taken into a separate test tube and added 2.0 ml Tris buffer and 50 µl DTNB. Immediately read the OD at 412nm.

The enzyme activity was expressed as n moles of glutathione oxidized per min per mg protein.

3.3.14.4. Assay of glutathione-s-transferase (EC 2.5.1.18)

Glutathione-S-transferase activity was determined by the method of Habig *et al.* (1974).

Reagents

1. Phosphate buffer :0.3 M, pH 6.5
2. 1-chloro-2, 4-dinitrobenzene (CDNB): 30mM
3. Reduced Glutathione (GSH): 30mM.

Procedure

The reaction mixture containing 1.0 ml of buffer, 0.1 ml of CDNB and 0.1 ml of tissue homogenate was made up to 2.5 ml with water. The reaction mixture was pre-incubated at 37°C for 5min. 0.1 ml of GSH was added and the change in the absorbance was measured at 340 nm for 3 min at 30 seconds intervals.

The enzyme activity was expressed as μ moles of CDNB conjugate formed/min./mg protein.

3.3.14.5. Assay of catalase (EC 1.11.1.6)

Catalase was assayed according to the method of Takahara *et al.* (1960).

Reagents

1. Phosphate buffer: 50 mM, pH 7.0
2. Hydrogen peroxide: 30 mM solution in the above buffer.

Procedure

To 2.45 ml of the phosphate buffer, 50 μ l of the enzyme source was added and the reaction was started by the addition of 1.0 ml of H₂O₂ solution. The decrease in absorbance was measured at 240 nm at 30 sec intervals for 2 min. The enzyme blank was run simultaneously with 1.0 ml of distilled water instead of hydrogen peroxide.

The enzyme activity was expressed as n moles of H₂O₂ decomposed per minute per mg protein.

3.3.14.6. Assay of superoxide dismutase (EC 1.15.1.1)

The superoxide dismutase was assayed according to the method of Misra and Fridovich (1972) based on the oxidation of epinephrine-adrenochrome transition by the enzyme.

Reagents

1. 0.1 M Carbonate-bicarbonate buffer: pH 10.2 containing 57mg/dl EDTA.
2. Epinephrine: 3 mM

Procedure

Taken 50 μ l of sample into the cuvette and add 1.5 ml buffer and 0.5 ml epinephrine mixed well and immediately read the change in optical density at 480 nm for 2 min in a Shimadzu-UV spectrophotometer.

One unit of SOD activity was the amount of protein required to give 50% inhibition of epinephrine auto oxidation.

3.3.15. Membrane-bound ATPases

3.3.15.1. Assay of Total ATPase

Total adenonine triphosphatase was assayed according to the method described Hokins *et al.* (1973) which was modified from the method of Evans (1969).

Reagents

- | | |
|-------------------------|------------------|
| 1. Imidazole-HCl buffer | : 300 mM, pH 7.1 |
| 2. Sodium chloride | : 1200 mM |
| 3. Potassium chloride | : 200 mM |
| 4. Magnesium chloride | : 60 mM |
| 5. ATP solution | : 40 mM |
| 6. TCA | : 4 % |

Procedure

For the assay of total ATPase activity the incubation medium in a total volume of 1.0 ml contained the following: 0.1 ml each of buffer, sodium chloride, potassium chloride, magnesium chloride and 0.1 ml of enzyme solution. The mixture was incubated at 37⁰C. The reaction was terminated by the addition of 2.0 ml of TCA. The suspension was centrifuged and phosphorous content of the supernatant was estimated.

The enzyme activity is expressed as μ moles of Pi liberated per mg proten

3.3.15.2. Assay of Na⁺/K⁺ -dependent ATPase

Na⁺/K⁺ dependent ATPase activity was measured from the amount of Pi released according to the method of Bonting (1970).

Reagents

1. 184 mM Tris buffer, pH 7.

2. 50 mM Magnesium Sulphate
3. 50 mM Potassium Chloride
4. 600 mM Sodium Chloride
5. 1.0 mM EDTA
6. 40 mM ATP
7. 10% TCA

Procedure

1 ml of Tris-buffer and 0.2 ml each of the above assay reagents were mixed together. So the assay medium, in the final volume of 2.0 ml contained, 92 mM Tris-buffer, 5 mM magnesium sulphate, 60 mM sodium chloride, 5 mM potassium chloride, 0.1 mM EDTA and 4.0 mM ATP. After 10 min equilibration at 37°C in an incubator, reaction was started by the addition of 0.2 ml of the enzyme solution. The assay medium was incubated for 30 minutes and at the end of the incubation period. The reaction was stopped by the addition of 2 volumes of ice cold 10% TCA. The phosphorus (Pi) liberated was estimated by the method of Fiske and Subbarow (1925).

The Enzyme activity was expressed as μ moles of Pi liberated per mg protein per hour.

3.3.15.3. Assay of Ca²⁺-dependent ATPase

Ca²⁺ Dependent ATPase was assayed by the method of Hjerten and Pan (1983).

Reagents

1. 0.125 M, pH 8.0 Tris-HCl buffer.

2. 0.05 M Calcium chloride
3. 0.01 M ATP
4. 10% TCA

Procedure

Tris-HCl buffer 0.1 ml, Calcium 0.1 ml, ATP solution 0.1 ml and distilled water 0.1 ml were taken in test tubes. 0.1 ml of enzyme preparation was added and the tubes were incubated at 37°C for 15 min. The reaction was arrested by the addition of 10% TCA to the incubation mixture. 0.1 ml enzyme source was added to the control tubes. The contents were centrifuged at 4000 rpm for 5 min. The supernatant was used for the estimation of inorganic phosphorous.

The Enzyme activity was expressed as μ moles of Pi liberated per mg. protein per min.

3.3.15.4. Assay of Mg^{2+} -dependent ATPase

Mg^{2+} -ATPase was assayed according to the method described by Ohinishi *et al.* (1982)

Reagents

- | | |
|-----------------------|------------------|
| 1. Tris-HCl buffer | : 0.375M, pH-7.6 |
| 2. Magnesium chloride | : 0.205 M |
| 3. ATP | : 0.01 M |

Procedure

Buffer 0.1 ml, Magnesium chloride 0.1 ml and distilled water 0.1ml were taken in test tubes. 0.1 ml of enzymed preparation was added. The fibre were then incubated at 37⁰C for 15 min. The reaction was stopped by the addition of 1.0 ml of 10% TCA. 0.1 ml of enzyme was added to control tubes. The tubes were centrifuged and the phosphorous content of the supernatant was estimated according to the method of fiske and subbarow. The enzyme activity is expressed as μ moles of Pi liberated per mg. protein per min.

3.3.16.Estimation of inorganic phosphorus

Inorganic Phosphorus was estimated by the method of Fiske and Subbarow (1925). The method is based on the formation of phosphomolybdic acid by the reaction between a phosphate and molybdic acid and its subsequent reduction to a dark blue phosphomolybdic acid, the intensity of which is proportional to the phosphate ion concentration.

Reagents

1. Ammonium molybdate reagent: 2.5 g of ammonium molybdate was dissolved in 100 ml of 3 N sulphuric acid.
2. Amino naphthol sulphonic acid (ANSA): 0.5 g of ANSA was dissolved in 195 ml of 15% sodium metabisulphite and 5.0 ml of 20% sodium sulphate was added for complete solubilization. The solution was filtered and stored in a brown bottle.
3. Standard Phosphorus: 35.1 mg of potassium dihydrogen phosphate was accurately weighed, dissolved and made up to 100 ml with distilled water.

Procedure

To suitable aliquots of the supernatant, 1.0 ml of ammonium molybdate reagent was added. 0.4 ml of ANSA was added after 10 min incubation at room temperature, standards and blank were also treated in the above manner. The blue colour developed was read after 20 min at 640 nm in a Shimadzu-UV Spectrophotometer.

The values were expressed as μg per mg. protein.

3.3.17. Pepsin

The activity of pepsin was estimated by the method of Anson (1938)

Reagents

1. Pepsin substrate: Haemoglobin in 0.06 N HCl
2. Sodium carbonate : 0.55 M
3. Trichloro acetic acid : 7%
4. Standard tyrosine : 10 mg of tyrosine dissolved in 100 ml of distilled water
5. Folin's ciocaltaue reagent

Procedure

1.0 ml of the substrate was added to 0.2 ml of the enzyme and incubated at 37⁰ C for 10 minutes. The reaction was terminated by adding 1.0 ml of 7% TCA, and the mixture was centrifuged at 1000 mg for 10 min. 1.0 ml of supernatant was made alkaline by 5.0 ml of 0.55 M sodium carbonate. Then 0.5 ml of 1N Folin's

reagent was added. The optical density was measured at 660 nm after 30 min. at room temperature in Shimadzu spectrophotometer.

Results were expressed as μ moles of tyrosine liberated/4h.

3.3.18 Estimation of Amino Acids

Total amino acids in the tissue was determined as per the procedure of Rajendra (1987).

Reagents

1. HCl : 6 N
2. HCl : 0.05 M
3. Buffer A: Dissolve tri sodium citrate (58.8 g) in 2 liters of double distilled water, add 210 ml ethanol of 99.5%, adjust the pH to 3.2 by adding 60% perchloric acid and make up to 3 liters using double distilled water.
4. Buffer B: Dissolve tri sodium citrate, 58.8 g and boric acid, 12.4 g in double distilled water, adjust the pH to 10 by adding 4 N NaOH, and make up the volume to 1 liter using double distilled water.
5. Phthalaldehyde (OPA) Buffer: Dissolve 122.1 g. of Na_2CO_3 , 40.7 g. of H_3BO_3 and 56.4 g. of K_2SO_4 in double distilled water and make up the volume to 3 liters.
6. Phthalaldehyde solution (OPA): Dissolve 400 mg OPA, 7 ml ethanol, 1 ml of 2- Mercaptoethanol and 2ml of 30% w/v Brij-35 in 500 ml OPA buffer.
5. Sodium hypochlorite solution: 4% w/v Sodium hypochlorite in OPA buffer. ie., 0.3 ml Sodium hypochlorite in 100 ml OPA buffer.

3.3.18.1 Total amino acids

3.3.18.1.1 Principle

The amount of each amino acid present within a given protein does not vary from molecule and can provide useful information about the nature of the protein molecule. In a typical analysis of the amino acid content of a protein, peptide bonds are broken by acid hydrolysis with 6 N HCl at 110⁰ C (24h) so that the released amino acids can be assayed. The amino acid tryptophan is not stable to acid digestion in the presence of even trace amounts of oxygen and is estimated separately by alkali digestion.

3.3.18.1.2 Sample preparation

Take (100 mg tissue) sample in a heat stable test tube. Add 10ml 6NHCl and heat seal the tube after filling with pure nitrogen gas. Carry out the hydrolysis at 110⁰C for 24 hours. After the hydrolysis is over, open the test tube. Remove the contents quantitatively and filter into a round bottom flask through Whatmann filter paper No. 42. Wash the filter paper 2-3 times with distilled water. Flash evaporate the contents of the flask to remove all traces of HCl, the process should be repeated for 2-3 times with distilled water. Dissolve the residue and make up to 10 ml with 0.05 M HCl.

3.3.18.1.3 HPLC Analysis

Filter the sample thus prepared again through a membrane filter of 0.45 µm and inject 20 µl of this to an amino acid analyzer (HPLC- LC 10 AS) equipped with cation exchange column packed with a strongly acidic cation exchange resin i.e., styrene di vinyl benzene co polymer with sulphonic group. The column used was Na type i.e., ISC- 07/S 1504 Na having a length of 19 cm and diameter 5mm.

The instrument was equipped with Shimadzu FL 6A fluorescence detector and Shimadzu CR 6A Chrompac recorder. The mobile phase of the system consists of two buffers, Buffer A and buffer B. A gradient system can be followed for the effective separation of amino acids. The oven temperature can be maintained at 60^o C. The total run was programmed for 62 minutes. The amino acid analysis can be done with non-switching flow method and fluorescence detection after post-column derivatization with o-phthalaldehyde. In the case of proline and hydroxyl proline, imino group is converted to amino group with hypochlorite.

Run an amino acid standard (Sigma chemical Co., St. Louis, USA) also to calculate the concentration of amino acids in the sample. Calibration of equipment using standards needs to be done before the start of analysis.

3.3.18.1.4 Quantification of amino acids

The standard and the sample were analyzed under identical conditions. The elution time of the amino acids of the sample was compared and identified with those of the standard. Quantification of amino acid was done by comparing the respective peak areas in the chromatogram of the sample and the standard.

The amount of each amino acid is expressed as nmol/g tissue.

3.3.18.2 Estimation of tryptophan

Tryptophan was estimated as per the method of Sastry and Tummuru (1985) after alkali hydrolysis of the sample using 5% sodium hydroxide at 110^oC for 24 hours.

The 5-hydroxy furfural resulting from sucrose under acidic conditions formed a pale green coloured condensation product with thioglycolic acid, which on treatment with tryptophan formed a pink coloured complex. The colour intensity is measured at 500 nm.

Reagents

1. NaOH : 5%
2. HCl : 6N
3. Sucrose :2.5%
4. Thioglycolic acid : 0.6%
5. H₂SO₄ :50%
6. HCl : 0.1N
7. Tryptophan Standard: 10µg/ ml solution

Procedure

Sample was hydrolyzed with 10 ml of 5% NaOH at 110⁰C for 24 hours in a sealed tube with pure nitrogen. The hydrolysate was neutralized to pH 7.0 with 6 N HCl using phenolphthalein indicator. The volume was made up to 25 ml with distilled water. The solution was then filtered through Whatman filter paper no.42 and filtrate was used for estimation.

The test tube containing 4 ml of 50% H₂SO₄, 0.1mlk of 2.5% sucrose and 0.1ml of 0.6% thioglycolic acid were added. These tubes were kept for 5minutes in water bath at 45-50⁰C and cooled. The sample was then added to the test tubes. A set of (0.1 to 0.8ml) standard tryptophan (10µg/ ml solution) was run in a similar way. The volume was made up to 5ml with 0.1N HCl and allowed to stand for 5 minutes for the development of colour. The absorbance was measured against a reagent blank at 500 nm in a spectrophotometer.

3.3.19 Estimation of urea

Urea was estimated by the method of Natelson *et al.* (1951) using diacetyl monoxime.

Reagents

1. Diacetyl monoxime reagent: 2.0 g of diacetyl monoxime was dissolved in 100 ml of 2.0% acetic acid.
2. Sulfuric acid –phosphoric acid mixture: 25 ml of concentrated sulfuric acid, 75 ml of 85% phosphoric acid and 70 ml of distilled water were mixed.
3. Sodium tungstate solution: 10%
4. Sulfuric acid: 0.67 N
5. Standard urea solution: 20 mg urea dissolved in 100 ml of distilled water. 200 µg/ml
6. Working standard: 1.0 ml diluted to 10 ml using distilled water. 20 µg/ml

Procedure

To 0.1 ml of blood was added 3.3 ml of water and mixed with 0.3 ml of 10% sodium tungstate and 0.3 ml of 0.67 N sulfuric acid reagent. The suspensions were centrifuged and to 1.0 ml of the supernatant, 1.0 ml of water, 0.4 ml of diacetylmonoxime and 2.6 ml of sulfuric acid-phosphoric acid reagent were added in that order. Aliquots of standard urea were also treated in a similar manner and heated in a boiling water bath for 30 minutes, cooled and the colour developed was measured at 480 nm in a Shimadzu-UV spectrophotometer.

The values were expressed as mg per dl blood.

3.3.20 Estimation of creatinine

Creatinine was estimated by the method of Slot (1965)

Reagents

1. Picric acid : 40mM
2. NaOH : 0.75 M
3. Sodium tungstate solution : 100g/liter of water
4. Sulphuric acid : .33 M
5. Creatinine standard : 10 mM of creatinine in 0.1 M HCl. Just before use
6. Working standard : 2 ml of stock standard was diluted to 100 ml with distilled water.

Procedure

To 0.1ml of serum 1.5 ml of distilled water and 0.5 ml of sodium tungstate were added. 1.0ml sulphuric acid was added with shaking and allowed to stand and centrifuged. To 3.0 ml of supernatant 1.0 ml picric acid and 1.0 ml NaOH were added. The color developed was read at 500 nm after 15 minutes.

The values are expressed as mg/ dl.

3.3.21 Lipoprotein fractionation

Addition of heparin-manganous chloride to serum caused the precipitation of LDL. The supernatant represented the HDL fraction.

3.3.21.1 Estimation of high density lipoprotein fraction

Total HDL was separated by the method of Burstein and Scholnick (1972).

Reagents

1. Heparin-Manganous chloride reagent: 3.167 gm of manganous chloride was added to 1.0 ml of heparin containing 20,000 units/ml. This was made up to 8.0 ml with water.

Procedure

0.1 ml of serum was added to 9 μ l of heparin-manganous chloride reagent and mixed well. The solution was allowed to stand at 4°C for 30 min. The supernatant represented HDL fraction. Aliquots were taken from HDL fraction for the estimation of cholesterol.

3.3.21.2 Estimation of low density lipoproteins

This differential analysis was made by the method of Brustein and Scholink (1972) using sodium dodecyl sulphate.

Reagent

1. Sodium dodecyl sulphate: 10% in 0.15 M sodium chloride pH 9.0

Procedure

To 0.1 ml of serum, 75 μ l of sodium dodecyl sulphate solution was added, which was taken in a polycarbonate centrifuge tube. The contents were swirled briefly and packed for 2 hrs in a water bath at 35°C. The contents were centrifuged in a refrigerated centrifuge at 10,000g for 30 min. VLDL got aggregated as a pellicle at the top. The supernatant was a mixture containing HDL and LDL cholesterol was estimated in 0.05 ml aliquot of the supernatant as described above.

3.3.22 Analysis of fatty acids (FAME)

Fatty acids were analyzed according to the method of AOAC (1975). Lipid content of the tissues was estimated by the method of Folch et al (1957). Methyl esters of fatty acids from animal and vegetable origin having 8-24 atoms are separated and detected by gas chromatography. Method is not applicable to epoxy, oxidised, or polymerized fatty acids. (AOAC-1975 Gas chromatography method.)

Reagents

1. Boron trifluoride reagent.
2. Methanolic sodium hydroxide solution.
3. Petroleum ether.
4. Sodium sulphate.

Procedure

Add sample (lipid of known weight) to flask followed by methanolic NaOH and boiling chip. Attach condenser, and reflux until fat globules disappear (usually 5-10 minutes). Add BF_3 solution from bulb or automatic pipette through condenser and continue boiling for 2 minutes. Remove heat, then condenser, and add 15ml saturated NaCl solution. Stopper flask and shake vigorously 15s while solution is still tepid. Transfer aqueous phase to 250ml separator. Extract with two 30ml portions of petroleum ether (b.p 60-80°C). Wash combined extracts with 20ml portions H_2O , dry over anhydrous Na_2SO_4 , filter and evaporate solvent under stream of nitrogen on steam bath.

Methyl esters of the fatty acid thus obtained were separated by gas liquid chromatography (Varian CP 3800. U.S.A) equipped with a capillary column (30m long and 0.54mm diameter) and a flame ionization detector in the presence of

hydrogen and air. The carrier gas was nitrogen and the flow rate was 4ml/min the chromatograph temperature started at 150°C and was increased 4 °C/min until a temperature of 250 °C was obtained. Fatty acids separated were identified by the comparison of retention times with those obtained by the separation of a mixture of standard fatty acids. Measurement of peak areas and data processing were carried out by electronic integrator. Individual fatty acids were expressed as a percentage of total fatty acids.

3.3.23 Estimation of minerals using Atomic Absorption Spectrophotometer.

Minerals were estimated according to the method of the AOAC (1980).

Reagents

1. Nitric acid
2. Perchloric acid
3. 1 & 2 in 9:4
4. Stock solution of sodium, potassium, calcium zinc, copper, magnesium, manganese, selenium and iron were prepared by diluting concentrated solution of 1000 mg/L (Merck).

Procedure

Samples size of 1 g of tissue and 1 ml serum were used for the experiment. To the sample containing flask, 7 ml of nitric acid and perchloric acid (9:4) mixture was added, covered with a watch glass and left at room temperature over night. The sample was then digested using a microwave digester (Milestone ETHOS PLUS lab station Closed Vessel Microwave Digestion System). The completely digested samples were allowed to cool at room temperature, filtered (glass wool) carefully transferred into a clean 50 ml volumetric standard flask and then diluted to the mark with ultra pure water (Milli Q, Millipore). The digested samples were

analyzed using Varian SpectrAA 220 Atomic Absorption Spectrometer equipped with a deuterium back ground corrector, for the determination of minerals viz sodium, potassium, calcium zinc, copper, magnesium, manganese, selenium and iron.

3.4 Statistical analysis

Results are expressed as mean \pm SD. One-way analysis of variance (ANOVA) was carried out, and the statistical comparisons among the groups were performed with Tukey's test using a statistical package program (SPSS 10.0 for Windows).

4. RESULTS AND DISCUSSION

4.1. ANTIULCEROGENIC POTENTIAL OF CHITOSAN AND GLUCOSAMINE

4.1.1. Gross pathology

In gross pathology of ulcer-induced stomach, ibuprofen caused ulcerations are apparently visible as dark red lesions (**Plate 4.1.2**). These are non-malignant wounds on the mucosal membrane by the action of gastric acid. In control animals, the mucosal membrane was lesion free (**Plate 4.1.1**). Chitosan and glucosamine treated animals showed lesser number of ulcerations on the mucosa (**Plates 4.1.3 and 4.1.4**) showing the antiulcer effect of these compounds.

4.1.2. Histopathology

In the ibuprofen ulcer models, more facial tissue differentiations were observed after administration of the active fraction in bleeding areas and mucosa than those of control group. On the other hand, inflamed cell infiltration and edema in lamina propria were pronounced in the test group when compared with the control group on *per os* administration, while the effects were less on subcutaneous administration. This observation can be interpreted as the co-existence of some compounds in the active fraction, which might affect the gastric mucosa on *per os* application (**Plate 4.1.8**).

In chitosan and glucosamine pre treated animals, histopathological results of the stomach shows a normal cell architecture (**Plates 4.1.9 and 4.1.10**) as that of control animals (**Plate 4.1.5**). All the tissue alterations were normalized by chitosan and glucosamine. Histology of the stomach of rats fed with chitosan or glucosamine alone did not show any characteristic change (**Plates 4.1.6 and 4.1.7**).




Plate 4.1.1. Gross pathology of the stomach of normal control rat



Plate 4.1.2. Ulcer induced stomach of albino rat by the administration of ibuprofen. Mucosal lesion is indicated by arrow




Plate 4.1.3. Gross pathology of the ulcer treated rat stomach with chitosan pre-oral administration, which shows lesser mucosal lesions compared to ulcerated rats.




Plate 4.1.4. Stomach of ulcerated rat treated by the pre-oral administration of glucosamine. Most of the ulcerated lesions were vanished as compared to ulcer-induced rats by the treatment with glucosamine.

4.1.3. Gastric mucosal lesions

Table 4.1.1 shows the number of lesions present in the gastric mucosa, of the normal and experimental groups of rats.

Experimental peptic ulcer may be assessed on the basis of number of gastric mucosal lesions. In the present study, a significant ($P<0.001$) increase in the number of lesions in the gastric mucosa was observed in Group III ulcer induced rats as compared to that of Group I control animals. This is in accordance with an earlier report (Lanza *et al.*, 1989), which showed that ibuprofen-induced lesion formation may be multifactorial, with stasis of gastric flow contributing significantly to the hemorrhagic as well as the necrotic aspects of the tissue injury. The products of the 5-lipoxygenase pathway may also play a key role in the development of this ulcer (Rainsford, 2005). The prior oral administration of chitosan and glucosamine resulted in significant ($P<0.001$) reduction in the number of lesions in the gastric mucosa of Group V and Group VI rats as compared to Group IV, ulcer induced rats, indicating the cytoprotective activity of chitosan and glucosamine. Mutoh *et al.* (1995) have postulated that the adoptive cytoprotection in cultured gastric mucus producing cells are mediated by glucosamine rich mucus released in response to a mild irritant. But animals administered with chitosan alone and glucosamine alone did not show adverse effects, which show that chitosan and glucosamine are non-toxic.

4.1.4. Volume of gastric juice, acidity and peptic activity

Volume of gastric juice, acid out put and peptic activity in normal and experimental groups of rats were detailed in **Table 4.1.1**.

Significant ($P<0.001$) increase in acidity and volume of the gastric juice was noticed in Group III ulcer induced rats as compared to that of Group I control animals. Also a significant decline in peptic activity was also observed. These findings concur with an earlier reported study (Welage, 2005). Increased production of hydrochloric acid in ulcerated condition might be a consequence of

Table 4.1.1: Number of lesions, volume of gastric juice, and acid output and pepsin activity of the gastric mucosa of normal and experimental groups of rats.

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
	Normal Control	Chitosan-Control A	Glucosamine-Control B	Ulcer-Induced C	A+C	B+C
No of lesions	-	-	-	8.78±1.42 ^{a,b,c}	2.34±0.09 ^d	1.25± 0.06 ^d
Volume of Gas Juice	1.45±0.05	1.39±0.03	1.37±0.04	2.76±0.09 ^{a,b,c}	1.87±0.06 ^d	1.55±0.05 ^d
Acid out put	156±12.3	146±11.2	138±14.7	293±18.2 ^{a,b,c}	205±16.6 ^d	172±15.5 ^d
Pepsin	723±58.1	712±55.3	698±63.5	576±41.9 ^{a,b,c}	668±53.2 ^d	682±54.7 ^d

Values are expressed as, Mean ± SD for six animals in each group.

Volume of gastric juice- ml/4h; Acid output- µEq/4h; Pepsin- µmol tyrosine liberated/4h.

^a P<0.001- compared with Group-I

^b P<0.001- compared with Group-II

^c P<0.001- compared with Group-III

^d P<0.001- compared with Group-IV

^e P<0.01 - compared with Group-III

^f P<0.01 - compared with Group-II

^g P<0.05 - compared with Group-II

^h P<0.05 - compared with Group-IV




Plate 4.1.5. Histopathology of the gastric mucosa of the normal rat.




Plate 4.1.6. Histopathology of the stomach of chitosan control rat




Plate 4.1.7. Histopathology of the stomach of glucosamine control rat




Plate 4.1.8a Illustrates the histopathology of the gastric mucosa of ulcer-induced rat (X100)




Plate 4.1.8b. Illustrates the histopathology of the gastric mucosa of ulcer-induced rat showing the cell necrosis (X200).




Plate 4.1.9. Histopathology of the stomach of chitosan + ibuprofen administered rat




Plate 4.1.10a. Illustrates the histopathology of the gastric mucosa of glucosamine + ibuprofen induced rat (X100)





Plate 4.1.10b Illustrates the histopathology of the gastric mucosa of glucosamine + ibuprofen administered rat showing near to normal architecture (X200)

increased permeability of the mucosa, which is an important process in the development of ulcer Bardhan *et al.* (1994). In the present study, the prior oral administration of chitosan and glucosamine resulted in significant ($P<0.001$) reduction in the volume of gastric juice and acid output in Group V and Group VI animals as compared to Group IV ulcer induced rats. Peptic activity was also maintained at near normal level. It probably did so by the neutralization of hydrochloric acid excessively secreted into the stomach. The neutralization of acid secretion in stomach has already been reported to accelerate ulcer healing (Weberg *et al.*, 1990).

4.1.5. Proteins

The levels of protein are featured in **Fig. 4.1.1**. The level of protein was significantly ($P<0.001$) decreased in the gastric mucosa of Group IV ulcerated rats as compared to that of Group I normal rats. This is indicated that administration of ibuprofen caused the corrosion of gastric mucosa, resulting in disruption and disintegration of gastric mucosal cells. Similar observations have been reported earlier also (Vanisree and Devaki, 1995). The ulcerative lesions observed in the gastrointestinal tract may also be associated with increased loss protein. According to the Davenport's concept of H^+ release, histamine stimulates cholinergic nerves with a resulting stimulation of acid secretion, an increase in mucosal blood flow, and an increase in capillary permeability (Davenport, 1967). The net effect would be the loss of protein. In the present study, oral pre-treatment with chitosan and glucosamine significantly ($P<0.001$) prevented the ibuprofen-induced erosion of protein in the gastric mucosa of Group V and Group VI rats, indicating the antiulcer effects of chitosan and glucosamine. The SDS-polyacrylamide gel electrophoretic separation (**Plate 4.1.11.**) also confirmed the loss of proteins in the ulcerated group as compared to normal rats. The oral chitosan and glucosamine pretreatment nullified the effect of ibuprofen and brought back the level of protein to the near normalcy.

Fig. 4.1.1: Protein levels in the gastric mucosa of normal and experimental groups of rats.



Mean ± SD for six animals in each group.

^a p<0.001- compared with Group-I, ^b p<0.001- compared with Group-II, ^c p<0.001- compared with Group-III,

^d p<0.001- compared with Group-IV.




Plate 4.1.11. Electrophoretic pattern of stomach protein of normal and experimental groups of rats.

- BSA- Bovine Serum Albumin**
- I- Normal control rats**
 - II- Chitosan control**
 - III- Glucosamine control**
 - IV- Ulcer induced**
 - V- Chitosan pre treated**
 - VI- Glucosamine pre treated**


4.1.6. Hexose, hexosamine and sialic acid

Fig. 4.1.2. depicts the levels of hexose and hexosamine and Fig. 4.1.3. shows the level of sialic acid in the normal and experimental groups of rats. Intra-gastric administration of ibuprofen caused significant ($P < 0.001$) depletion in the level of gastric mucosal protein as compared to that of Group I control animals. Significant ($P < 0.001$) reduction in the levels of mucosal glycoprotein components (hexose, hexosamine and sialic acid) were also observed in the ulcerated mucosa of Group IV animals as compared to Group I normal rats. Prior oral administration of chitosan and glucosamine in Group V and Group VI animals significantly ($P < 0.001$) prevented all these alterations and maintained the rats at near normal status. The normal rats receiving chitosan or glucosamine alone did not show any significant change when compared with the normal rats, showing that it does not *per se* have any adverse effects.

Mucus is said to protect the gastrointestinal tract from infective, chemical and physical insults. The glycoprotein component of mucus, which is responsible for the characteristic viscous gel forming property, is believed to be important for the functional role of mucus (Allen *et al.*, 1997). This mucosal barrier lessens the stomach wall friction during contraction, improves the buffering acid and acts as an effective barrier to back diffusion of hydrogen ions (Pfeiffer, 1981). The carbohydrate side chains of mucosal glycoprotein comprise over 80% by weight of the glycoprotein molecule. Each chain consists of sugars such as hexose, amino sugars and sialic acid (Zarzycki *et al.*, 1986). In this study, a significant ($P < 0.001$) reduction in the levels of glycoprotein components were observed in the gastric mucosa of Group IV ibuprofen-administered animals as compared to that of Group I rats. Administration of ibuprofen might have affected the cellular processes of exocytosis and apical expulsion or pinching off the apical membrane or exfoliation responsible for the secretion of glycoproteins from the surface epithelial cells by causing the erosion of the epithelial cells.

In the present study, oral pre-treatment with chitosan and glucosamine significantly ($P < 0.001$) prevented the ibuprofen-induced depletion of glycoprotein

Fig. 4.1.2: Levels of hexose and hexosamine in the gastric mucosa of normal and experimental groups of rats.




Mean ± SD for six animals in each group.

^a p<0.001 - compared with Group-I, ^b p<0.001 - compared with Group-II, ^c p<0.001 - compared with Group-III,

^d p<0.001 - compared with Group-IV

Fig. 4.1.3: Level of sialic acid in the gastric mucosa of normal and experimental groups of rats.



Mean ± SD for six animals in each group.

^a p<0.001- compared with Group-I, ^b p<0.001- compared with Group-II, ^c p<0.001- compared with Group-III

^d p<0.001- compared with Group-IV

Table 4.1.2: Activities of membrane bound ATPases of the gastric mucosa of normal and experimental groups of rats

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
	Normal Control	Chitosan-Control A	Glucosamine-Control B	Ulcer-Induced C	A+C	B+C
Total ATPase	6.73±0.58	6.5±0.54	6.52±0.05	5.58±0.47 ^{e,f,g}	6.39±0.53 ^h	6.42±0.54 ^h
Na-K ATP	4.48±0.38	4.29±0.37	4.33±0.41	3.57±0.31 ^{e,f,g}	4.20±0.42 ^h	4.21±0.44 ^h
Ca-ATP	3.72±0.35	3.80±0.31	3.86±0.33	2.61±0.28 ^{a,b,c}	3.70±0.32 ^d	3.74±0.34 ^d
Mg-ATP	4.25±0.37	4.52±0.38	4.48±0.47	3.19±0.32 ^{a,b,c}	4.47±0.38 ^d	4.58±0.45 ^d

Mean ± SD for six animals in each group. Values are expressed as μ moles of Pi liberated per mg protein per hour.

^a $P < 0.001$ - compared with Group-I

^b $P < 0.001$ - compared with Group-II

^c $P < 0.001$ - compared with Group-III

^d $P < 0.001$ - compared with Group-IV

^e $P < 0.01$ - compared with Group-I

^f $P < 0.01$ - compared with Group-III

^g $P < 0.05$ - compared with Group-II

^h $P < 0.05$ - compared with Group-IV

Table 4.1.3: Levels of reduced glutathione (GSH) and lipid peroxides (LPO) in the gastric mucosa of normal and experimental groups of rats.

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
	Normal Control	Chitosan-Control A	Glucosamine-Control B	Ulcer-Induced C	A+C	B+C
GSH	5.02±0.36	4.82±0.23	4.89±0.25	2.12±0.12 ^{a,b,c}	4.21±0.15 ^d	4.73±0.18 ^d
LPO	1.02±0.02	0.98±0.02	0.96±0.01	2.45±0.05 ^{a,b,c}	1.46±0.03 ^d	1.18±0.01 ^d

Mean ± SD for six animals in each group. Values are expressed as: LPO, nmol MDA/mg protein; GSH, nmol/g wet tissue

^a $P < 0.001$ - compared with Group-I

^b $P < 0.001$ - compared with Group-II

^c $P < 0.001$ - compared with Group-III

^d $P < 0.001$ - compared with Group-IV

fatty acids (PUFA), has been linked with altered membrane structure and enzyme inactivation. The level of lipid peroxidation was found to be significantly ($P<0.001$) higher in the gastric mucosa of Group IV ulcer-induced rats as compared to that of normal controls. Significant increases in the levels of lipid peroxides in the ulcer induced gastric mucosa have already been reported (Yoshikawa *et al.*, 1986, Yoshikawa *et al.*, 1987). Lipid peroxidation worsens the mucosal injury. The exact sources of peroxy radicals in peptic ulcer are not yet thoroughly understood. Xanthine oxidase has been considered as one of the major source of free radicals (Yoshikawa *et al.*, 1990). Another source may be neutrophils, since administration of non-steroidal anti-inflammatory drugs has already been reported to cause neutrophil infiltration into the gastric mucosa (Amagase *et al.*, 2003). Jimenez *et al.* (2004) proposed that the active oxygen species, derived both from xanthine oxidase and activated neutrophils, could play a role in the pathogenesis of gastric injury induced by ibuprofen. The metabolism of arachidonic acid via the lipoxygenase and cyclooxygenase pathways may also result in the formation of reactive oxygen species and other free radicals in the ulcerated gastric mucosa (Argentieri *et al.*, 1994)

Prior administration of chitosan and glucosamine resulted in a significant ($P<0.001$) reduction in the level of lipid peroxidation towards near normalcy as compared to Group IV ulcer-induced rats, establishing their cytoprotective effect. The unpaired electron present in the ibuprofen-generated free radicals might have been trapped and subsequently dismuted by chitosan and glucosamine by the free radical scavenging capability (Yan *et al.*, 2006; Sal'nikova *et al.*, 1990).

4.1.9. Reduced glutathione (GSH)

Here, GSH level was significantly ($P<0.001$) decreased in ulcerated group as compared to normal rats (**Table 4.1.3**). GSH has a direct antioxidant function. It functions by reaction with superoxide radicals, peroxy radicals and singlet oxygen, followed by the formation of oxidized glutathione and other disulphides (Anandan *et al.*, 1999). Depletion of GSH results in enhanced lipid peroxidation, and excessive lipid peroxidation can cause increased GSH consumption

Anandan *et al.*, 2004), as observed in the present study This is in accordance with a previously reported study (Lu *et al.*, 1989), which indicated that the depletion of gastric GSH by diethyl maleate produced gastric ulceration. This reduction might have resulted from the oxidation of GSH to ibuprofen-induced generation of free radicals. The depletion of GSH further enhances the susceptibility of the gastric mucosal cells to oxygen metabolites and acid mediated cell damage. Hiraishi *et al.* (1994) has reported that exogenous GSH protects cultured gastric mucosal cells from oxidant-induced damage.

in the present study, the pretreatment with chitosan and glucosamine resulted in significant ($P<0.001$) elevation of the GSH level, which protects against oxidative damage by the way of regulating the redox status of proteins in the cell membrane (Anandan *et al.*, 2004). This increase in intracellular GSH in turn may protect the gastric cells against ibuprofen mediated cellular injury by decreasing their susceptibility to free radicals and acid.

4.1.10. Mucosal antioxidant enzymes

Activities of glutathione-dependent antioxidant enzymes [glutathione peroxidase (GPX) and glutathione s-transferase (GST)] and antiperoxidative enzymes [catalase (CAT) and super oxide dismutase (SOD)] were significantly ($P<0.001$) lower in the gastric mucosa of Group IV ulcer-induced rats as compared to that of Group I control animals (**Table 4.1.4**). GPX, an antioxidant enzyme, offers protection to the mucosal membrane from peroxidative damage (Prabha *et al.*, 2003). A significant decrease in the activity of GPX makes mucosal membrane susceptible to ibuprofen-induced peroxidative damage, which leads to changes in the gastric mucosal composition and function. GST, another scavenging enzyme, binds to many different lipophilic compounds (Anandan *et al.*, 1999); so it would be expected to bind ibuprofen and act for GSH conjugation reactions. The significant ($P<0.001$) decrease in GST activity noted in this study might have been due to the decreased availability of GSH.

Table 4.1.4: Activities of glutathione-S-transferase (GST), glutathione peroxidase (GPx), superoxide dismutase (SOD) and catalase (CAT) in the gastric mucosa of normal and experimental groups of rats.

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
	Normal Control	Chitosan-Control A	Glucosamine-Control B	Ulcer-Induced C	A+C	B+C
GST	5.46±0.38	5.31±0.37	5.34±0.33	3.18±0.25 ^{a,b,c}	4.61±0.32 ^d	4.95±0.34 ^d
GPx	179±14.2	184±14.4	195±16.8	108±7.3 ^{a,b,c}	153±13.4 ^d	172±15.5 ^d
SOD	5.87±0.27	5.95±0.28	6.02±0.32	2.25±0.18 ^{a,b,c}	4.46±0.20 ^d	5.18±0.22 ^d
CAT	4.08±0.21	3.90±0.20	3.95±0.26	1.25±0.09 ^{a,b,c}	3.49±0.16 ^d	3.82±0.18 ^d

Mean ± SD for six animals in each group. Values are expressed as: GST, μmol 1-chloro-2,4 dinitrobenzene conjugate formed /min /mg protein; GPx, nmol GSH oxidized/min/mg protein; SOD, one unit of the superoxide dismutase activity is the amount of protein required to give 50% inhibition of epinephrine autooxidation; CAT, μmol of H_2O_2 consumed/min/mg protein.

^a $P < 0.001$ - compared with Group-I

^b $P < 0.001$ - compared with Group-II

^c $P < 0.001$ - compared with Group-III

^d $P < 0.001$ - compared with Group-IV

Significant reduction ($P<0.001$) noticed in the activities of CAT and SOD in experimentally induced ulcerated mucosa is in line with earlier reported studies (Prabha *et al.*, 2003; Raghavendran *et al.*, 2004), indicating that decline in the activities of these antiperoxidative enzymes leads to the formation of peroxy radicals, which are harmful to the gastric mucosa. In the present study, the prior oral administration of chitosan and glucosamine significantly ($P<0.001$) prevented the ibuprofen-induced alterations in the activities of glutathione-dependent antioxidant enzymes (GPX and GST) and antiperoxidative enzymes (SOD and CAT) in the gastric mucosa of Group IV rats. It probably did so by counteraction of ibuprofen-induced free radicals (Balasubramanian *et al.*, 2004) due to the antioxidant nature.


The results of the present study indicates that the overall antiulcerogenic effect of chitosan and glucosamine is probably related to its ability to maintain near to the normal status the activities of the mucosal antioxidant enzymes and the level of GSH which protects mucosa against oxidative damage, by decreasing lipid peroxidation.

4.1.11. Minerals

Mineral status of the stomach tissue of normal and experimental groups of rats were described in **Fig. 4.1.4** and **Table 4.1.5**. Rats treated with ibuprofen showed significant elevation ($P<0.001$) in concentration of calcium in the stomach tissue. The accumulation of intracellular calcium levels also play an important role in the DNA fragmentation and ulcer induced by ibuprofen. An increase in the level of intracellular calcium can simply cause cellular damage via activation of calcium dependent degenerative enzymes including calcium dependent phospholipases, proteases and endonucleases. This could lead to destruction of membranes, critical protein and DNA fragmentation (Timbrell, 2000)

The level of sodium was found to decrease significantly ($P<0.001$) in the ulcerated group. But, levels of potassium, magnesium and zinc were found to

Fig. 4.1.4: Sodium and potassium levels in the stomach of normal and experimental groups of rats.



Mean ± SD for six animals in each group.

^a p<0.001- compared with Group-I, ^b p<0.001- compared with Group-II, ^c p<0.001- compared with Group-III,

^d p<0.001- compared with Group-IV

Table 4.1.5: Mineral status of the stomach of normal and experimental groups of rats

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
	Normal Control	Chitosan-Control A	Glucosamine-Control B	Ulcer-Induced C	A+C	B+C
Calcium	71±0.21	74±0.22	73±0.22	110±0.38 ^{a,b,c}	80±0.27 ^d	78±0.25 ^d
Magnesium	60±0.18	65±0.19	63±0.18	98±0.35 ^{a,b,c}	68±0.21 ^d	66±0.20 ^d
Zinc	55±0.15	59±0.17	57±0.16	92±0.34 ^{a,b,c}	61±0.19 ^d	59±0.16 ^d

Mean ± SD for six animals in each group. Values are expressed as mg/g tissue.

^a $P < 0.001$ - compared with Group-I

^b $P < 0.001$ - compared with Group-II

^c $P < 0.001$ - compared with Group-III

^d $P < 0.001$ - compared with Group-IV

increase significantly ($P<0.001$) in this group. Increased peroxidation may be the cause of increment in the magnesium level. An earlier reported study (Bironaite and Ollinger, 1997) also supports this. Oral pre treatment of chitosan and glucosamine significantly ($P<0.001$) maintained the levels of these minerals to the normalcy.

4.1.12. Amino acids

There was a significant decrease ($P<0.001$) in the amino acid levels of ulcer-induced rats as compared to normal control animals (**Table 4.1.6.**). This may be due to the cytodeterioration brought about by the ibuprofen created free radicals. In, Groups V and Groups VI animals, due to the administration of chitosan and glucosamine, the levels of these amino acids were maintained to near normalcy. Inhibition of the transsulfuration reactions from homocysteine to cysteine may lead to a decrease in the synthesis of GSH, which would produce an important impact on normal biochemistry and physiology (Sang and Young, 2005). Here, the cytoprotective effect of chitosan and glucosamine plays a good role.

Table 4.1.6: Levels of various amino acids in the stomach of normal and experimental groups of rats

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
	Normal Control	Chitosan-Control A	Glucosamine-Control B	Ulcer-Induced C	A+C	B+C
Glycine	1800±81.25	1825±85.34	1831±86.16	1685±76.52 ^{a,b,c}	1790±78.37 ^d	1798±78.63 ^d
Serine	470±24.32	475±25.51	478±25.82	401±19.34 ^{a,b,c}	465±23.63 ^d	468±24.52 ^d
Valine	340±16.28	351±19.68	357±20.34	269±12.93 ^{a,b,c}	331±14.57 ^d	337±15.69 ^d
Histidine	520±25.24	528±26.46	532±26.51	445±19.37 ^{a,b,c}	511±24.91 ^d	515±24.87 ^d
Tryptophan	89±6.93	95±7.64	97±8.82	45±2.38 ^{a,b,c}	78±5.64 ^d	84±5.98 ^d
Threonine	290±20.35	301±22.56	305±22.72	180±16.19 ^{a,b,c}	278±18.82 ^d	287±20.14 ^d
Leucine	155±8.62	160±9.14	163±11.17	91±6.52 ^{a,b,c}	148±7.16 ^d	150±7.85 ^d

Mean ± SD for six animals in each group. Values are expressed as nmol/g tissue

^a P<0.001- compared with Group-I

^b P<0.001- compared with Group-II

^c P<0.001- compared with Group-III

^d P<0.001- compared with Group-IV

4.2. ANTI HEPATOTOXIC POTENTIAL OF CHITOSAN AND GLUCOSAMINE

4.2.1 Diagnostic marker enzymes

Antibiotic-related liver injuries cover most of the clinical and pathological expressions of hepatic dysfunction, including cytotoxic hepatitis (Westphal *et al.*, 1994). There was a significant ($P < 0.001$) elevation observed in the levels of alanine aminotransferase (ALT), aspartate aminotransferase (AST), lactate dehydrogenase (LDH), alkaline phosphatase (ALP) and acid phosphatase (ACP) in the serum of Group IV antitubercular drugs-administered rats as compared to Group I normal controls (**Table 4.2.1**), which is in line with an earlier reported study (Imogen *et al.*, 1995). Elevated levels of these enzymes in serum are presumptive markers of drug-induced necrotic lesions in the hepatocytes (Amr and Alaa, 2005). Enhanced susceptibility of hepatocytes cell membrane to the isoniazid and rifampicin-induced peroxidative damage might have resulted in increased release of these diagnostic marker enzymes into the systemic circulation. In the present study, co-administration of chitosan and glucosamine significantly ($P < 0.001$) maintained the levels of these diagnostic marker enzymes in the serum of Group V and Group VI animals at near normalcy as compared to Group IV rats, indicating the cytoprotective effect of chitosan and glucosamine (Anandan *et al.*, 2004, Sal'nikova *et al.*, 1990). It probably did so by preventing the antitubercular drugs-induced necrotic damage by the membrane stabilizing action (Filipovic-Grcic *et al.*, 2001). Reports by Anandan *et al.* (2004) showed that chitosan administration prevented experimental ulcer in rats by its cytoprotectivity.

Amino transferases are important class of enzymes linking carbohydrate and amino acid metabolism, thereby clearly establishing the relationship between the intermediates of the citric acid and amino acids. Alanine amino transferase and

Table 4.2.1: Levels of alanine aminotransferase (ALT), aspartate aminotransferase (AST), lactate dehydrogenase (LDH), acid phosphatase (ACP) and alkaline phosphatase (ALP) levels in the serum of normal and experimental groups of rats

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
	Normal Control	Chitosan-Control A	Glucosamine-Control B	Hepatotoxicity C	A+C	B+C
ALT	121±7.23	112±6.31	128±7.45	314±29.61 ^{a,b,c}	133±9.54 ^d	153±12.65 ^d
AST	102±9.02	94.5±5.69	115±10.23	297±26.74 ^{a,b,c}	98.4±5.23 ^d	127±10.65 ^d
LDH	178±14.92	182±16.0	185±16.25	284±25.63 ^{a,b,c}	193±17.56 ^d	210±19.85 ^d
ACP	14.3±0.98	12.7±1.02	14.2±1.13	36.8±2.41 ^{a,b,c}	16.5±1.18 ^d	19±1.21 ^d
ALP	98.4±7.46	94.1±7.21	97.5±7.35	224±18.0 ^{a,b,c}	118±9.77 ^d	123.6±10.1 ^d

ALT, AST and LDH=μmoles of pyruvate liberated/hr/l; ACP and ALP=μmoles of phenol liberated/hr/l

Values are expressed as the mean ± SD for six animals.

^a P<0.001- compared with Group-I

^b P<0.001- compared with Group-II

^c P<0.001- compared with Group-III

^d P<0.001- compared with Group-IV


aspartate amino transferase are well known diagnostic indicators of liver disease. The levels of ALT and AST in the liver of normal and experimental groups of rats were shown in **Fig. 4.2.1**. Antitubercular drugs administration in Group IV animals significantly ($P<0.001$) lowered the levels of ALT and AST as compared to Group I normal rats. Co-administration of chitosan and glucosamine brought back the levels of these diagnostic marker enzymes to the normal level. Significant increase in the levels of ALT [$(P<0.001)$, $(P<0.01)$] and AST ($P<0.001$) was observed in Group V and Group VI animals. In cases of liver damage with hepatocellular lesions and parenchymal cell necrosis, these marker enzymes are released from the damaged tissues into the blood stream. So, in the present study, the activities of these enzymes were significantly lower in the liver of Group IV, antitubercular drugs-administered rats as compared to that of control rats. Observations of significant decrease in the activities these enzymes have already been reported in the liver of isoniazid and rifampicin intoxicated rats (Saraswathy *et al.*, 1998).

The fall in alanine amino transferase activity is almost always due to hepatocellular damage and is usually accompanied by a lowering in the activity of aspartate amino transferase. Increased protein catabolism and urea formation that are seen in antitubercular drugs-induced hepatocellular damage may also be responsible for the decline of these amino transferases activities in liver. In the present study, the activities of alanine amino transferase and aspartate amino transferase were maintained at near normal level in the orally chitosan and glucosamine co-administered rats as compared to that of Group IV antitubercular drugs-induced hepatotoxic rats, showing the normalizing effect of chitosan and glucosamine on protein metabolism.

4.2.2. Histopathology

Liver sections of control rats revealed the normal hepatic architecture and the typical hexagonal hepatic lobules (**Plate 4.2.1**). Liver tissue of isoniazid and rifampicin treated rats showed severe vacuolation and degeneration of hepatocytes in the three zones of lobules. Membrane disintegration with intense

Fig. 4.2.1: Levels of alanine aminotransferase (ALT) and aspartate aminotransferase (AST) in the liver of normal and experimental groups of rats.



Values are expressed as the mean ± SD for six animals.

^a $P < 0.001$ - compared with Group-I, ^b $P < 0.001$ - compared with Group-II, ^c $P < 0.001$ - compared with Group-III,

^d $P < 0.001$ - compared with Group-IV, ^e $P < 0.01$ - compared with Group-IV

vacuolation accompanied with cytoplasmic rarefaction resulting in loss of polyhedral structure in the liver of rat treated with antitubercular drugs (**Plate 4.2.4**). There was severe infiltration of inflammatory cells in the portal areas and around the necrotic cell. These are in line with an earlier reported study (Victor and John, 2006). Congested blood vessels and sinusoids were also observed in some cases. Central vein dilation, enlargement of portal areas and bile duct proliferation were seen. However, the other abnormalities like necrosis, macrovesicular steatosis and inflammation were also occasionally present.

Co-administration with chitosan and glucosamine caused reversal of such pathology with no evidence of necrotic areas (**Plates 4.2.5. and 4.2.6.**), which establishes the cytoprotectivity of these marine compounds. Liver architecture of rats, fed with chitosan and glucosamine alone did not show any characteristic change (**Plates 4.2.2. and 4.2.3.**).

4.2.3. Urea, albumin, globulin and albumin/globulin

Drug induced hepatotoxicity is associated with failure to maintain serum albumin level whereas serum globulins, particularly γ -globulin, which is formed in the reticuloendothelial system are increased. Serum albumin level is an index of severity and prognosis in patients suffering from chronic hepatitis. So, in the present study, the level of globulin was elevated to a significantly ($P<0.05$) and levels of albumin and albumin/globulin ratio were found to be decreased significantly ($P<0.001$) in isoniazid and rifampicin induced group as compared to normal rats (**Table 4.2.2**). Formation of urea, creatine and creatinine are the mode of disposal of nitrogen. Liver is the most important organ in the maintenance of blood ammonia levels through the urea cycle (Stryer, 1995). In hepatotoxic condition, due to the failure of the liver to convert amino acids and ammonia to urea, a significant ($P<0.001$) decrease in urea (**Table 4.2.2**) was observed (Reicher and Paumgartner, 1980). There is an increased catabolism of proteins coupled with the diminished ability of kidneys to excrete the nitrogenous waste. This could be possibly the reason for the lowered level of urea in hepatotoxic rats. This shoulders with a previous report (Eule *et al.*, 1986).




Plate 4.2.1. Histopathological observation of the liver of normal rat.




Plate 4.2.2. Histopathological observation of the liver of chitosan control rat.




Plate 4.2.3. Histopathological architecture of the liver of glucosamine control rat.




Plate 4.2.4a Illustrates the histopathology of the liver of hepatotoxic rat showing cell necrosis




Plate 4.2.4b. Illustrates the histopathology of the liver of hepatotoxicity induced rat. Necrosis is visible.




Plate 4.2.5a Illustrates the histopathology of the liver of the rat co-administered with chitosan along with antitubercular drugs.




Plate 4.2.5b. Illustrates the histopathology of the liver of chitosan co-administered with isoniazid and rifampicin.




Plate 4.2.6b. Shows the histopathological architecture of rat liver in which glucosamine co-administered with antitubercular drugs.




Plate 4.2.6a Illustrates the histopathology of the liver of the liver of the rat, glucosamine co-administered with isoniazid and rifampicin.

Table 4.2.2: Levels of urea (mg/dl), albumin(g/dl), globulin(g/dl), albumin/globulin ratio (g/dl) in the serum of normal and experimental groups of rats

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
	Normal Control	Chitosan-Control A	Glucosamine-Control B	Hepatotoxicity C	A+C	B+C
Urea	22.1±1.90	20.8±2.20	20.7±2.10	11.4±1.30 ^{a,b,c}	18.6±1.55 ^d	17.9±1.51 ^d
Albumin	4.12±0.32	4.21±0.30	4.24±0.33	2.18±0.23 ^{a,b,c}	3.95±0.28 ^d	3.92±0.26 ^d
Globulin	3.26±0.35	3.28±0.33	3.30±0.36	3.89±0.37 ^{e,f,g}	3.31±0.29 ^h	3.34±0.32 ^h
Albumin/globulin	1.25±0.08	1.19±0.09	1.17±0.09	0.53±0.03 ^{a,b,c}	1.04±0.07 ^d	1.03±0.08 ^d

Values are expressed as the mean ± SD for six animals.

^a P<0.001- compared with Group-I, ^b p<0.001- compared with Group-II, ^c p<0.001- compared with Group-III,

^d P<0.001- compared with Group-IV, ^e p<0.05 - compared with Group-I, ^f p<0.05 - compared with Group-II,

^g P<0.05 - compared with Group-III, ^h p<0.05 - compared with Group-IV

Creatinine level (**Table 4.2.3**) was also found to be decreased at significant range ($P<0.001$). Co-administration of chitosan and glucosamine in the present study significantly prevented antitubercular drugs-induced alterations in the levels of creatinine ($P<0.001$) (**Table 4.2.3**) albumin ($P<0.001$), globulin ($P<0.05$), albumin/globulin ratio ($P<0.001$) and urea ($P<0.001$) in Group V and Group VI rats as compared to that of Group IV rats (**Table 4.2.2**). It probably did so by reducing the accumulation of toxic antitubercular drugs derived metabolites, which may contribute to the changes in the rough endoplasmic reticulum and the disturbance of protein metabolism in liver.

4.2.4. Serum glucose

Serum glucose level of experimental and normal groups of rats were described in **Table 4.2.3**. The level of glucose in the serum of hepatotoxic rats were found to be decreased significantly ($P<0.001$). Administration of isoniazid and rifampicin brought about hypoglycaemia, which may be due to the inactivated pancreatic β cell function and to the inhibition in the activities of the key gluconeogenic enzymes or increase the activities of key glucose utilizing enzymes. The changes in the level of glucose in the serum were normalized significantly by the co-administration of chitosan and glucosamine ($P<0.001$). Since, chitosan and glucosamine were of saccharide family, they can increase the gluconeogenesis (Krisana *et al.*, 2004). This may be the reason for the hyperglycaemic action of these compounds.

4.2.5. Proteins

The oral administration of antitubercular drugs to rats induces liver cell damage leading to periportal infiltration of inflammatory cells, disruption of lobular architecture, and liver cell necrosis and causes severe disturbance of RNA and protein metabolism. The disaggregation of polyribosomes in isoniazid and rifampicin-induced hepatitis is associated with the inhibition of protein synthesis, which may be partially responsible for the fatty liver, though probably not for the necrosis, although it contributes to the disabling of the cell. Accumulation of

Table 4.2.3: Levels of creatinine and glucose in the serum of normal and experimental

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
	Normal Control	Chitosan-Control A	Glucosamine-Control B	Hepatotoxicity C	A+C	B+C
Creatinine	1.01±0.08	1.17±0.06	1.19±0.07	0.63±0.03 ^{a,b,c}	1.14±0.11 ^d	1.13±0.13 ^d
Glucose	88.4±6.31	86.7±6.34	86.5±6.32	64.58±4.95 ^{a,b,c}	77.8±5.68 ^e	77.7±5.63 ^e

Values are expressed as the mean ± SD for six animals. Creatinine and glucose- mg/dl

^a $P < 0.001$ - compared with Group-I

^b $P < 0.001$ - compared with Group-II

^c $P < 0.001$ - compared with Group-III

^d $P < 0.001$ - compared with Group-IV

^e $P < 0.01$ - compared with Group-IV

metabolites of antitubercular drugs may also contribute to the changes in rough endoplasmic reticulum and to the disturbance of protein metabolism. My findings confirm the same pattern, and showed a significant decrease ($P < 0.001$) in protein content in the serum and liver of Group IV, toxicity induced animals as compared to that of Group I normal rats (**Figs 4.2.2. and 4.2.3.**). The rats co-administered with chitosan and glucosamine showed a significant increase in liver and serum protein content as compared to hepatotoxic rats.


4.2.6. Electrophoretic pattern

In the serum of antitubercular drugs induced rats, the level of albumin was decreased while the level of globulin was slightly increased. The SDS-polyacrylamide gel electrophoretic separation of serum proteins showed the significant reduction in the amount of serum protein especially albumin fraction (**Plate 4.2.7.**). Serum albumin level is an index of severity and prognosis in patients with chronic hepatic disease. In patients with acute hepatitis, serum albumin level is slightly depressed whereas globulin level is mildly increased (Devaki *et al.*, 1992). In the present study, albumin/globulin ration in antitubercular drugs-induced rats, was less than the normal control rats. When, the SDS-page electrophoretic pattern of liver (**Pate 4.2.8.**) is considered, similar changes were observed. The albumin and globulin levels were maintained to near normalcy in chitosan and glucosamine administered rats.

4.2.7. Glycoprotein components

Levels of glycoprotein conjugates in the liver were showed in **Table 4.2.4.** Group IV, isoniazid and rifampicin administered rats showed a decrease in the levels of hexose, hexosamine and sialic acid. This is in line with an earlier reported study (Leinweber and Mahrt, 1976). The inhibition of protein synthesis in isoniazid and rifampicin-administered rats will disturb the glycoprotein synthesis. Thus, the glycoprotein conjugates (hexose, hexosamine and sialic acid) were found significantly decreased ($P < 0.001$) in the Group IV rats. Since, chitosan and glucosamine can induce glycosylation, co-administration of these compounds

Fig. 4.2.2: Levels of total protein in the serum of normal and experimental groups of rats.




Values are expressed as the mean \pm SD for six animals.

^a $P < 0.001$ - compared with Group-I, ^b $P < 0.001$ - compared with Group-II, ^c $P < 0.001$ - compared with Group-III,

^d $P < 0.001$ - compared with Group-IV

Fig. 4.2.3: Levels of protein in the liver of normal and experimental groups of rats.



Values are expressed as the mean ± SD for six animals.

^a $P < 0.001$ - compared with Group-I, ^b $P < 0.001$ - compared with Group-II, ^c $P < 0.001$ - compared with Group-III,

^d $P < 0.001$ - compared with Group-IV, ^e $P < 0.01$ - compared with Group-IV




Plate 4.2.7. Electrophoretic pattern of serum protein of normal and experimental groups of rats.

BSA- Bovine Serum Albumin

I- Normal control rats

II- Chitosan control

III- Glucosamine control

IV- Hepatotoxicity induced

V- Chitosan co-administered

VI- Glucosamine co-administered




Plate 4.2.8. Electrophoretic pattern of liver protein of normal and experimental groups of rats.

BSA- Bovine Serum Albumin

- I- Normal control rats**
- II- Chitosan control**
- III- Glucosamine control**
- IV- Hepatotoxicity induced**
- V- Chitosan co-administered**
- VI- Glucosamine co-administered**

Table 4.2.4: Levels of hexose, hexosamine and sialic acid in the liver of normal and experimental groups of rats.

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
	Normal Control	Chitosan-Control A	Glucosamine-Control B	Hepatotoxicity C	A+C	B+C
Hexose	24.7±1.55	28.9±1.68	32.7±2.14	15.6±1.21 ^{a,b,c}	24.3±1.45 ^d	27.5±1.99 ^d
Hexosamine	8.18±0.49	10.7±0.65	12.8±0.73	5.11±0.44 ^{a,b,c}	7.87±0.56 ^d	9.24±0.57 ^d
Sialic acid	0.32±0.02	0.36±0.03	0.34±0.0.3	0.21±0.01 ^{a,b,c}	0.28±0.02 ^d	0.27±0.03 ^e

Values are expressed as the mean ± SD for six animals. Values are expressed in mg/g tissue.

^a $P < 0.001$ - compared with Group-I

^b $P < 0.001$ - compared with Group-II

^c $P < 0.001$ - compared with Group-III

^d $P < 0.001$ - compared with Group-IV

^e $P < 0.01$ - compared with Group-IV

may accelerate the synthesis of hexose, amino sugars and sialic acid. As the result, tissue glycoprotein conjugate levels were brought back to normal level in Group V and Group VI animals.

4.2.8. Lipid peroxidation

Lipid peroxidation of membranes is regulated by the availability of substrate in the form of polyunsaturated fatty acids (PUFA), the availability of inducers such as free radicals and the excited state molecules to initiate propagation, the antioxidant defense status of environment and the physical status of membrane lipids (Pepicelli *et al.*, 2005). Antibiotic therapy can favor free radical production in excess of basal rates. Many antibiotics that depend on bound metals for their activity are able to generate free radicals and cause cellular damage (Doroshov & Hochstein, 1982). In the present study, a significant ($P < 0.001$) increase in the level of lipid peroxides in the serum and liver tissue of antitubercular drugs-treated rats was observed (**Table 4.2.5**). This is in corroboration with an earlier investigation (Saraswathy and Shyamala Devi, 2001), which suggested that the high vulnerability of liver to peroxidative damage is mainly due to a decline in the level of free radicals for scavengers. Free radicals initiate lipid peroxidation of biological membranes. Isoniazid and rifampicin induced hepatitis is due to their biotransformation to reactive metabolites that are capable of binding to cellular macromolecules (Georgeiva *et al.*, 2004). As an alternative to inducing cellular damage by covalent binding, there is evidence that these antitubercular drugs causes cellular damage through the induction of oxidative stress, a consequence of dysfunction of hepatic antioxidant defense system. The role of oxidative stress in the mechanism of isoniazid and rifampicin-induced hepatitis has been reported by Attri *et al.* (2000). My findings confirm the same pattern and show significant increase in the level of lipid peroxidation in the serum and liver tissue of Group IV antitubercular drugs administered rats as compared to that of Group I control rats. The combination of isoniazid and rifampicin treatment in experimental animals enhanced lipid peroxidation, indicating increased oxidative stress in liver (Skakun and Yun, 1992).

Table 4.2.5: Levels of lipid peroxides (LPO) in the serum and liver of normal and experimental groups of rats.

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
	Normal Control	Chitosan-Control A	Glucosamine-Control B	Hepatotoxicity C	A+C	B+C
Serum-LPO	1.43±0.11	1.36±0.09	1.45±0.11	3.42±0.22 ^{a,b,c}	1.57±0.10 ^d	1.86±0.15 ^d
Liver- LPO	0.96±0.07	0.87±0.04	1.02±0.06	1.91±0.165 ^{a,b,c}	0.98±0.05 ^d	1.21±0.06 ^d

Values are expressed as the mean ± SD for six animals. Serum LPO- nmoles of MDA/mi; Liver LPO- nmole MDA/mg protein.

^a $P < 0.001$ - compared with Group-I

^b $P < 0.001$ - compared with Group-II

^c $P < 0.001$ - compared with Group-III

^d $P < 0.001$ - compared with Group-IV

Antioxidants are necessary for preventing the formation of free radicals and they inhibit some of the deleterious actions of reactive oxygen species that damage lipids, DNA and proteins (Eidelman *et al.*, 2002). Co-treatment with chitosan and glucosamine resulted in significant ($P<0.001$) reduction in the level of lipid peroxidation (Shevtsova, 2000) towards near normalcy, as compared with the levels in the Group IV antitubercular drugs-administered rats, establishing the antioxidant nature (Mohur and Cook, 1957) by blocking isoniazid and rifampicin-induced lipid peroxidation.

4.2.9. Glutathione-dependent antioxidant system

Activities of glutathione-dependent antioxidant enzymes [glutathione peroxidase (GPx) and glutathione S-transferase (GST)] and the level of reduced glutathione were significantly ($P<0.001$) lower in the liver tissue of Group IV rats as compared to Group I control animals (**Tables 4.2.6 and 4.2.8**). Glutathione antioxidant system plays a fundamental role in cellular defense against reactive free radicals and other oxidant species. Excessive lipid peroxidation can cause increased glutathione consumption (Onyema *et al.*, 2006), as observed in the present study. The cellular tripeptide, GSH (γ -glutamyl cysteinyl glycine) thwarts peroxidative damage by neutralizing the free radicals. Reduction noticed in the activities of GPx and GST in antitubercular drugs -induced hepatotoxicity condition might be due to decreased availability of their substrate, reduced glutathione. This in turn may lead to increase the accumulation of oxidative free radicals and also enhance the susceptibility of liver cell membrane to peroxidative damage.

In the present study, co-treatment with chitosan and glucosamine significantly prevented the antitubercular drugs-induced aberrations in the level of GSH ($p<0.001$) and the activities of GPx ($P<0.001$) and GST ($P<0.001$, $P<0.05$) in the liver tissue of Group V and Group VI rats as compared to that of Group IV rats. Benassi *et al.* (2006) had earlier reported that chitosan administration could increase the level of GSH in hepatotoxic condition.

4.2.10. Antiperoxidative enzymes

Activities of antiperoxidative enzymes [super oxide dismutase (SOD) and catalase (CAT)] were significantly ($P<0.001$) lower in the liver of antitubercular drugs-administered rats as compared to Group I control rats (**Table 4.2.6**). SOD and CAT, responsible for the destruction of peroxides have a specific role in protecting tissues against oxidative damage. Reduction in the activities of these enzymes in antitubercular drugs-induced hepatotoxicity may lead to the formation of O^{2-} and H_2O_2 , which in turn can form hydroxyl radical (OH^{\cdot}) and bring about a number of reactions harmful to the cellular and subcellular membranes. The generation of free radical scavenging enzymes in antitubercular drugs-induced hepatotoxicity might have exceeded the ability of these free radical scavenging enzymes to dismute the radicals, resulting in peroxidative damage to liver cell membranes.

In this study, co-treatment with chitosan and glucosamine significantly ($P<0.001$) maintained the activities of these antiperoxidative enzymes at near normalcy. They probably did so by their antioxidant action (Xing *et al.*, 2005; Sa'nikova *et al.*, 1990) against isoniazid and rifampicin-induced lipid peroxidation. The unpaired electron present in the hydroxyl free radical, which is mainly responsible for the antitubercular drugs-induced necrotic damage to the hepatocytes, might have been trapped and subsequently dismuted by the antioxidant property of chitosan and glucosamine. Earlier Yan *et al.* (2006) reported that chitosan had a potent reducing effect on the production of free radicals in rats exposed to cytotoxic substance.

4.2.11. Membrane-bound ATPases

There was a significant ($P<0.001$) reduction noticed in the activities of the membrane bound ATPases in the liver of Group IV, antitubercular drugs administered rats as compared to that of Group I control animals (**Table 4.2.7**). This is in accordance with an earlier reported study (Saraswathy *et al.*, 1998), which indicated that isoniazid and rifampicin induced hepatitis is characterized by

Table-4.2.6: Levels of catalase (CAT), glutathione s-transferase (GST), glutathione peroxidase (GPx) and super oxide dismutase (SOD) in the liver of normal and experimental groups of rats.

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
	Normal Control	Chitosan-Control A	Glucosamine-Control B	Hepatotoxicity C	A+C	B+C
CAT	7.33±0.42	7.25±0.39	7.28±0.46	3.05±0.24 ^{a,b,c}	6.84±0.37 ^d	5.46±0.27 ^d
GST	9.63±0.58	10.4±0.61	9.42±0.55	6.29±0.31 ^{a,b,c}	8.77±0.59 ^d	7.23±0.47 ^e
GPx	7.59±0.47	7.76±0.45	7.43±0.52	4.83±0.24 ^{a,b,c}	6.99±0.36 ^d	6.02±0.32 ^d
SOD	6.21±0.55	6.44±0.47	6.18±0.42	3.04±0.18 ^{a,b,c}	5.97±0.46 ^d	4.83±0.39 ^d

Values are expressed as the mean ± SD for six animals.

CAT, μmol of H_2O_2 consumed/min/mg protein; GST, μmol 1-chloro-2,4 dinitrobenzene conjugate formed /min /mg protein; GPx, nmol GSH oxidized/min/mg protein; SOD, one unit of the superoxide dismutase activity is the amount of protein required to give 50% inhibition of epinephrine autooxidation.

^a $P < 0.001$ - compared with Group-I, ^b $P < 0.001$ - compared with Group-II, ^c $P < 0.001$ - compared with Group-III,

^d $P < 0.001$ - compared with Group-IV, ^e $P < 0.05$ - compared with Group-IV

Table 4.2.7: Activities of membrane bound ATPases in the liver of normal and experimental groups of rats.

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
	Normal Control	Chitosan-Control A	Glucosamine-Control B	Hepatotoxicity C	A+C	B+C
Total ATPase	3.34±0.18	3.45±0.15	3.23±0.22	2.08±0.11 ^{a,b,c}	3.08±0.19 ^d	2.67±0.14 ^d
Ca ²⁺ -ATP	0.74±0.05	0.69±0.05	0.78±0.04	0.51±0.02 ^{a,b,c}	0.67±0.04 ^d	0.63±0.05 ^d
Mg ²⁺ -ATP	1.42±0.11	1.49±0.14	1.35±0.12	1.10±0.07 ^{a,b,e}	1.31±0.11 ^f	1.28±0.10 ^f
Na ⁺ -K ⁺ ATP	1.18±0.08	1.27±0.09	1.1±0.07	0.47±0.03 ^{a,b,c}	1.12±0.07 ^d	0.86±0.07 ^d

Mean ± SD for six animals in each group. Values are expressed as μ moles of Pi liberated per mg protein per hour.

^a P<0.001- compared with Group-I

^b P<0.001- compared with Group-II

^c P<0.001- compared with Group-III

^d P<0.001- compared with Group-IV

^e P<0.01- compared with Group-III

^f P<0.05- compared with Group-IV

Table 4.2.8: Levels of total thiols and free thiol (GSH) in the liver of normal and experimental groups of rats.

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
	Normal Control	Chitosan-Control A	Glucosamine-Control B	Hepatotoxicity C	A+C	B+C
Total thiols	398±27.4	412±24.5	385±28.2	237±17.5 ^{a,b,c}	353±23.7 ^d	319±25.7 ^d
Free thiol (GSH)	4.84±0.32	5.18±0.34	4.97±0.31	3.38±0.28 ^{a,b,c}	4.95±0.33 ^d	4.65±0.32 ^d

Mean ± SD for six animals in each group. Values are expressed as nmoles/g wet tissue.

^a $P < 0.001$ - compared with Group-I

^b $P < 0.001$ - compared with Group-II

^c $P < 0.001$ - compared with Group-III

^d $P < 0.001$ - compared with Group-IV

a severe derangement of subcellular metabolism and structural alterations in liver cell membrane. Since membrane-bound ATPases are lipid dependent as well as –SH dependent enzymes, alterations in membrane lipid may lead to changes in membrane fluidity, which in turn affects concentration of intracellular ions and conservation of transmembrane electrochemical gradient. The functional state of liver depends on its active Na^+ , K^+ sequestering property. Earlier Tasduq *et al.* (2005) have reported that the appearance of antitubercular drugs induced hepatic injury is dependent on intracellular Ca^{2+} concentrations. Isoniazid and rifampicin intoxication has been previously reported to be associated with alteration in membrane-bound ATPases (Skakun and Tabachuk, 1992). Co-administration of chitosan and glucosamine restored the activities of membrane bound ATPases at near normal level. This could be due to the ability of chitosan and glucosamine to protect the ‘SH’ groups from oxidative damage though inhibition of peroxidation of membrane lipids. The membrane stabilizing property of chitosan is already reported by Filipovic-Grcic *et al.* (2001).

4.2.12. Hepatic thiols

The levels of membrane protective thiols were found to decrease significantly ($P < 0.001$) in antitubercular drugs-administered rats (Table 4.2.8). The remarkable loss in total, Ca^{2+} and Na^+K^+ ATPase activities found in Groups IV animals may also be due to the deteriorations of SH because of increased peroxidative damage caused by isoniazid and rifampicin administration. Co-administration of chitosan and glucosamine was found to restore the levels of thiols in Group V and Group VI animals. This is due to the free radical scavenging property of these compounds

4.2.13. Lipid components

Tables 4.2.9 and 4.2.10 show the levels of lipid components in serum and liver of normal and experimental groups of rats respectively. Serum HDL and LDL cholesterol levels were shown in Fig. 4.2.4. Significant ($P < 0.001$) increases in

Table 4.2.9: Levels of lipid components in the serum or normal and experimental groups of rats

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
	Normal Control	Chitosan-Control A	Glucosamine-Control B	Hepatotoxicity C	A+C	B+C
Cholesterol	54.5±0.48	48.1±0.24	58.7±0.31	78.9±0.62 ^{a,b,c}	56.1±0.27 ^d	59±0.29 ^d
Triglycerides	47.4±0.36	40.6±0.26	48.1±0.28	95.8±0.76 ^{a,b,c}	54.9±0.34 ^d	55.3±0.35 ^d
Free fatty acids	10.2±0.09	8.34±0.51	9.87±0.58	18.1±0.12 ^{a,b,c}	10.5±0.98 ^d	11.4±1.2 ^d
Phospholipids	108±9.8	94.2±8.72	98.3±8.91	128±10.40 ^{b,c,e}	102±9.16 ^d	107±9.54 ^f

Values are expressed as Mean ± SD for six animals in each group. Cholesterol, triglycerides, free fatty acids and phospholipids = mg/dl

^a P<0.001- compared with Group-I, ^b P<0.001- compared with Group-II, ^c P<0.001- compared with Group-III,

^d P<0.001- compared with Group-IV, ^e P<0.01- compared with Group-I, ^f P<0.01- compared with Group-IV

Table 4.2.10: Levels of lipid components in the liver normal and experimental groups of rats

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
	Normal Control	Chitosan-Control A	Glucosamine-Control B	Hepatotoxicity C	A+C	B+C
Cholesterol	9.18±0.81	8.05±0.54	8.71±0.58	15.4±0.98 ^{a,b,c}	10.4±0.87 ^d	10.5±0.88 ^d
Triglycerides	18.5±1.25	15.3±0.95	18.8±1.23	29.9±1.75 ^{a,b,c}	18.2±1.29 ^d	20.3±1.58 ^d
Free fatty acids	0.94±0.05	0.83±0.04	1.02±0.05	1.73±0.11 ^{a,b,c}	0.88±0.04 ^d	1.1±0.06 ^d
Phospholipids	27.3±1.02	28.5±2.15	27.2±1.03	24.2±1.84 ^{e,f,g}	27.6±2.03 ^h	27.4±2.01 ^h

Values are expressed as Mean ± SD for six animals in each group. Cholesterol, triglycerides, free fatty acids and phospholipids = mg/g tissue

^a $P < 0.001$ - compared with Group-I, ^b $P < 0.001$ - compared with Group-II, ^c $P < 0.001$ - compared with Group-III,


^d $P < 0.001$ - compared with Group-IV, ^e $P < 0.01$ - compared with Group-II, ^f $P < 0.05$ - compared with Group-I,

^g $P < 0.05$ - compared with Group-III, ^h $P < 0.05$ - compared with Group-IV

the levels of cholesterol, triglycerides and free fatty acids were observed in the serum and liver of Group IV, isoniazid and rifampicin-administered rats as compared to that of Group I control rats. A slight decline in the level of HDL-cholesterol with concomitant rise in LDL-cholesterol level was also noted in Group IV antitubercular drugs-administered rats (**Fig. 4.2.4.**). The rats co-administered with chitosan and glucosamine showed significantly near normal levels of lipid components in serum and liver of Group V and Group VI rats as compared to that of Group IV hepatotoxicity induced rats, indicating the antilipidemic effect of chitosan. A significant rise in the level of phospholipids was noted in serum ($P<0.01$) of Group IV antitubercular drugs-administered rats. This was paralleled by a slight reduction ($P<0.05$) in the level of phospholipids in liver tissue of Group IV rats. Chitosan co-administration resulted in the prevention of antitubercular drugs-induced phospholipids depletion in the liver of experimental groups of rats. It also exerted a significant ant ilipid peroxidative effect by blocking the induction of isoniazid and rifampicin-mediated lipid peroxidation in serum and liver of Group IV rats. The rats administered with chitosan alone did not show any adverse effects, indicating that chitosan is non-toxic.

The major disorder encountered in antitubercular drugs-induced hepatitis is fatty accumulation in the liver, which develops either due to excessive supply of lipids to the liver or interference with lipid deposition. The pathogenesis is multifactorial, reflecting complex biosynthetic, enzymatic and catabolic derangement in lipoprotein metabolism. In the present study, the levels of total cholesterol and LDL-cholesterol were significantly ($P<0.001$) higher in Group III isoniazid and rifampicin-administered rats as compared to that of normal control animals, indicating the antitubercular drugs-induced hypercholesterolemic condition. Also the level of HDL-cholesterol was slightly reduced in Group III rats. This is in line with an earlier study (Gibson *et al.*, 1995). Increased cholesterol levels in the liver might due to increased uptake of LDL from the blood by the tissues (Kissler *et al.*, 2005). The abnormal cholesterol deposition is favored by the dangerous tendency of cholesterol to passive exchange between the plasma lipoproteins and the cell membranes (Brown and Goldstein, 1986).

Fig. 4.2.4: Levels of HDL-cholesterol and LDL-cholesterol in the serum of normal and experimental groups of rats.



Values are expressed as the mean ± SD for six animals.

^a $P < 0.001$ - compared with Group-I, ^b $P < 0.001$ - compared with Group-II, ^c $P < 0.001$ - compared with Group-III,

^d $P < 0.001$ - compared with Group-IV

In the present study, co-administration with chitosan significantly ($P<0.001$) reduced the antitubercular drugs-induced elevation in the levels of total cholesterol and LDL-cholesterol in Group IV rats as compared to that of Group III rats. Also the level of HDL-cholesterol maintained at near normalcy. Earlier Yao and Chiang (2002) have shown that chitosan intake is effective in lowering the plasma total cholesterol, VLDL cholesterol and LDL cholesterol levels in experimental animals. In the present study, a slight reduction in the level of total cholesterol and LDL-cholesterol were also observed in Group II chitosan and glucosamine fed normal animals, establishing the hypocholesterolemic property of them. The hepatoprotective effect of chitosan and glucosamine is probably related to its ability to inhibit the lipid accumulation in the liver tissue by the antilipidemic property (Xing *et al.*, 2005).

The levels of triglycerides were significantly ($P<0.001$) higher in serum and liver tissue of Group III rats as compared to Group I control animals (**Tables 4.2.9 and 4.2.10**). Increased lipolysis of depot triglycerides liberates free fatty acids from adipose tissue stores (Kruger *et al.*, 1967; Stenberg, 1976) and the free fatty acids liberated by the adipose tissue are also taken up by the liver tissue, leading to the hypertriglyceridemic condition. In the present study, a significant ($P<0.001$) reduction was noticed in the levels of triglycerides and free fatty acids in serum and liver of Group IV chitosan and glucosamine administered rats as compared to that of Group III isoniazid and rifampicin-induced hepatotoxic rats, indicating the hypolipidemic nature of chitosan and glucosamine. The hypolipidemic property of chitosan has already been reported in high fat diet fed experimental animals (Xing *et al.*, 2005).

The levels of phospholipids were significantly ($P<0.01$) decreased in the liver tissue of Group III isoniazid and rifampicin-administered rats as compared to that of normal control animals. This is in accordance with earlier study (Karthikeyan, 2005), which indicates that hepatic injury related alterations in lipid composition of liver tissue appears to occur due to the destruction of hepatocytes. The intracellular calcium concentration has been reported to rise in antitubercular drugs-induced hepatitis (Tasduq *et al.*, 2005). This intracellular Ca^{2+} is an

inducer of phospholipase A₂, which degrades membrane phospholipids. The increased peroxidation of membrane phospholipids releases free fatty acid by the action of phospholipase A₂. Hence, the significant ($P<0.001$) elevation noticed in free fatty acid content in isoniazid and rifampicin-administered rats might be due to enhanced breakdown of membrane phospholipids. Recent studies (Karthikeyan, 2005) indicate that accelerated degradation of membrane phospholipids is very likely the biochemical basis for the antitubercular induced hepatocellular injury.

In the present study, the co-administration of chitosan significantly ($P<0.05$) reduced the antitubercular drugs-induced degradation of membrane phospholipids in liver, establishing its membrane stabilizing effect. It probably did so by preventing changes in fatty acid composition and inhibiting the peroxidative deterioration of hepatocellular membranes. It is possible that stabilization of hepatocellular membranes might prolong the viability of hepatocytes.

4.2.14. Fatty acid composition

Table 4.2.11 shows the fatty acid composition of the liver of normal and experimental groups of rats. Fatty Administration of isoniazid and rifampicin leads to their biotransformation to toxic metabolites like ROS (reactive oxygen species) and the prime target of ROS is PUFA (Georgieva *et al.*, 2004). Depletion of PUFA alters the fatty acid composition of tissues in isoniazid and rifampicin intoxicated rats. The levels of all fatty acids were found to increase except arachidonic acid in Group IV, hepatotoxic animals. Arachidonic acid is released from cell membrane phospholipids by the effect of calcium dependent phosphatases (PL) A₂. Cyclooxygenase products of arachidonic acid are present in much higher concentrations in inflamed tissues than in healthy tissues (Emken *et al.*, 1997). Administration of isoniazid and rifampicin may increase the membrane PLA₂ activity, liberating arachidonic acid from phospholipids. Thus the arachidonic acid levels in the antitubercular drugs induced rats are lowered than the control animals. This decrease may also be due to the channeling of arachidonic acid for the production of prostaglandins. Arachidonic acid

Table 4.2.11: Fatty acid composition of the liver of normal and experimental groups of rats

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
	Normal Control	Chitosan-Control A	Glucosamine-Control B	Hepatotoxicity C	A+C	B+C
Palmitic acid	12.52±1.24	10.58±1.21	11.63±1.39	29.24±2.89 ^{a,b,c}	17.91±1.26 ^d	19.26±1.98 ^e
Palmitoleic acid	16.34±1.58	18.62±1.71	17.46±1.82	9.53±1.06 ^{a,b,c}	14.53±1.21 ^d	11.39±1.23 ^e
Stearic acid	11.23±1.13	12.65±1.19	11.84±1.16	19.57±1.62 ^{a,b,c}	12.36±1.15 ^d	13.54±1.31 ^e
Oleic acid	11.16±1.25	9.87±1.08	10.08±1.10	18.68±1.38 ^{a,b,c}	12.62±1.20 ^d	13.81±1.19 ^e
Arachidonic acid	2.87±0.05	1.98±0.03	2.11±0.04	4.89±0.07 ^{a,b,c}	3.12±0.03 ^d	3.56±0.04 ^f
Eicosapentaenoic acid	22.86±1.98	23.75±2.02	23.02±2.03	15.64±1.52 ^{a,b,c}	20.86±1.19 ^d	19.46±1.12 ^f
Docosapentaenoic acid	15.49±1.38	14.72±1.36	13.88±1.41	26.85±2.56 ^{a,b,c}	16.54±1.53 ^d	17.62±1.28 ^e

Values are expressed as Mean ± SD for six animals in each group. Values are expressed in wt %

^a P<0.001- compared with Group-I, ^b P<0.001- compared with Group-II, ^c P<0.001- compared with Group-III,

^d P<0.001- compared with Group-IV, ^e P<0.001- compared with Group-V, ^f P<0.05- compared with Group-IV

hydrolysed from phospholipids is used for the production of eicosanoids (Holtzman *et al.*, 1991). Chitosan and glucosamine administration to antitubercular drugs induced rats inhibit the release of arachidonic acid in the liver, which may be because chitosan and glucosamine inhibit the activity of PLA₂.

4.2.15. Mineral composition

Mineral composition of the liver of normal and experimental groups of rats were given in **Fig. 4.2.5.** and **Table 4.2.12.** The intracellular concentration of sodium and potassium were regulated by a transport system in the cell membrane. This converts chemical energy by the hydrolysis of ATP to drive the vectorial movement of sodium out of the cell and potassium into the cell against an electrochemical gradient (Skou, 1971). Active calcium transport and resultant low calcium concentration are the necessary conditions for active sodium/potassium ionic transport. Since, sodium and calcium are thought to be competitive at a number of membrane sites, it seems likely that a high concentration of Ca²⁺ in liver cells of ischemic rats would compete with sodium specific sites at the inner surface of the membrane (Vincenzi, 1971) and this may lead to decrease in sodium content. Also, failure of sodium pump results in a depletion of liver sodium and rise in potassium concentration was observed in the present study. This leads to hyponatremia and hyperkalemia, which are the most common electrolyte abnormalities in hepatotoxic condition.

Isoniazid and rifampicin administration increased the level of calcium in the liver tissue. Elevation in Ca²⁺ is known to be involved in the initiation and progression of hepatocellular damage (Bellemo and Orrenius, 1985). Calcium is a messenger of great importance to cells. Antitubercular drugs administration could maximally increase Ca²⁺ influx in whole liver tissue. Since plasma membrane Ca²⁺-ATPase extrudes Ca²⁺ from the cytoplasm of all the cells (Howard *et al.*, 1994) blocking of intracellular Ca²⁺-ATPase under the condition of isoniazid and rifampicin induced hepatotoxicity results in fast cellular Ca²⁺ accumulation. Thus influx of calcium into the cell is uncontrolled under antitubercular drugs induced toxicity,

Table 4.2.12: Levels of calcium, magnesium and zinc in the liver of normal and experimental groups of rats

Parameters	Group I	Group II	Group III	Group IV	Group V	Group VI
	Normal Control	Chitosan-Control A	Glucosamine-Control B	Hepatotoxicity C	A+C	B+C
Calcium	302±6.32	299±5.84	301±5.68	355±10.36 ^{a,b,c}	312±9.63 ^d	314±9.81 ^e
Magnesium	291±5.12	288±5.03	290±5.11	338±9.25 ^{a,b,c}	301±7.94 ^d	304±7.82 ^e
Zinc	92±3.56	90±3.49	91±3.51	144±5.32 ^{a,b,c}	98±4.68 ^d	100±4.85 ^e

Mean ± SD for six animals in each group. Values are expressed as mg/g tissue

^a P<0.001- compared with Group-I


^b P<0.001- compared with Group-II

^c P<0.001- compared with Group-III

^d P<0.001- compared with Group-IV

^e P<0.01- compared with Group-IV

Fig. 4.2.5: Levels of sodium and potassium in the liver of normal and experimental groups of rats.



Values are expressed as the mean ± SD for six animals.

^a $P < 0.001$ - compared with Group-I, ^b $P < 0.001$ - compared with Group-II, ^c $P < 0.001$ - compared with Group-III,

^d $P < 0.001$ - compared with Group-IV, ^e $P < 0.01$ - compared with Group-IV

leading to the disturbance in equilibrium between intracellular and extra cellular calcium concentrations. The level of Mg^{2+} was also found to be raised significantly in the hepatotoxic group. The intracellular concentration of calcium regulates the activity of the Mg^{2+} . Bironaite and Ollinger (1997) reported that peroxidation can influence the alteration of Mg^{2+} .

Co-administration of chitosan and glucosamine significantly ($P < 0.001$) prevented the isoniazid and rifampicin induced alterations in the levels of these minerals to the near normal level. It may be done by protecting the plasma membrane bound ATPases from the free radical attack by their free radical scavenging capacity. An earlier report (Wada *et al.*, 1997) supports this finding, that chitosan can reduce the whole-body retention of Calcium in rats.

In conclusion, the results of the present study indicate that the co-treatment with chitosan and glucosamine prevent antitubercular drugs-induced hepatotoxicity in rats. The overall hepatoprotective effect of chitosan and glucosamine is probably due to its membrane stabilizing action, or to a counteraction of free radicals by its antioxidant nature, or to its ability to maintain near to the normal status the activities of the free radical scavenging enzymes and the level of reduced glutathione, which protects cell membrane against peroxidative damage by decreasing lipid peroxidation and strengthening the liver cell membrane.

5. SUMMARY AND CONCLUSION

The protective effect of chitosan and glucosamine against ibuprofen-induced gastric ulcer was studied. Administration of ibuprofen had brought about the ulcerated mucosal lesions in the experimental rats. It is evident by the increased acidity, number of lesions and decreased peptic activity. The gross pathological and histopathological observations also confirmed the mucosal tissue damage.

Pre-oral treatment with chitosan and glucosamine significantly decreased the ibuprofen induced gastric lesions and acidity and brought back the peptic activity to the normal level. This shows the cytoprotective effect of these compounds. Chitosan and glucosamine might be neutralized the increased acid secreted by the stomach due to ibuprofen irritant.

The levels of antioxidant enzymes, antiperoxidative enzymes, reduced glutathione, lipid peroxidation and membrane bound ATPases were also altered by the ibuprofen intake. Chitosan and glucosamine pre-treatment maintained these levels to near normalcy. The results of the present study indicate that the pre-oral administration of chitosan and glucosamine maintain near to the normal status the activities of the mucosal antioxidant enzymes and the level of GSH, which protect mucosa against oxidative damage by decreasing the lipid peroxidation and strengthening the mucosal barrier, and which are the first line of defense against exogenous and endogenous ulcerogenic agents. Pre-treatment with chitosan and glucosamine significantly maintained the activities of membrane bound ATPases by the counteraction of ibuprofen-induced free radicals by the free radical scavenging property of chitosan and glucosamine.

Administration of ibuprofen deteriorated protein and glycoprotein levels due to the corrosion of gastric mucosa, resulting in the disruption and disintegration of gastric mucosal cells. This was reflected in the electrophoretic pattern also. Oral pre-treatment with chitosan and glucosamine maintained these levels. This may be because; the incorporation of glucosamine units into mucosal glycoprotein might have resulted in strengthening of the mucosal barrier.

Histopathological study showed that ibuprofen induction was characterized by severe cell necrosis, which is normalized by the administration of chitosan and glucosamine showing their cytoprotectivity. The level of sodium was decreased and potassium, calcium, magnesium and zinc were increased due to ibuprofen administration. The accumulation of intracellular calcium levels also play an important role in the DNA fragmentation and ulcer induced by ibuprofen. Increased peroxidation may be the cause of increment in the magnesium level. Chitosan and glucosamine pre-treatment maintained the levels of these minerals to the normal level. Changes in the amino acid levels brought about by the ibuprofen induced cell damage were also restored to normal level by the pre-oral administration of chitosan and glucosamine.

Observations in this study indicate that the oral pre-treatment of chitosan and glucosamine can prevent ibuprofen-induced peptic ulcer in rats. The overall antiulcerogenic activity of chitosan and glucosamine is probably related to the ability to neutralize the hydrochloric acid secreted into the stomach, and to the maintenance of GSH level by decreasing lipid peroxidation and to the capability to strengthen the mucosal barrier by increasing mucosal glycoprotein synthesis and the free radical scavenging property. On comparison, glucosamine was found to have better antiulcerogenic property than chitosan.

The protective effect of chitosan and glucosamine against antitubercular drugs-induced hepatotoxicity was also studied. Administration of isoniazid and rifampicin, most widely used antitubercular drugs, caused severe toxicity to liver. It increased the levels of diagnostic marker enzymes in the serum. Enhanced susceptibility of hepatocytes cell membrane to the isoniazid and rifampicin-induced peroxidative damage might have resulted in increased release of these diagnostic marker enzymes into the systemic circulation. In cases of liver damage with hepatocellular lesions and parenchymal cell necrosis, these marker enzymes are released from the damaged tissues into the blood stream. As a result the levels of liver ALT and AST were decreased in the antitubercular drugs-induced rats. Co-administration of chitosan and glucosamine maintained the levels of diagnostic marker enzymes in the serum and liver to near normalcy.



They probably did so by preventing the antitubercular drugs induced necrotic damage by the membrane stabilizing action normalizing the protein metabolism.

Histopathological investigations showed severe vacuolation and degeneration of hepatocytes in the three zones of lobules. Membrane disintegration with intense vacuolation accompanied with cytoplasmic rarefaction resulting in loss of polyhedral structure in the liver of rat treated with antitubercular drugs. Treatment with chitosan and glucosamine caused reversal of such pathology with no evidence of necrotic areas.

The oral administration of antitubercular drugs led to periportal infiltration of inflammatory cells, disruption of lobular architecture, and liver cell necrosis and causes severe disturbance of RNA and protein metabolism. Accumulation of metabolites of antitubercular drugs may also contribute to the disturbance in protein metabolism. The eletrophoretic pattern of hepatotoxic rats was also found disturbed especially in albumin fraction. Drug induced hepatotoxicity is associated with failure to maintain serum albumin level whereas serum globulins, particularly γ -globulin, which is formed in the reticuloendothelial system are increased. In hepatotoxic condition, due to the failure of the liver to convert amino acids and ammonia to urea, a significant decrease in urea was observed. The inhibition of protein synthesis in isoniazid and rifampicin-administered rats disturbed the glycoprotein synthesis. Thus, the glycoprotein conjugates (hexose, hexosamine and sialic acid) were found decreased in hepatotoxic animals. Alterations in all these factors were maintained to near normal level by chitosan and glucosamine co-administration. Since, chitosan and glucosamine were N-acetyl sugars they induced glycosylation, which helped glycoprotein synthesis. These muco-compounds can decrease the accumulation of reactive metabolites of isoniazid and rifampicin and balance protein metabolism.

Administration of isoniazid and rifampicin was characterized by increased lipid peroxidation and decreased levels of glutathione-dependent antioxidant enzymes, reduced glutathione and antiperoxidative enzymes. Co-administration of chitosan and glucosamine along with antitubercular drugs brought back the

levels of these enzymes, lipid peroxidation and reduced glutathione to normal status. The antioxidant property of chitosan and glucosamine prevented the formation of free radicals and they inhibited some of the deleterious actions of reactive oxygen species that damage lipids, DNA and proteins

The antitubercular drugs-induced hepatitis is characterized by the fatty accumulation in the liver, which develops either due to excessive supply of lipids to the liver or interference with lipid deposition. The levels of total cholesterol and LDL-cholesterol were increased whereas HDL-cholesterol was decreased in antitubercular drugs induced animals. Increased cholesterol levels in the liver might due to increased uptake of LDL from the blood by the tissues. The abnormal cholesterol deposition is favored by the dangerous tendency of cholesterol to passive exchange between the plasma lipoproteins and the cell membranes. Animals co-administered with chitosan and glucosamine maintained the levels of total, HDL and LDL cholesterol levels to the near normal level. The alterations in triglycerides, free fatty acids and phospholipids in the hepatotoxic rats were also normalized by chitosan and glucosamine co-administration. This may be due to their ability to inhibit the lipid accumulation in the liver tissue by the antilipidemic property and membrane stabilizing effect.

Chitosan and glucosamine co-administration rectified the deviations, caused by the antitubercular drugs administration, in the levels of membrane bound ATPases, hepatic thiols, minerals and fatty acid components.

From all the results obtained in the present study, it can be concluded that co-administration of chitosan and glucosamine can effectively prevent the isoniazid and rifampicin induced hepatotoxicity in rats. Comparatively, chitosan was found to have better results than glucosamine in alleviating the hepatic disorders.

6. REFERENCES

- Abul, K. N., Pillai, K.K., Pal, S.N., Aqil, M. Free radical scavenging and hepatoprotective activity of jigrine against galactosamine induced hepatopathy in rats. *J. Ethnopharmacol.* 2005; 97: 521-525.
- Agnihotri, S.A., Aminabhavi, T.M. Formulation and evaluation of novel tableted chitosan microparticles for the controlled release of clozapine. *J. Microencapsul.* 2004 ;21: 709-18
- Agrawal, N.M. Epidemiology and prevention of non-steroidal anti-inflammatory drug effects in the gastrointestinal tract. *Brit. J. Rheumatol.* 1995; 34: 5-10.
- Allan, G.G., Altman, L.C., Bensinger, R.E., Ghosh, D.K., Hirabayashi, Y., Neogi, A.N., Neogi,S. In *Chitin, Chitosan and Related Enzymes*, Academic Press, Inc. Orland, 1984, pp 119-133.
- Allan, G.G., Fox, J.R., Crosby, G.D. and Sarkanen, K.V. In *Sixth Fundamental Research Symposium*. Technical Division of the British Paper and Board Industry Federation. William Clowe, London. 1977; 75-79.
- Allan, G.G., Fox, J.R., Kong, N. In Proc. First Int. Conf. Chtin/Chitosan, MIT Cambridge, 1978; pp 64-78.
- Allan, G.G., Friedhoff, J.F., Koppela M, Laine, J.E. and Powell, J.C. *ACS. Symp. Ser* 1975; 10: 172-180.
- Allan,G.G., Crosby,G.D., Lee,J.H., Miller, N.L., Reif, W.H. In *Proc. Symp. Man-made Polymers in Paper making*, Helsinki, 1972; pp 85-96
- Allen, A., Newton, J., Oliver, L., Jordan, N., Strugala, V., Pearson, J.P., Dettmar, P.W. Mucus and H. pylori. *J. Physiol. Pharmacol.* 1997; 48: 297-305.
- Amagase, K., Yokota, M., Tsukimi, Y., Okabe, S. Characterization of "unhealed gastric ulcers" produced with chronic exposure of acetic acid ulcers to indomethacin in rats. *J. Physiol. Pharmacol.* 2003; 54: 349-360.
- Amr, A., Alaa A.H.,. Oxidative stress mediates drug-induced hepatotoxicity in rats: a possible role of DNA fragmentation. *Toxicology* 2005; 208: 367–375.
- Anandan, R., Rekha, R.D., Devaki, T. Protective effect of *Picrorhiza kurroa* on mitochondrial glutathione antioxidant system in D-galactosamine–induced hepatitis in rats. *Curr. Sci.* 1999; 76: 1543-1545.

- Anandan, R., Suseela Mathew, Viswanathan Nair, P.G. Antiulcerogenic effects of chitin and chitosan on mucosal antioxidant defense system in HCl-ethanol induced ulcer in rats. *J. Pharm. Pharmacol.* 2004; 56: 265-269.
- Angell, M., Kassirer, J. P. Alternative medicine — the risks of untested and unregulated remedies. *N. Engl. J. Med.* 1998; 339: 839-841
- Anson, M.L. The estimation of pepsin, trypsin, papain and cathepsin with haemoglobin. *J. Gen. Physiol.* 1938; 22: 79-89
- AOAC. *Official Methods of Analysis*, 13th edtn., Association of Official Analytical Chemists, Washington, DC, USA., 1980.
- AOAC. *Official Methods of Analysis*, 9th edtn., Association of Official Analytical Chemists, Washington, DC, USA., 1975.
- Argentieri, D.C., Ritchie, D.M., Ferro, M.P., Kirchner, T., Wachter, M.P., Anderson, D.W., Rosenthale, M.E., Capetola, R.J. Tepoxalin: a dual cyclooxygenase/5-lipoxygenase inhibitor of arachidonic acid metabolism with potent anti-inflammatory activity and a favorable gastrointestinal profile. *J. Pharmacol. Exp. Ther.* 1994; 271: 1399-1408.
- Attri, S., Rana, S.V., Vaiphei, K., Sodhi, C.P., Katyal, R., Goel, R.C., Nain, C.K., Singh, K., 2000. Isoniazid and rifampicin induced oxidative hepatic injury protection by N-acetylcysteine. *Hum. Exp. Toxicol.* 19, 517-522.
- Austin, P.R., Brine, C.J., Castle, J.E., Zikakis, J.P. *Science* 1981; 212:749-753
- Balasubramanian, T., Somasundaram, M., Felix, A.J. Taurine prevents ibuprofen-induced gastric mucosal lesions and influences endogenous antioxidant status of stomach in rats. *Sci. World J.* 2004; 4: 1046-1054.
- Barbara, K. Membrane-based processes performed with use of chitin/chitosan materials. *Sep. Purif. Technol.* 2005; 41: 305-312.
- Bardhan, K.D., Ahlberg, J., Hislop, W.S., Lindholmer, C., Long, R.G., Morgan, A.G., Sjostedt, S., Smith, P.M., Stig, R., Wormsley, K.G. Rapid healing of gastric ulcers with lansoprazole. *Aliment. Pharmacol. Ther.* 1994; 8: 215-220.
- Barletter, G.R. *J. Biol. Chem.* 1959; 234: 466.
- Bellemo, G., Orrenius, S. Altered thiol and calcium homeostasis in oxidative hepatocellular injury. *Hepatology* 1985; 5: 876–82.
- Benassi, J.C., Laus, R., Geremias, R., Lima, P.L., Menezes, C.T., Laranjeira, M.C., Wilhelm-Filho, D., Favere, V.T., Pedrosa, R.C., 2006. Evaluation of remediation of coal mining wastewater by chitosan microspheres using biomarkers. *Arch. Environ. Contam. Toxicol.* 50, 23-30.

- Billi de Catabbi, S.C., Faletti, A., Fuentes, F., San Martin de Viale, L.C., Cochon, A.C. Hepatic arachidonic acid metabolism is disrupted after hexachlorobenzene treatment. *Toxicol. Appl. Pharmacol.* 2005 ; 204:187-95
- Bironaite, D., and Ollinger, K. Hepato toxicity of rhein involves impairment of mitochondrial functions. *Chem. Biol. Interact.* 1997; 103: 35-50.
- Black, M., Mitchell, J.R., Hyman, P.D. Isoniazid-associated hepatitis in 114 patients. *Gastroenterology* 1975; 69: 289-302.
- Blower, A.L. Considerations for nonsteroidal anti-inflammatory drug therapy: Safety. *Scand. J. Rheumatol.* 1996; 25: 13-26.
- Bonting, S.L. Sodium potassium activated adenosine triphosphatase and carbon transport. In: Membrane and Iron Transport, Vol. 1, (ed Bittar EE), Wiley-Interscience, London 1970; pp 257-363.
- Boxenbaum, H.G., Riegelman, S. Determination of isoniazid and metabolites and biological fluids. *J. Pharm. Sci.* 1974; 63: 1191-1197.
- Braham, R. The effect of glucosamine supplementation on people experiencing regular knee pain. *Br. J. Sports Med.* 2003; 37: 45-49.
- Brown, M.S., Goldstein, J.L. A receptor mediated pathway for cholesterol homeostasis. *Science* 1986; 232, 34 -47.
- Burstein, M., Scholnick, H.R. Precipitation of chylomicron and very low density lipoprotein from human serum with sodium lauryl sulphate. *Life Sci.* 1972; 11:177-184.
- Campbell, F.L. *Ann. Ent. Soc. Amer.* 1929; 22: 395-402.
- Canapp, S.O. Jr, McLaughlin, R.M. Jr, Hoskinson, J.J., Roush, J.K., Butine, M.D., Scintigraphic evaluation of dogs with acute synovitis after treatment with glucosamine hydrochloride and chondroitin sulfate. *Am. J. Vet. Res.* 1999; 60: 1552-7
- Cantoni, L., Valaperta, R., Ponsoda, X., Castell, J.V., Barelli, D., Rizzardini, M., Mangolini, A., Hauri, L., Villa, P. Induction of hepatic hem oxygenase-1 by diclofenac in rodents: role of oxidative stress and cytochrom P-450 activity. *Hepatology* 2003, 38; 776-783.
- Castell, J.V., Gomez, M.J., Miranda, M.A., Morera, I.M. Photolytic degradation of ibuprofen. Toxicity of the isolated photoproducts on fibroblasts and erythrocytes. *Photochem. Photobiol.* 1987; 46: 991-6.
- Chi, C.L., Wing S.L., Kwok, C.C., Cheuk, M. T., Wing, W.Y., Chi, K.C., Man, Y.W. Initial experience on Rifampicin and Pyrazinamide vs Isoniazid in the

treatment of Latent Tuberculosis Infection among patients with silicosis in Hong Kong. *Chest* 2003; 124; 2112-2118.

Corinne, B., Lucio, R., Franchimont, P. Stimulation of proteoglycan production by glucosamine sulfate in chondrocytes isolated from human osteoarthritic articular cartilage in vitro. *Osteoarthr. Cartilage* 1998; 6; 427-134.

Crupi, V., Majolino, D., Migliardo, P., Mondello, M.R., Germano, M.P., Pergolizzi, S. FT-IR molecular evidence of liver damage by chemical agents. *Vibr. Spectrosc.* 2001; 25: 213-222.

Cvetkoric, D., Begic-Janeva A., Protic S. *Lagosl Physiol Pharmacol Acta.* 1996, 32: 17-24

Daren, A., Scroggie, M.D., Allison Albright, M.D. Mark, D., Harris, M.D. The effect of glucosamine-chondroitin supplementation on glycosylated hemoglobin levels in patients with type 2 diabetes mellitus: A placebo-controlled, double-blinded. randomized clinical trial. *Arch. Intern. Med.* 2003; 163:1587-1590.

Davenport, H.W. Salicylate damage to the gastric mucosal barrier. *N. Engl. J. Med.* 1967; 276: 1307-1312.

Davies, D.M. *Textbook of adverse drug reactions*, 2nd ed., Oxford University Press, New York, 1981; pp 136-138

Dawe, S., Lockwood, D.N., Creamer, D. A case of post-partum borderline tuberculoid leprosy complicated by a median nerve abscess, peptic ulceration and rifampicin-induced haemolytic renal failure. *Lepr. Rev.* 2004; 75:181-187.

Dembinski, A., Warzecha, Z., Ceranowicz, P., Brzozowski, T., Dembinski, M., Konturek, S.J., Pawlik, W.W. Role of capsaicin-sensitive nerves and histamine H1, H2, and H3 receptors in the gastroprotective effect of histamine against stress ulcers in rats. *Eur. J. Pharmacol.* 2005; 508: 211-221.

Dettmer, N. The therapeutic effect of glycosaminoglycan polysulfate (Arteparon) in arthroses depending on the mode of administration [German]. *Z. Rheumatol.* 1979; 38:163-181.

Devaki, T., Venmadhi, S., Govindaraju, P. *Med Sci Res.* 1992; 20; 725-730

Dokmeci, D. Ibuprofen and Alzheimer's disease. *Folia Med. Plovdiv.* 2004; 46: 5-10.

Dominique, L. Drug-induced liver diseases. *J. Hepatol.* 2000; 32: 77-88.

Doroshov, J., Hochstein, P. Redox cycling and the mechanism of action of antibiotics in neoplastic diseases. In: *Pathology of Oxygen*, Academic press, New York, 1982; pp. 245-250.

- Eidelman, R.A., Lamas, G.S., Hennerkens, C.H. The new national education program guidelines. *Arch. Intern. Med.* 2002; 162: 2033-2036
- Ellman, G.L.: Tissue sulfhydryl groups. *Arch. Biochem. Biophys.* 1959., 82, 70-77.
- Emery, P. Clinical implications of selective cyclooxygenase-2 inhibition. *Scand. J. Rheumatol.* 1996; 25: 23-28
- Emken, E.A., Adlot, R.O., Duval, S.M., Nelson, G.J.. Influence of dietary arachidonic acid on metabolism *in vivo* of 8 cis, 11 cis, 14-eicosatrienoic acid in humans. *Lipids* 1997; 32: 441-444.
- Eule, H., Beck, H., Evers, H., Fischer, P., Kwiatkowski, H., Merkel, S., Reech, R., Sieler, R., Thomas, E., Weinecke, W. Daily ultrashort chemotherapy and intermittent short-term chemotherapy with 4 drugs of communicable pulmonary tuberculosis treated for the first time. Results of a cooperative multicenter study. *Z. Erkr. Atmungsorgane* 1986;167:29-41
- Evans, D.J. *J. Bacteriol.* 1969; 100: 914-916
- Fabio, P., He'le'ne, T., Michel, T. Enzymatic synthesis of D-glucosaminic acid from D-glucosamine. *Carbohydr. Res.* 2005; 340: 139-141.
- Felix, S., Hans-Martin, B., Karlheinz, S., Dimitrios, V., Gerhard, S., Helmut, K.S., Detlef, S. Hepatitis induced by Kava (*Piper methysticum rhizoma*). *J. Hepatol.* 2003; 39: 62-67
- Fenton, D.M, Eveleigh, D.E. *J. Gen. Microbiol.* 1981; 126: 156-162
- Filipovic-Grcic, J., Skalko-Basnet, N., Jalsenjak, I. Mucoadhesive chitosan-coated liposomes: characteristics and stability. *J. Microencapsul.* 2001; 18: 3-12.
- Fiske, C.H., Subbarow, Y. The colorimetric determination of phosphorus. *J. Biol. Chem.* 1925; 66: 375-400.
- Folch, J., Lees, M., Stanely, G.H.S. A simple method for isolation and purification of total lipids from animal tissues. *J. Biol. Chem.* 1957; 226: 497-509
- Georgieva, N., Gadjeva, V., Tolekova, A. New Isonicotinoylhydrazones With SSA protect against oxidative-hepatic injury of Isoniazid. *TJS*; 2: 37-43, 2004.
- Gibson, J.C., Lee, W.H., Stephan, Z.F. The ansamycins: a novel class of hypolipidemic agents with a high affinity for lipoproteins. *Atherosclerosis* 1995; 112: 47-57.
- Goldman, A.L., Braman, S.S. Isoniazid: a review with emphasis on adverse effects. *Chest* 1972; 62: 71-77.

- Gomez-Lechon, M.J., Ponsoda, X.O.C.E., Donato, T., Castell, J.V., Jover, R.. Diclofenac induces apoptosis in hepatocytes by alteration of mitochondrial function and generation of ROS. *Biochem. Pharmacol.* 2003; 66: 2155-2167.
- Gorss, P. Konrad, E., Mager, H. In *Chitin and Chitosan*, The Japanese Society on Chitin and Chitosan, Tottori Univ. Tottori, Japan, 1982, pp 205-209
- Gupta, S., Husser, R.C., Geske, R.S., Welty, S.E., Smith, C.V. Sex differences in diquat-induced hepatic necrosis and DNA fragmentation in Fisher 344 rats. *Toxicol. Sci.* 2000; 54: 204-211.
- Guzman, R.O, Ramirez, M.D., Campo, M., Fernandez, L.I., Romero, G.M. Hepatotoxicity induced by paroxetine (Spanish), *Med. Clin(Barc)*. 2005; 124: 399-405.
- Habig, W.H., Pabst, M.J. and Jackoby, W.B.C.: Glutathione-S-transferases: The first enzymatic step in enzymatic stepping mercapturic acid formation. *J. Biol. Chem.* 1974; 249: 7130-7139.
- Han, L.K., Kimura, Y., Okuda, H. Reduction in fat storage during chitin-chitosan treatment in mice fed a high-fat diet. *Int. J. Obes. Relat. Metab. Disord.* 1999; 23: 174-179.
- Hiraishi, H., Terano, A., Ota, S., Mutoh, H., Sugimoto, T., Harada, T., Razandi, M., Ivey, K.J. Protection of cultured rat gastric cells against oxidant-induced damage by exogenous glutathione. *Gastroenterol.* 1994; 106: 1199-1207.
- Hiroaki, O., Hajime, N., Yasuyuki, S., Shinichi, A., Norifumi, K., Yoshio, Y., Junji, Y. Overexpression of Thioredoxin Prevents acute Hepatitis caused by Thioacetamide or lipopolysaccharide in Mice. *Hepatology* 2003; 37: 1015-1025.
- Hjerten, S., Pan, H. Purification and characterization of two forms of low affinity calcium ion ATPase from erythrocyte membranes *Biochim. Biophys. Acta.* 1983; 755: 457-466.
- Hokin, L.E., Dhal, J.L., Deukprea, J.D., Dixon, J.F., Hackney, J.F., Perdue, J.E. *J. Biol. Chem.* 1973; 248: 2593-2596
- Hollenz, M., Labenz, J. Gastrointestinal complications under NSAID treatment in the doctor's office (German). *MMW Fortschr Med.* 2004; 146: 42-44.
- Holt, G.A. *Food & Drug Interactions*. Chicago: Precept Press, 1998, 147-149
- Holtzman, M.J. 1991. Arachidonic acid metabolism. *Annu. Rev. Respir. Dis.*, 143,186-203.
- Horn, W.T., Menahan, L.A. A sensitive method for determination of free fatty acids in plasma. *J. Lipid Res.* 1981; 122: 377-381.

- Forward, A., Barley, N. F., Legon, S. and Walter, J. R. Plasma membrane calcium pump iso forms in human and rat liver. *Biochem. J.* 1994; 303: 275-279.
- Hussain, Z., Kar, P., Husain, S.A. Antituberculosis drug-induced hepatitis: risk factors, prevention and management. *Indian J. Exp. Biol.* 2003; 41:1226-1232
- Imogen, M., Julia W., Sarah F., Roger W. Anti-tuberculous therapy and acute liver failure. *The Lancet* 1995; 345: 555-556.
- Ishiyama, H., Ogino, K., Shimomura, Y., Kanke, T., Hobara, T. Hepatotoxicity of diethyldithiocarbamate in rats. *Pharmacol. Toxicol* 1990; 67: 426-430.
- Itaya, K. J. *Lipid Res.* 1977; 23: 377-379
- Jasmer, R.M., Saukkoner, J.J., Blumberg, H.M. Short course rifampicin and pyrazinamide compared with isoniazid for latent tuberculosis infection: a multicenter clinical trial. *Ann. Intern. Med.* 2002, 137:640-647.
- Jeuniaux, C. *Bull. Soc. Chim. Biol.* 1965; 47: 2267-2269.
- Jeuniaux, C. *Chitine et Chitinolyse* 1963, Masson, Paris, pp.215-223
- Jianfeng, X., Baoru, Y., Yaling, D., Bingwen, W., Junxian, W., Heikki, P.K. Effects of sea buckthorn (*Hippophae rhamnoides* L.) seed and pulp oils on experimental models of gastric ulcer in rats. *Fitoterapia* 2002; 73: 644–650.
- Jimenez, M.D., Martin, M.J., Alarcon de la Lastra, C., Bruseghini, L., Esteras, A., Herrerias, J.M., Motilva, V. Role of L-arginine in ibuprofen-induced oxidative stress and neutrophil infiltration in gastric mucosa. *Free Radic. Res.* 2004; 38: 903-911.
- Jing, S.B., Li, L., Ji, D., Takiguchi, Y., Yamaguchi, T. Effect of chitosan on renal function in patients with chronic renal failure. *J. Pharm. Pharmacol.* 1997 ;49: 721-723.
- Kalender, Y., Yel, M., Kalender, S. Doxorubicin hepatotoxicity and hepatic free radical metabolism in rats. The effects of vitamin E and catechin,. *Toxicology* 2005; 209: 39-45.
- Karthikeyan, S. Isoniazid and rifampicin treatment on phospholipids and their subfractions in liver tissue of rabbits. *Drug. Chem. Toxicol.* 2005; 28, 273-280.
- Kaufmann, P, Torok M, Hanni A, Roberts P, Gasser R, Krahenbuhl S. Mechanisms of benzarone and benzbromarone-induced hepatic toxicity. *Hepatology* 2005; 41: 925-935.
- Khor, E., Lim, L.Y. Implantable applications of chitin and chitosan. *Biomaterials* 2003; 24: 2339–2349.

- Kim, S.T., Kim, J.D., Ahn, S.H., Ahn, G.S., Lee, Y.I., Jeong, Y.S. Hepatoprotective and antioxidant effects of *Alnus japonica* extracts on acetaminophen-induced hepatotoxicity in rats. *Phytother. Res.* 2004; 18: 971-975
- Kim, Y.C., Jung, Y.S., Kim, S.K. Effect of betaine supplementation on changes in hepatic metabolism of sulfur-containing amino acids and experimental cholestasis induced by alpha-naphthylisothiocyanate. *Food Chem. Toxicol.* 2005; 43: 663-670.
- King J. In: *Lactate dehydrogenase in practical clinical enzymology*, (Van. D., Ed.) Nostrand Co., London ; 1965: pp 83-93.
- Kissler, H.J., Hauffen, J., Hennig, R., Gepp, H., Schwille, P.O.,. Glucose and lipid metabolism after liver transplantation in inbred rats: consequences of hepatic denervation. *Metabolism* 2005; 54: 881-890.
- Kittur, F.S., Harish Prashanth, K.V., Udaya Sankar, K., Tharanathan, R.N. *Carbohydr. Pol.* 2002; 49: 185-193.
- Knapczyk, J., Krowczyński, L., Krazek, J., Brzeski, M., Nurberg, E., Sehenk, D., Struszezyk, H. In *Chitin and Chitosan*, Elsevier Applied Science, London, 1989, pp 657-664
- Kong, N. A Feasibility Study Of New Routes To Marine Polymers, Chitin And Chitosan, M.S.Thesis, Univ. of Washington, 1975.
- Krisana, S, Khatcharin, S, Tawun, R, Nongnuj, M., Werasak, U., Supot, H.. Solvation structure of glucosamine in aqueous solution as studied by Monte Carlo simulation using ab initio fitted potential. *Chem. Phys. Let.* 2004, 395, 233-238
- Krishnan, G., Sundara Rajulu, G. Z. *Naturf.* 1964; 19 B: 640-652.
- Kruger, F.A., Leighty, E.G., Weissler, A.M.,. Catecholamine stimulation of myocardial lipolysis and fatty acid reesterification. *J. Clin. Invest.* 1967; 46: 1080-1081.
- Kunike, G. *J.Soc.Dyers Col.* 1926; 42: 228-230.
- Kweon, D.K., Song, S.B., Park, Y.Y. Preparation of water-soluble chitosan/heparin complex and its application as wound healing accelerator. *Biomaterials* 2003; 24: 1595-1601.
- Lammler, U.K. *Nature* 1970; 227: 680-685.
- Lanza, F.L., Fakouhi, D, Rubin, A., Davis, R.E., Rack, M.F., Nissen, C., Geis, S. A double-blind placebo-controlled comparison of the efficacy and safety of 50, 100, and 200 micrograms of misoprostol QID in the prevention of ibuprofen-

- induced gastric and duodenal mucosal lesions and symptoms. *Am. J. Gastroenterol.* 1989; 84: 633-636.
- Larrey, D., Vial, T., Pauwels, A., Castot, A., Biour, M., David, M. Hepatitis after Germander (*Teucrium chamaedrys*) administration: another instance of herbal medicine hepatotoxicity. *Ann. Intern. Med.* 1992;117:129–32.
- Leinweber, B., Mahrt, R.,. Animal experimental studies on the problem of liver damage by antitubercular agents (rifampicin and isoniazid) by means of the galactosamine model. *Verh. Dtsch. Ges. Inn. Med.* 1976; 82: 300-303.
- Lenaerts, A.J., Johnson, C.M., Marrieta, K.S., Gruppo, V., Orme, I.M. Significant increases in the levels of liver enzymes in mice treated with anti-tuberculosis drugs. *Int. J. Antimicrob. Agents* 2005; 16: 201-206.
- Litkowski, L.J., Christensen, S.E., Adamson, D.N., Van Dyke, T., Han, S.H., Newman, K.B. Analgesic efficacy and tolerability of oxycodone 5 mg/ibuprofen 400 mg compared with those of oxycodone 5 mg/acetaminophen 325 mg and hydrocodone 7.5 mg/acetaminophen 500 mg in patients with moderate to severe postoperative pain: a randomized, double-blind, placebo-controlled, single-dose, parallel-group study in a dental pain model; *Clin. Ther.* 2005; 27: 418-29
- Loes, M., Wikholm, G., Shields, M., Steinman, D. In *Arthritis: The Doctors' Cure*. Keats Publishing, New Canaan, CT. , 1998; pp 625-636.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L., Randall, R.J. Protein Measurement with the Folin phenol reagent. *J. Biol. Chem.* 1951; 193: 265-275
- Lu, S.C., Kuhlenkamp, J., Robert, A., Kaplowitz, N. Role of glutathione status in protection against ethanol-induced gastric lesions. *Pharmacol.* 1989; 38: 57-60.
- Maddrey, W.C. Drug-induced hepatotoxicity. *J. Clin. Gastroenterol.* 2005; 39: S 83-89
- Madhavan, P. Chitin and chitosan and their novel applications. In: *Popular Science Lecture Series*. Central Institute of Fisheries Technology, Cochin, India, 1992; pp 6-7.
- Majeti, N.V. Kumar, R. A Review of chitin and chitosan applications. *React. Funct. Polym.* 2000; 46: 1-27
- Mandell, G.L., Petri, W.A. Jr. Antimicrobial Agents: Drugs used in the chemotherapy of tuberculosis, *Mycobacterium avium* complex disease and leprosy. In *Goodman and Gilman's The Pharmacological Basis of Therapeutics*, 9th ed. New York: McGraw-Hill, 1996, pp 1158-1160

- Masubuchi, Y., Makayama, S., Horie, T. Role of mitochondrial permeability transition in dichlofenac-induced hepatocyte injury in rats. *Hepatology* 2002; 35: 544-551.
- Mc Alindon, T. E., LaValley, M. P., Gulin, J. P., Felson, D. T.. Glucosamine and chondroitin for treatment of osteoarthritis: a systematic quality assessment and meta-analysis. *J. Am. Med. Assoc.* 2000; 283: 1469-1475.
- Misra, H.P. and Fridovich T. The role of superoxide ion in the auto-oxidation of epinephrine and a simple assay for superoxide dismutase. *J. Biol. Chem.* 1972; 247: 3170-3175.
- Mitra, S.K., Venkataranganna, M.V., Sundaram, R., Gopumadhavan, S. Protective effect of HD-03, a herbal formulation, against various hepatotoxic agents in rats, *Journal of Ethnopharmacology* 1998; 63 : 181–186
- Mohankumar, K., Ramasamy, P. Activities of membrane bound phosphatases, transaminases and mitochondrial enzymes in white spot syndrome virus infected tissues of *Fenneropenaeus indicus*. *Virus Res.* 2006; 118: 130-135.
- Mohur, A., Cook, I.J.Y. Simple methods for measuring serum levels of glutamic-oxalo acetic and glutamic-pyruvic transaminase in routine laboratories. *J. Clin. Pathol.* 1957; 10: 394-399.
- Monaghan, R.L., Eveleigh, D.E., Tewari, R.P., Reese, E.T. *Nature New Biol.*; 1972: 245: 78-80
- Mori, T., Okumura, M., Matsuura, M., Ueno, K., Tokura, S., Okamoto, Y., Minami, S., Fujinaga, T. Effects of chitin and its derivatives on the proliferation and cytokine production of fibroblasts in vitro. *Biomaterials* 1997; 18: 947–51.
- Mutoh, H., Ota, S., Hiraishi, H., Ivey, K.J., Terano, A., Sugimoto, T., Adaptive cytoprotection in cultured rat gastric mucus producing cells. Role of mucus and prostaglandin synthesis. *Dig. Dis. Sci.* 1995; 40: 872-878.
- Muzzarelli, R., Baldassarre, V., Conto, F., Ferrara, P., Biagini, G., Gazzanelli, G., Vasi, V. Biological activity of chitosan: ultrastructural study. *Bioamaterials* 1988; 9: 247-252.
- Muzzarelli, R.A.A. Chitosan-based dietary foods. *Carbohydr. Polym.* 1996; 29: 309-316.
- Muzzarelli, R.A.A. In *Chitin*, Pergamon Press, Oxford, 1977, pp 5-44.
- Muzzarelli, R.A.A. In *Proc. First Int. Conf. Chitin/Chitosan* MIT, Cambridge. 1978. pp 1-3
- Muzzarelli, R.A.A., a. In *Chitin*. Pergamon Press, Oxford. 1977, pp 45-85.

- Muzzarelli, R.A.A., Tanfani, F., Emameli, M, Mariotti, S. a. *N*-(carboxymethylidene) chitosans and *N*-(carboxymethyl) chitosans: Novel chelating polyampholytes obtained from chitosan glyoxylate *Carbohydr. Res* 1982; 107: 199-214.
- Muzzarelli, R.A.A., Tanfani, F., Emameli, M., Mariotti, S. The characterization of *N*-methyl, *N*-ethyl, *N*-propyl, *N*-butyl and *N*-hexyl chitosans, novel film-forming polymers *J.Membr. Sci.* 1982; 16: 295-308.
- Muzzarelli, R.A.A., Tanfani, F., Mariotti, S, Emameli,M. . *N*- (*o*-carboxybenzyl) chitosans: Novel-chelating polyampholytes. *Carbohydr. Polym.* 1983; 2: 145-157.
- Naga Chalasani. Statins and hepatotoxicity: Focus on patients with fatty liver. *Hepatology* 2005; 41: 690-695
- Natelson, S., Scott, M.L., Beffa, C.,. A rapid method for the estimation of urea in biologic fluids. *Am. J. Clin. Pathol.* 195; 121: 275-281.
- Niebes, P. Determination of enzymes and degradation products of glycosamino glycan metabolism in healthy and various subjects. *Clin. Chim. Acta* 1972; 42: 399-408
- Nishimura, K., Nishimura, S., Nishi, N., Saiki, I., Tokura, S., Azuma, I. Immunological activity of chitin and its derivatives. *Vaccine* 1984; 2: 93–99.
- Ohkawa, H., Onishi, N., Yagi, K. Assay for lipid peroxides in animal tissue by thiobabaturic acid reaction. *Anal. Biochem.* 1979; 95: 351-358.
- Ohnishi, T., Suzuki, R., Suzuki, Y., Ozawa, K. *Biochim. Biophys. Acta* 1982; 684: 67-71
- Okamoto, Y., Minami, S., Matsushashi, A., Sashiwa, H., Saimoto, H., Shigemasa, Y., Tanigawa, T., Tanaka, Y., Tokura, S. Polymeric *N*- acetyl- *D*-glucosamine (Chitin) induces histionic activation in dogs. *J. Vet. Med. Sci.* 1993; 55: 739–742.
- Onyema, O.O., Farombi, E.O., Emerole, G.O., Ukoha, A.I., Onyeze, G.O. Effect of vitamin E on monosodium glutamate induced hepatotoxicity and oxidative stress in rats. *Indian J. Biochem. Biophys.* 2006; 43: 20-24.
- Pagila, D.E., Valentine, W.N. Studies on the glutathione characterization of erythrocyte glutathione peroxidase. *J. Lab. Clin. Med.* 1967; 70: 158-169.
- Parekh, A.C., Jung, D.H. Cholestrol determination with ferric acetate- uranyl acetate and sulphuric acid – ferrus sulphate reagents. *Anal. Chem.* 1970; 42: 1423-1427.
- Patwardhan, B., Warude, D., Pushpangadan, P., Bhatt, N. Ayurveda and traditional medicine: A comparative overview. *eCAM* 2005; 2: 465-473.

- Pepicelli, O., Fedele, E., Berardi, M., Raiteri, M., Levi, G., Greco, A., Ajmone-Cat, M.A., Minghetti, L. Cyclo-oxygenase-1 and -2 differently contribute to prostaglandin E2 synthesis and lipid peroxidation after in vivo activation of N-methyl-D-aspartate receptors in rat hippocampus. *J Neurochem.* 2005; 93: 1561-1567.
- Pfeiffer, C. J. Experimental analysis of hydrogen ion diffusion in gastrointestinal mucus glycoprotein. *Am. J. Physiol.* 1981; 240: 176-182.
- Polat, O., Karaman, A.I. Pain control during fixed orthodontic appliance therapy; *Angle Orthod.* 2005; 75: 214-219.
- Porchezian, E., Ansari, S.H. Hepatoprotective activity of *Abutilon indicum* on experimental liver damage in rats. *Phytomedicine* 2005; 12: 62–64.
- Prabakan, M., Anandan, R., Devaki, T. Protective effect of *Hemidesmus indicus* against rifampicin and isoniazid-induced hepatotoxicity in rats. *Fitoterapia* 2000; 71: 55-59
- Prabha, T., Dora Babu, M., Priyambada, S., Agrawal, V.K., Goel, R.K. Evaluation of *Pongamia pinnata* root extract on gastric ulcers and mucosal offensive and defensive factors in rats. *Indian J. Exp. Biol.* 2003; 41: 304-310.
- Raghavendran, H.R., Sathivel, A., Devaki, T. Efficacy of brown seaweed hot water extract against HCl-ethanol induced gastric mucosal injury in rats. *Arch. Pharm. Res.* 2004; 27: 449-453.
- Rainsford, K.D. The ever-emerging anti-inflammatories. Have there been any real advances? *J. Physiol. Paris.* 2002; 95:11-19.
- Rajendra, W. High performance liquid chromatographic determination of amino acids in biological samples by precolumn derivatization with O-phthdialdehyde. *J. Liquid Chromatogr.* 1987; 10:941–955.
- Ray, S.D., Sorge, C.L., Kamendulis, L.M., Corcoran, G.B. Ca⁽⁺⁺⁾- activated DNA fragmentation and dimethylnitroamine induced hepatic necrosis: effects of Ca⁽⁺⁺⁾-endonuclease and poly (ADP-ribose) polymerase inhibitors in mice. *J. Pharm. Exp. Ther.* 1992; 263: 387-394.
- Reginster, J.Y., Rita, D., Lucio, C.R., Richard, L.L., Eric, L., Olivier, B., Giampaolo, G., Yves, H., Jane, E.D., Christiane, G. Long-term effects of glucosamine sulphate on osteoarthritis progression: a randomised, placebo-controlled clinical trial. *The Lancet* 2001; 357: 251-256
- Reichen J, Paumgartner G. Excretory function of the liver. *Int Rev Physiol* 1980; 21: 103-150.

- Rice, E.W. Triglycerides in serum, In Standard methods in clinical chemistry, (Roedrick, P. and Mc Donald, R. P., Eds), Academic press. New York, 1970: pp 215-218
- Rodriguez, A., Buckholz, P. Antimicrobial agents. *J. Microbiol.* 2003; 14: 15-18.
- Rossi, S. Australian Medicines Handbook 2004. Adelaide: Australian Medicines Press. 2004; pp 156-157
- Rostom, A., Goldkind, L., Laine, L. Nonsteroidal anti-inflammatory drugs and hepatic toxicity: a systematic review of randomized controlled trials in arthritis patients. *Am. J. Gastroenterol.* 2005; 100: 1043-1050
- Rudall, K.M. *J. Polym Sci* 1969; 28: 83-88
- Rutherford, F.A., Dunson, W.A. In *Chitin, Chitosan and Related Enzymes*. Academic Press, Inc. Orlando, 1984; pp 135-143.
- Sal'nikova, S.I., Drogovoz, S.M., Zupanets, I.A. The liver-protective properties of D-glucosamine. *Farmakol. Toksikol.* 1990; 53: 33-35.
- Salvatore, S., Antonina, C., Vincenza, M.C., Nunzio, G.M., Agata, P., Matilde Amico-Roxas, Alfredo, B., Giuseppe, C. Effects of adrenomedullin on the contraction of gastric arteries during reserpine-induced gastric ulcer. *Peptides* 2003; 24: 117-122.
- Sang, K.K., Young, C.K. Effects of betaine supplementation on hepatic metabolism of sulfur-containing amino acids in mice. *J. Hepatol.* 2005; 42: 907-913.
- Santhosh, S., Anandan, R. Sini, T.K., Mathew, P.T., Thankappan, T.K. Biochemical studies on the antiulcer effect of glucosamine on antioxidant defense status in experimentally induced peptic ulcer in rats. *J. Clin. Biochem. Nutr.* 2005; 37: 61-66.
- Santhosh, S., Sini, T.K., Anandan, R., Mathew, P.T. Effect of chitosan supplementation on antitubercular drugs-induced hepatotoxicity in rats *Toxicology* 2006; 219: 53-59.
- Sapelli, P.L., Baldassarre, V., Muzzarelli, R.A.A., Emanulli, M. In *Chitin and Nature and Technology*. Plenum Press, New York, 1986; pp 507-512.
- Saraswathy S.D., Shyamala Devi, C.S. Modulating effect of Liv.100, an ayurvedic formulation on antituberculosis drug-induced alterations in rat liver microsomes. *Phytother. Res.* 2001; 15: 501-505
- Saraswathy SD, Shyamala Devi CS. Hepatoprotective effect of Liv.100, a polyherbal formulation, on mitochondrial enzymes in anti-tubercular drug-induced liver damage in rats. *J. Clin. Biochem. Nutr.* 1999; 26: 27-34.

- Saraswathy, S.D., Suja, V., Gurumurthy, P., Shymala Devi, C.S. Effect of Liv.100 against antitubercular drugs (Isoniazid, Rifampicin and Pyrazinamide) induced hepatotoxicity in rats. *Indian J. Pharmacology* 1998; 30: 233-238.
- Sasaki, T., Natsui, S. *Rinsho Kagaku* 1972; 1: 346-348
- Sastry, C.S.P., Ktummuru, M.. Spectrophotometric determination of tryptophan in proteins. *Journal of Food Science and Technology* 1985; 22: 146-147
- Scelzi, S., Giubilei, G., Bartoletti, R., Di, L.F., Mondaini, N., Crisci, A.; Nephrogenic adenoma of bladder after ibuprofen abuse. *Urology* 2004; 64:1030-1033
- Schuppan, D., Jia, J.D., Brinkhaus, B., Hahn, E.G. Herbal products for liver diseases: a therapeutic challenge for the new millenium. *Hepatology*: 1999; 30:1099–104.
- Senthilkumar, R., Nalini, N. Effect of glycine on tissue fatty acid composition in an experimental model of alcohol-induced hepatotoxicity; *Clin. Exp. Pharmacol. Physiol.* 2004; 31: 456-461.
- Setnikar, I., Giacchetti, C., Zanolo, G. Pharmacokinetics of glucosamine in the dog and in man. *Arzneimittelforschung* 1986; 36: 729-735
- Setnikar, I., Giacchetti, C., Zanolo, G. Absorption, distribution and excretion of radioactivity after a single intravenous or oral administration of [¹⁴C] glucosamine to the rat. *Pharmatherapeutica* 1984; 3: 538-550
- Setnikar, I., Palumbo, R., Canali, S., Zanolo, G.. Pharmacokinetics of glucosamine in man. *Arzneimittelforschung* 1993; 43: 1109-1113
- Shakile, A.O., Kramer, D., Mazariegos, G., Fung, J.J., Rakela, J. Acute liver failure: clinical features, outcome analysis and applicability of prognostic criteria. *Liver Transpl.* 2000; 6: 163-169.
- Shannon, M.W., Lovejoy, F.H. Jr. Isoniazid. In: Haddad LM, Winchester JF, eds. *Clinical management of poisoning and drug overdose*. 2d ed. Philadelphia: Saunders, 1990: pp 970-975
- Sheikh, N.M., Philen, R.M., Love, L.A. Chaparral-associated hepatotoxicity. *Arch. Intern. Med.* 1997; 157:913–9.
- Shepherd, R., Reader, S., Falshaw, A. Chitosan functional properties. *Glycoconjugate J.* 1997; 14: 535-542.
- Shevtsova, O.I., 2000. Effect of chitosan on lipid peroxidation in toxic liver injury. *Ukr. Biokhim Zh.* 72, 88-90.

- Shi, J., Aisaki, K., Ikawa, Y., Wake, K.,. Evidence of hepatocyte apoptosis in rat liver after the administration of carbon tetrachloride. *Am. J. Pathol.* 1998; 53: 515-525
- Silvana, D., Petrovic, S.D., Dubravko, B., Marjan N., Alfonso G., Irmgard, M. Evaluation of *Tanacetum larvatum* for an anti-inflammatory activity and or the protection against indomethacin-induced ulcerogenesis in rats *J. Ethnopharmacol.* 2003; 87: 109–113
- Sini, T.K., Santhosh, S., Mathew, P.T. Study of the influence of processing parameters on the production of carboxymethylchitin. *Polymer* 2005; 46: 3128-3131.
- Skakun N.P., Slivka, Y. The correction of hepatotoxicity of antitubercular preparations with tocopherol acetate and riboxin. *Eksp. Klin. Farmakol.* 1992; 55: 52-54.
- Skakun, N.P., Tabachuk, O.P. The comparative action of isoniazid, rifampicin and ethambutol on liver function. *Eksp. Klin. Farmakol.* 1992; 55: 45-47.
- Skou, N. Spatial orientation of Na⁺ K⁺ dependent enzymatic reaction in the cell membrane, In *Biomembranes, Vol 2*, (Manson, L. A., Ed.), Plenum press. New York, 1971; pp.165-186.
- Slot, C. *Scand. J. Clin. Lab. Invest.* 1965; 17: 381-384
- Stenberg, D., Interconvertible enzymes in adipose tissue regulated by cyclic AMP-dependent protein kinase. *Adv. Cyclic Nucleotide Res.* 1976; 7: 157-198.
- Stickel, F., Seitz, H.K. The efficacy and safety of comfrey. *Public Health Nutr.* 2000; 3: 501–8.
- Stryer, L. In: *Biochemistry*, 4th edition, Freeman WH and company, New York, 1995, pp 626-652.
- Sundara Rajulu, G., Jeuniaux, C., Poulicek, V., Voss-Foucart. In *Chitin and Chitosan, The Japanese Society of Chitin and Chitosan*, Tottori Univ., Tottori, Japan, 1982; pp 430-435.
- Synowiecki, J., Al-Khateeb, N.A. Production, properties, and some new applications of chitin and its derivatives. *Crit. Rev. Food Sci. Nutr.* 2003; 43: 145-71.
- Takahara, S., Hamilton, B.H., Nell, J.V., Kobra, T.Y., Ogawa, Y. and Nishimura, E.T.: Hypocatalasemia: A new genetic carried state. *J. Clin. Invest.* 1960; ,29: 610-619.
- Takeuchi, K., Okabe, S., Takagi, K. A new model of stress ulcers in the rat with pylorus ligation and its pathogenesis. *Am. J. Dig. Dis.* 1976; 21: 782-788.

- Tan, P.V., Nyasse, B., Dimo, T., Mezui, C. Gastric cytoprotective anti-ulcer effects of the leaf methanol extract of *Ocimum suave* (Lamiaceae) in rats, *J. Ethnopharmacol.* 2002; 82; 69-74.
- Tanigawa, T., Tanaka, Y., Sashiwa, H., Saimoto, H., Shigemasa, Y. 1992. Various biological effects of chitin derivatives. In: Brine CJ, Sandford PA, Zikakis JP, editors. *Advances in chitin and chitosan*, New York: Elsevier; 1992; pp 206–215.
- Tannenbaum, H., Davis, P., Russell, A.S., Atkinson, M.H., Maksymowych, W., Huang, S.H., Bell, M., Hawker, G.A., Juby, A., Vanner, S., Sibley, J. An evidence-based approach to prescribing NSAIDs in musculoskeletal disease: a Canadian consensus. *Canadian Medical Association Journal* 1996; 155: 77-88
- Tannock, L.R., Peter J. L., Thomas, N., W., Alan, C.. Arterial smooth muscle cell proteoglycans synthesized in the presence of glucosamine demonstrate reduced binding to LDL. *J. Lipid Res.* 2002; 43: 149-157.
- Tasduq, S.A., Peerzada, K., Koul, S., Bhat, R., Johri, R.K. Biochemical manifestations of anti-tuberculosis drugs induced hepatotoxicity and the effect of silymarin. *Hepatol. Res.* 2005; 31: 132-135.
- Timbrell, J. Toxic responses to foreign compounds. In: *Principles of Biochemical Toxicity*, third ed. Taylor and Francis Limited, London, Philadelphia, Hong Kong, 2000, pp. 190–252.
- Timothy, M., Margaret, F., Michael, L., Melissa, L., Karim, K. Effectiveness of Glucosamine for Symptoms of Knee Osteoarthritis: Results from an Internet-Based Randomized Double-Blind Controlled Trial. *Am. J. Med.* 2004; 117: 643-649.
- Tokura, S., Ueno, K., Miyazaki, S., Nishi, N. Molecular weight dependent antimicrobial activity by chitosan. *Macromol. Symp.* 1997; 120: 1–9.
- Tomiyama, N., Takeda, M., Watanabe, M., Kobayashi, H., Harada, T. A further study on the reliability of toxicokinetic parameters for predicting hepatotoxicity in rats receiving a 28-day repeated administration of DDT. *J Toxicol. Sci.* 2004; 29: 505-516.
- Troy C. S., Stephen, P. A., James, M. W. The role of l-thyroxine and hepatic reductase activity in isoniazid-induced hepatotoxicity in rabbits, *Pharmacol. Res.* 1998; 38: 199-207.
- Tyagi, P., Sharma, B.C., Sarin, S.K. Cholestatic liver injury due to ibuprofen. 2005. *Indian J Gastroenterol.* 2005; 24: 77-8

- Van der Lubber, I.M., Verhoef, J.C., Borchard, G., Junginger, H.E. Chitosan and its derivatives in mucosal drug and vaccine delivery. *Eur. J. Pharm. Sci.* 2001; 14: 201–207.
- Van Wisselingh, E. *Jahrb.Wiss. Bot.* 1928; 22: 401-409
- Vane, J.R. Mechanism of action of NSAIDs. *Brit. J. Rheumatol.* 1996; 35: 1-3.
- Vanisree, A.J., Devaki, T. Biochemical studies evaluating the antiulcerogenic potential of *Ocimum sanctum* in experimentally induced peptic ulcer in rats. *J Clin. Biochem. Nutr.* 1995; 19: 79-87.
- Victor, J.N., John R. S. Current concepts drug related hepatotoxicity. *N. Engl. J. Med.* 2006; 354: 731-739.
- Vincenzi, F.F.. A calcium pump in red cell membranes, In Cellular mechanisms for calcium transfer and homeostasis, (Nicholas, G. and Wasserman, R. H., Eds.), Academic press. New York, 1971; pp.135-148.
- Wada, M. The adverse reactions of anti-tuberculosis drugs and its management. *Nippon Rinsho.* 1998; 56: 3091-3095.
- Wada, M., Nishimura, Y., Watanabe, Y., Takita, T., Innami, S. Accelerating effect of chitosan intake on urinary calcium excretion by rats. *Biosci. Biotec. Biochem.* 1997; 61:1206-1208.
- Wagner, W.D. More sensitive assay discriminating galactosamine and glycosamine in mixtures. *Anal. Biochem.* 1979; 94: 394-397
- Warren, L. The thiobarbituric acid assay of sialic acid. *J. Biol. Chem.* 1959; 234: 1971-1975
- Wason, S., Lacouture, P.G., Lovejoy, F.H. Jr. Single high-dose pyridoxine treatment for isoniazid overdose. *JAMA* 1981; 246:1102-1104.
- Weberg, R., Berstad, K., Berstad, A. Acute effects of antacids on gastric juice components in duodenal ulcer patients. *Eur. J. Clin. Invest.* 1990; 20: 511-515.
- Welage, L.S. Overview of pharmacologic agents for acid suppression in critically ill patients. *Am. J. Health. Syst. Pharm.* 2005; 62: S4-S10.
- Werbach, M.R. In *Foundations of Nutritional Medicine*. Tarzana, CA: Third Line Press, 1997; pp 231–232.
- Westphal, J.F., Vetter, D., Brogard, J.M.,. Hepatic side effects of antibiotics. *J. Antimicrob. Chemother.* 1994; 33: 387-401.

- Wu, T., Zivanovic, S., Draughon, F.A., Sams, C.E. Chitin and chitosan--value-added products from mushroom waste. *J. Agric. Food. Chem.* 2004; 52: 7905-7910.
- Xing, R., Liu, S., Guo, Z., Yu, H., Wang, P., Li, C., Li, Z., Li, P. Relevance of molecular weight of chitosan and its derivatives and their antioxidant activities in vitro. *Bioorg. Med. Chem.* 2005; 13: 1573-1577.
- Yan, Y., Wanshun, L., Baoqin, H., Bing, L., Chenwei, F. Protective effects of chitosan oligosaccharide and its derivatives against carbon tetrachloride-induced liver damage in mice. *Hepatol. Res.* 2006; 35: 178-184.
- Yao, H.T., Chiang, M.T.,. Plasma lipoprotein cholesterol in rats fed a diet enriched in chitosan and cholesterol. *J. Nutr. Sci. Vitaminol.* 2002; 48: 379-383.
- Yasser, S.E., Ahmed, A.B. *Analytica Chimica Acta* 2002; 462: 125-131.
- Yeel, Y.N., So, H.L., Liu, E.S.L., Shin, V.Y., Cho, C.H. Effect of polysaccharides from *Angelica sinensis* on gastric ulcer healing. *Life Sciences* 2003; 72: 925-932.
- Yoshida, E.M., McLean, C.A., Cheng, E.S., Blanc, P.D., Somberg, K.A., Ferrell, L.D. Chinese herbal medicine, fulminant hepatitis, and liver transplantation. *Am. J. Gastroenterol.* 1996; 91: 2647-8.
- Yoshikawa, T., Yoshida, N., Naito, Y., Takemura, T., Miyagawa, H., Tanigawa, T., Kondo, M. Role of oxygen radicals in the pathogenesis of gastric mucosal lesions induced by water-immersion restraint stress and burn in rats. *J. Clin. Biochem. Nutr.* 1990; 8: 227-234.
- Yoshikawa, T., Miyagawa, H., Yoshida, N., Sugino, S., Kondo, M. Increase in lipid peroxidation in rat gastric mucosal lesions induced by water-immersion restraint stress. *J. Clin. Biochem. Nutr.* 1986; 1: 271-277.
- Yoshikawa, T., Yoshida, N., Miyagawa, H., Takemura, T., Tanigawa, T., Sugino, S., Kondo, M. Role of lipid peroxidation in gastric mucosal lesions induced by burn shock in rats. *J. Clin. Biochem. Nutr.* 1987; 2: 163-170.
- Yu Shao, Rama, A., Mike, M., Uwe, K., Stanley, L. A stability-indicating HPLC method for the determination of glucosamine in pharmaceutical formulations. *Journal of Pharmaceutical and Biomedical Analysis* 2004; 35: 625-631
- Zarzycki W, Badurski J, Jaworowska E, Sarosiek J. Secretion of gastric mucus following maximal pentagastrin stimulation. *Dtsch. Z. Verdau. Stoffwechselkr.* 1986; 46:209-216.
- Zhang, Z., Gildersleeve, J., Yang, Y.Y., Xu, R., Loo, J.A., Uryu, S., Wong, C.H., Schultz, P.G. A new strategy for the synthesis of glycoproteins. *Science* 2004; 303: 371-373.

Zupanets, I.A., Drogovoz,, S.M., Bezdetko, N.V., Rechkiman, I.E., Semenov, A.N.:
The influence of glucosamine on the antiexudative effect of nonsteroidal
antiinflammatory agents [Russian]. *Farmakol. Toksikol.* 1991; 54: 61-63

LIST OF PUBLICATIONS

- Effect of chitosan supplementation on antitubercular drugs-induced hepatotoxicity in rats, *Toxicology* 2006; 219: 53-59.
- Biochemical studies on the antiulcer effect of glucosamine on antioxidant defense status in experimentally induced peptic ulcer in rats. *Journal of Clinical Biochemistry and Nutrition* 2005; 37: 61-66.
- Study of the influence of processing parameters on the production of carboxymethylchitin. *Polymer* 2005; 46: 3128-3131
- Protective effect of glucosamine against ibuprofen-induced peptic ulcer in rats. *Journal of Gastroenterology and Hepatology* (In Press)
- Effect of different organic acids in silage preparation from shrimp shell waste. *Fishery Technology* (In Press)
- Preparation and properties of glucosamine and carboxymethylchitin from prawn shell. *Journal of Applied Polymer Science* (Accepted for publication)
- Hepatoprotective activity of chitosan against isoniazid and rifampicin-induced toxicity in experimental rats. *European Journal of Pharmacology* (Accepted for publication)



Effect of chitosan supplementation on antitubercular drugs-induced hepatotoxicity in rats

S. Santhosh^{a,*}, T.K. Sini^a, R. Anandan^b, P.T. Mathew^a

^a Fish Processing Division, Central Institute of Fisheries Technology, Cochin 682 029, India

^b Biochemistry and Nutrition Division, Central Institute of Fisheries Technology, Cochin 682 029, India

Received 9 September 2005; received in revised form 21 October 2005; accepted 1 November 2005

Available online 5 December 2005

Abstract

We have studied the protective effect of chitosan on isoniazid- and rifampicin-induced hepatotoxicity with respect to the changes in the levels of diagnostic marker enzymes (in serum), lipid components and lipid peroxidation (in serum and liver). The oral administration of antitubercular drugs caused a significant elevation in the levels of diagnostic marker enzymes and cholesterol, triglycerides, free fatty acids and lipid peroxidation in serum and liver of experimental rats. There was a slight decline in the level of phospholipids in liver tissue also observed. Co-administration of chitosan significantly prevented the antitubercular drugs-induced elevation in the levels of serum diagnostic marker enzymes (alanine amino transferase, aspartate amino transferase, lactate dehydrogenase, acid phosphatase and alkaline phosphatase) in experimental groups of rats. It exerted a significant antilipidemic effect against isoniazid- and rifampicin-induced hepatitis by maintaining the levels cholesterol, triglycerides, free fatty acids and phospholipids in serum and liver at near normalcy. A tendency to prevent the isoniazid- and rifampicin-induced lipid peroxidation was also observed. The results of the present study indicated that the hepatoprotective effect of chitosan might be ascribable to its antilipidemic effect and/or antioxidant property.

© 2005 Elsevier Ireland Ltd. All rights reserved.

Keywords: Chitosan; Isoniazid; Rifampicin; Hepatotoxicity; Diagnostic marker enzymes; Lipids; Lipid peroxidation

1. Introduction

Tuberculosis (TB) is a hazardous disease, which gradually swallows the life span human beings. The global prevalence of tuberculosis was 32% (1.86 billion people) and the global case fatality rate was 23% (Mindie and Gabriel, 2002). The number of estimated cases of tuberculosis was 7.96 million in 1997, with 80% of all

incident tuberculosis cases being found in 22 countries and more than 40% in five South-East Asian countries. Isoniazid was the first effective bactericidal drug used to treat tuberculosis and till far, an important part of most antitubercular drug regimens. Rifampicin, which is another effective bactericidal drug, was added to the regimen in 1962 and has remained the most effective antitubercular combination along with isoniazid (Snider et al., 1984). Although both these drugs are very potent against the tuberculosis bacillus, both are well-known hepatotoxic drugs (Steele et al., 1991).

Hepatitis has been reported to occur in 0.46% of patients receiving these antitubercular drugs (Alexander et al., 1982) and the rate of hepatotoxic reaction was

* Corresponding author at: "Souparnika", Nemmara College (P.O.), Nellikattuparambu, Nemmara, Palakkad 678 508, Kerala, India.
Tel.: +91 4923 243558.

E-mail address: santhoshsethu@rediffmail.com (S. Santhosh).

reported much higher in Indian patients (Ramachandran, 1980) as compared to that reported from developed countries (Mindie and Gabriel, 2002; O'Brien et al., 1983) even when similar dose schedule was used. Nelson et al. (1976) have postulated that one of the isoniazid metabolites acts as an acetylating agent, causing injury to the macromolecules of hepatocytes. Rifampicin, a powerful inducer of drug metabolizing enzymes in man and rats contributes to the hepatotoxicity of isoniazid by enhancing the rate of production of toxic metabolites. However, the exact mechanisms responsible for injury to hepatocytes caused by isoniazid or rifampicin or the combination are still not clearly understood.

Marine organisms produce many bioactive substances, which are having a lot of potential applications. Chitosan, an important polysaccharide of marine origin, is prepared from the shells of crustaceans (Sini et al., 2005). It has profound applications in the fields of clarification and purification, chromatography, paper and textiles, photography, food and nutrition, agriculture, pharmaceutical and medical, cosmetics, biodegradable membranes and biotechnology (Nishimura et al., 1984). Chitosan has been reported to possess immunological (Mori et al., 1997), antibacterial (Tokura et al., 1997) and wound healing (Okamoto et al., 1993) properties. Chitosan is also an effective and adequate haemostatic agent even under the most severe conditions of anticoagulation. Chitosan solution is found beneficial for healing "athletes foot" conditions (Allan et al., 1984). Both hard and soft contact lenses can be made from chitosan. Sapelli et al. (1986) studied the application of chitosan in dentistry.

Han et al. (1999) reported that supplementation of chitosan reduces high fat diet-induced lipidemia by its antilipidemic property. The free radical quenching property of this marine polysaccharide has also been studied in detail (Xing et al., 2005). Reports by Filipovic-Grcic et al. (2001) indicated the membrane stabilizing property of chitosan. In our laboratory, Anandan et al. (2004) observed the antiulcerogenic potential of chitosan against HCl-ethanol-induced peptic ulcer in rats. Since chitosan can wrap solid particles in liquids to bring them together and agglomerate, it is effectively utilized for the controlled release of drug (Krisana et al., 2004). Though the beneficial effects of chitosan have been extensively studied, the antihepatotoxic potential of chitosan on lipid metabolism in antitubercular drugs-induced hepatotoxicity has not yet been explored.

In the present study, an attempt has been made to assess the antihepatotoxic effects of chitosan on isoniazid- and rifampicin-induced hepatotoxicity was studied with respect to changes in the levels of diagnostic

marker enzymes, lipid components and lipid peroxidation.

2. Materials and methods

2.1. Chemicals and drugs

Isoniazid and rifampicin were obtained by M/s. Lupin Pharmaceuticals Ltd., Mumbai. Bovine serum albumin, tetraethoxy propane and cholesterol were from M/s Sigma Chemical Company, St. Louis, MO, USA. Chitosan (M_w 750,000 Da; viscosity 8 cps; deacetylation rate 85–87%; purity 98.6%) used in this study was prepared from chitin in our laboratory (Madhavan, 1992).

2.2. Animals

Wistar strain male albino rats, weighing 120–150 g, were used for the experiment. The animals were housed in polypropylene cages under hygienic conditions and maintained at normal room temperature. The animals were allowed to have food and water ad libitum. The experiment was conducted according to the guidelines of the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA), New Delhi, India and approved by the Institutional Animal Ethics Committee (IAEC).

2.3. Experimental design

The experimental animals were divided into four groups of six rats each. Rats in Group I (normal control) were given only the standard diet. In Group II (drug control), rats were treated with chitosan (100 mg/kg body weight/day, intragastric intubation for 30 days). In Group III, hepatotoxicity was induced by oral administration of isoniazid and rifampicin (200 mg each/kg body weight/day) for a period of 30 days. Group IV animals were co-administered with chitosan (100 mg/kg body weight/day, intragastric intubation for 30 days) along with antitubercular drugs described for Group III.

At the end of the experiment, i.e. 24 h after the last dose of antitubercular drugs, rats were killed and blood was collected without any anticoagulant. The serum separated was used for the determination of alanine aminotransferase [ALT] (Mohur and Cook, 1957), aspartate amino transferase [AST] (Mohur and Cook, 1957), lactate dehydrogenase [LDH] (King, 1965a), alkaline phosphatase [ALP] (King, 1965b) and acid phosphatase [ACP] (King, 1965b). HDL-cholesterol, and LDL-cholesterol in serum were separated according to the method of Burstein and Scholnick (1972). The liver tissue was excised immediately and washed with ice-cold saline. A portion of liver tissue was homogenized with 0.1 M Tris-HCl, pH 7.4 and the homogenate was used for the determination of lipid peroxidation (Ohkawa et al., 1979).

Lipids were extracted from liver tissue according to the method of Folch et al. (1957) using a chloroform-methanol

Cholesterol content was determined by the method of Parekh and Jung (1970), triglycerides by the method of (1970), free fatty acids by the method of Horn and Mahan (1981). The phospholipids content were estimated by method of Fiske and Subbarow (1925) as inorganic phosphorus liberated after perchloric acid digestion.

Statistical analysis

Results were expressed as mean \pm S.D. One-way analysis of variance (ANOVA) was carried out, and the statistical comparisons among the groups were performed with Tukey's test using a statistical package program (SPSS 10.0 for Windows).

Results

Table 1 shows the levels of serum diagnostic marker enzymes (ALT, AST, LDH, ALP and ACP) in normal and experimental groups of rats. There was a significant

elevation noticed in the levels of serum diagnostic marker enzymes in Group III antitubercular drugs-administered rats as compared to that of normal rats. Co-administration of chitosan significantly ($P < 0.001$) prevented the antitubercular drugs-induced rise in the levels of these diagnostic marker enzymes in serum of Group IV animals as compared to those Group III rats, indicating the cytoprotective activity of chitosan. There were no significant changes observed in the levels of diagnostic marker enzymes in Group II chitosan alone administered rats.

Tables 2 and 4 show the levels of lipid components and lipid peroxidation in serum and liver of normal and experimental groups of rats, respectively. Significant ($P < 0.001$) increases in the levels of cholesterol, triglycerides and free fatty acids were observed in the serum and liver of Group III isoniazid- and rifampicin-administered rats as compared to that of Group I control

Table 1
Levels of alanine aminotransferase (ALT), aspartate aminotransferase (AST), lactate dehydrogenase (LDH), acid phosphatase (ACP) and alkaline phosphatase (ALP) levels in the serum of normal and experimental groups of rats

Parameters	Group I control	Group II chitosan (A)	Group III hepatotoxicity (B)	Group IV (A + B)
ALT	121 \pm 7.23	112 \pm 6.31	314 \pm 29.61 ^{a,b}	133 \pm 9.54 ^c
AST	102 \pm 9.02	94.5 \pm 5.69	297 \pm 26.74 ^{a,b}	98.4 \pm 5.23 ^c
LDH	178 \pm 14.92	182 \pm 16.0	284 \pm 25.63 ^{a,b}	193 \pm 17.56 ^c
ACP	14.3 \pm 0.98	12.7 \pm 1.02	36.8 \pm 2.41 ^{a,b}	16.5 \pm 1.18 ^{c,d,e}
ALP	98.4 \pm 7.46	94.1 \pm 7.21	224 \pm 18.0 ^{a,b}	118 \pm 9.77 ^{c,d,e}

ALT, AST and LDH = μ mol pyruvate liberated/h/l; ACP and ALP = μ mol phenol liberated/h/l; A: chitosan, 100 mg/kg body weight/day (intragastric intubation) for 30 days; B: 200 mg each/kg body weight/day (intragastric intubation) for 30 days; A + B: chitosan co-administered (as in A) with antitubercular drugs (as in B) for 30 days. Values are expressed as the mean \pm S.D. for six animals.

^a $P < 0.001$ significantly different compared with Group I control animals.

^b $P < 0.001$ significantly different compared with Group II control animals.

^c $P < 0.001$ significantly different compared with Group III hepatotoxicity induced rats.

^d $P < 0.01$ significantly different compared with Group II animals.

^e $P < 0.05$ significantly different compared with Group I control rats.

Table 2
Levels of cholesterol, triglycerides, free fatty acids (FFA), phospholipids and lipid peroxides levels in the serum of normal and experimental groups of rats

Parameters	Group I control	Group II chitosan (A)	Group III hepatotoxicity (B)	Group IV (A + B)
Cholesterol	54.5 \pm 0.48	48.1 \pm 0.24 ^a	78.9 \pm 0.62 ^{a,b}	56.1 \pm 0.27 ^{a,b,c}
Triglycerides	47.4 \pm 0.36	40.6 \pm 0.26 ^a	95.8 \pm 0.76 ^{a,b}	54.9 \pm 0.34 ^{a,b,c}
FFA	10.2 \pm 0.09	8.34 \pm 0.51 ^a	18.1 \pm 0.12 ^{a,b}	10.5 \pm 0.98 ^{b,c}
Phospholipids	108 \pm 9.8	94.2 \pm 8.72	128 \pm 10.4 ^{b,d}	102 \pm 9.16 ^c
Lipid peroxides	1.43 \pm 0.11	1.36 \pm 0.09	3.42 \pm 0.22 ^{a,b}	1.57 \pm 0.10 ^{c,e}

Cholesterol, triglycerides, FFA and phospholipids = mg/dl; lipid peroxides = nmol MDA/ml; A: chitosan, 100 mg/kg body weight/day (intragastric intubation) for 30 days; B: 200 mg each/kg body weight/day (intragastric intubation) for 30 days; A + B: chitosan co-administered (as in A) with antitubercular drugs (as in B) for 30 days. Values are expressed as the mean \pm S.D. for six animals.

^a $P < 0.001$ significantly different compared with Group I control animals.

^b $P < 0.001$ significantly different compared with Group II animals.

^c $P < 0.001$ significantly different compared with Group III hepatotoxicity induced animals.

^d $P < 0.01$ significantly different compared with Group I rats.

^e $P < 0.05$ significantly different compared with Group II rats.

Table 3

Levels of LDL-cholesterol and HDL-cholesterol in the liver of normal and experimental groups of rats

Parameters	Group I control	Group II chitosan (A)	Group III hepatotoxicity (B)	Group IV (A + B)
LDL-cholesterol	41.7 ± 1.93	37.4 ± 1.58 ^d	50.1 ± 2.68 ^{a,b}	40.8 ± 2.36 ^{e,f}
HDL-cholesterol	25.9 ± 0.96	27.3 ± 1.02	22.5 ± 1.25 ^{a,b}	25.3 ± 1.01 ^e

LDL-cholesterol and HDL-cholesterol = mg/dl; A: chitosan, 100 mg/kg body weight (intra-gastric intubation) for 30 days; B: 200 mg each/kg body weight/day (intra-gastric intubation) for 30 days; A + B: chitosan co-administered (as in A) with antitubercular drugs (as in B) for 30 days. Values are expressed as the mean ± S.D. for six animals.

^a $P < 0.001$ significantly different compared with Group I control animals.

^b $P < 0.001$ significantly different compared with Group II animals.

^c $P < 0.001$ significantly different compared with Group III hepatotoxicity induced rats.

^d $P < 0.01$ significantly different compared with Group I control animals.

^e $P < 0.01$ significantly different compared with Group II rats.

^f $P < 0.05$ significantly different compared with Group II rats.

rats. A slight decline in the level of HDL-cholesterol with concomitant rise in LDL-cholesterol level was also noted (Table 3). The rats co-administered with chitosan showed significantly near normal levels of lipid components in serum and liver of Group IV rats as compared to that of Group III hepatotoxicity induced rats, indicating the antilipidemic effect of chitosan. There was a significant rise ($P < 0.01$) in the level of phospholipids was noted in serum Group III antitubercular drugs administered rats. This was paralleled by a slight reduction in the level of phospholipids in liver tissue of Group III rats. Chitosan co-administration resulted in the prevention of antitubercular drugs-induced phospholipids depletion in the liver of experimental groups of rats. It also exerted a significant antilipidperoxidative effect by blocking the induction of isoniazid- and rifampicin-mediated lipid peroxidation in serum and liver of Group

IV rats (Tables 2 and 4). The rats administered with chitosan alone did not show any adverse effects, indicating that chitosan is non-toxic.

4. Discussion

Antibiotic-related liver injuries cover most of the clinical and pathological expressions of hepatic dysfunction, including cytotoxic hepatitis (Westphal et al., 1994). There was a significant ($P < 0.001$) elevation observed in the levels of AST, ALT, LDH, ALP and ACP in the serum of Group III antitubercular drugs-administered rats as compared to Group I normal controls (Table 1), which is in line with an earlier reported study (Imogen et al., 1995). Elevated levels of these enzymes in serum are presumptive markers of drug-induced necrotic lesions in the hepatocytes (Amr and Alaa, 2005). Enhanced sus-

Table 4

Levels of cholesterol, triglycerides, free fatty acids (FFA), phospholipids and lipid peroxides levels in the liver of normal and experimental groups of rats

Parameters	Group I control	Group II chitosan (A)	Group III hepatotoxicity (B)	Group IV (A + B)
Cholesterol	9.18 ± 0.81	8.05 ± 0.54	15.4 ± 0.98 ^{a,b}	10.4 ± 0.87 ^{b,c,f}
Triglycerides	18.5 ± 1.25	15.3 ± 0.95 ^d	29.9 ± 1.75 ^{a,b}	18.2 ± 1.29 ^{c,e}
FFA	0.94 ± 0.05	0.83 ± 0.04 ^f	1.73 ± 0.11 ^{a,b}	0.88 ± 0.04 ^c
Phospholipids	27.3 ± 1.80	28.5 ± 2.15	24.2 ± 1.84 ^{c,f}	27.6 ± 2.03 ^e
Lipid peroxides	0.96 ± 0.07	0.87 ± 0.04	1.91 ± 0.165 ^{a,b}	0.98 ± 0.05 ^c

Cholesterol, triglycerides, FFA and phospholipids = mg/g tissue; lipid peroxides = nmol MDA/mg protein; A: chitosan, 100 mg/kg body weight (intra-gastric intubation) for 30 days; B: 200 mg each/kg body weight/day (intra-gastric intubation) for 30 days; A + B: chitosan co-administered (as in A) with antitubercular drugs (as in B) for 30 days. Values are expressed as the mean ± S.D. for six animals.

^a $P < 0.001$ significantly different compared with Group I control animals.

^b $P < 0.001$ significantly different compared with Group II animals.

^c $P < 0.001$ significantly different compared with Group III hepatotoxicity induced rats.

^d $P < 0.01$ significantly different compared with Group I control animals.

^e $P < 0.01$ significantly different compared with Group II rats.

^f $P < 0.05$ significantly different compared with Group I control rats.

^g $P < 0.05$ significantly different compared with Group III hepatotoxicity induced rats.

permeability of hepatocytes cell membrane to the isoniazid- and rifampicin-induced peroxidative damage might have resulted in increased release of these diagnostic marker enzymes into the systemic circulation. In the present study, co-administration of chitosan maintained the levels of these diagnostic marker enzymes in the serum of Group IV animals at near normalcy as compared to Group III rats, indicating the cytoprotective effect of chitosan (Anandan et al., 2004). It probably did so by preventing the antitubercular drugs-induced necrotic damage by its membrane stabilizing action (Filipovic-Grcic et al., 2001). Reports by Anandan et al. (2004) showed that chitosan administration prevented experimental ulcer in rats by its cytoprotective action.

The major disorder encountered in antitubercular drugs-induced hepatitis is fatty accumulation in the liver, which develops either due to excessive supply of lipids to the liver or interference with lipid deposition. The pathogenesis is multifactorial, reflecting complex biosynthetic, enzymatic and catabolic derangement in lipoprotein metabolism. In the present study, the levels of total cholesterol and LDL-cholesterol were significantly ($P < 0.001$) higher in Group III isoniazid and rifampicin-administered rats as compared to that of normal control animals (Tables 2–4), indicating the antitubercular drugs-induced hypercholesterolemic condition. Also the level of HDL-cholesterol was slightly reduced in Group III rats (Table 3). This is in line with an earlier study (Gibson et al., 1995). Increased cholesterol levels in the liver might be due to increased uptake of LDL from the blood by the tissues (Kissler et al., 2005). The abnormal cholesterol deposition is favored by the dangerous tendency of cholesterol to passive exchange between the plasma lipoproteins and the cell membranes (Brown and Goldstein, 1986).

In the present study, co-administration with chitosan significantly ($P < 0.001$) reduced the antitubercular drugs-induced elevation in the levels of total cholesterol and LDL-cholesterol in Group IV rats as compared to that of Group III rats. Also the level of HDL-cholesterol was maintained at near normalcy. Earlier Yao and Chiang (2002) have shown that chitosan intake was effective in lowering the plasma total cholesterol, VLDL-cholesterol and LDL-cholesterol levels in experimental animals. In the present study, a slight reduction in the level of total cholesterol and LDL-cholesterol were also observed in Group II chitosan-fed normal animals, establishing the hypocholesterolemic property of chitosan. The hepatoprotective effect of chitosan is probably related to its ability to inhibit the lipid accumulation in the liver tissue by its antilipidemic property (Xing et al., 2005).

The levels of triglycerides were significantly ($P < 0.001$) higher in serum and liver tissue of Group III rats as compared to Group I control animals (Tables 2 and 4). Increased lipolysis of depot triglycerides liberates free fatty acids from adipose tissue stores (Kruger et al., 1967; Stenberg, 1976) and the free fatty acids liberated by the adipose tissue are also taken up by the liver tissue, leading to the hypertriglyceridemic condition. In the present study, a significant ($P < 0.001$) reduction was noticed in the levels of triglycerides and free fatty acids in serum and liver of Group IV chitosan-administered rats as compared to that of Group III isoniazid- and rifampicin-induced hepatotoxic rats, indicating the hypolipidemic nature of chitosan. The hypolipidemic property of chitosan has already been reported in high fat diet fed experimental animals (Xing et al., 2005).

The levels of phospholipids were significantly ($P < 0.01$) decreased in the liver tissue of Group III isoniazid- and rifampicin-administered rats as compared to that of normal control animals (Table 4). This is in accordance with an earlier study (Karthikeyan, 2005), which indicates that hepatic injury related alterations in lipid composition of liver tissue appears to occur due to the destruction of hepatocytes. The intracellular calcium concentration has been reported to rise in antitubercular drugs-induced hepatitis (Tasduq et al., 2005). This intracellular Ca^{2+} is an inducer of phospholipase A_2 , which degrades membrane phospholipids. The increased peroxidation of membrane phospholipids releases free fatty acid by the action of phospholipase A_2 . Hence, the significant ($P < 0.001$) elevation noticed in free fatty acid content in isoniazid- and rifampicin-administered rats might be due to enhanced breakdown of membrane phospholipids. Recent study (Karthikeyan, 2005) indicates that accelerated degradation of membrane phospholipids is very likely the biochemical basis for the antitubercular drugs-induced hepatocellular injury.

In the present study, the co-administration of chitosan significantly ($P < 0.05$) reduced the antitubercular drugs-induced degradation of membrane phospholipids in liver, establishing its membrane stabilizing effect. It probably did so by preventing changes in fatty acid composition and inhibiting the peroxidative deterioration of hepatocellular membranes. It is possible that stabilization of hepatocellular membranes might prolong the viability of hepatocytes.

Lipid peroxidation of membranes is regulated by the availability of substrate in the form of polyunsaturated fatty acids (PUFA), the availability of inducers such as free radicals and the excited state molecules to

initiate propagation, the antioxidant defense status of environment and the physical status of membrane lipids (Pepicelli et al., 2005). Antibiotic therapy can favor free radical production in excess of basal rates. Many antibiotics that depend on bound metals for their activity are able to generate free radicals and cause cellular damage (Doroshov and Hochstein, 1982). In the present study, a significant ($P < 0.001$) increase in the level of lipid peroxides in the liver tissue of antitubercular drugs-treated rats (Table 4) was observed. This is in corroboration with an earlier investigation (Saraswathy and Shyamala Devi, 2001), which suggested that the high vulnerability of liver to peroxidative damage is mainly due to a decline in the level of free radicals for scavengers. Free radicals initiate lipid peroxidation of biological membranes. The combination of isoniazid and rifampicin treatment in experimental animals enhanced lipid peroxidation, indicating increased oxidative stress in liver (Skakun and Slivka, 1992).

Antioxidants are necessary for preventing the formation of free radicals and they inhibit some of the deleterious actions of reactive oxygen species that damage lipids, DNA and proteins (Eidelman et al., 2002). Co-treatment with chitosan resulted in significant ($P < 0.001$) reduction in the level of lipid peroxidation towards near normalcy, as compared with the levels in the Group II antitubercular drugs-administered rats. It probably did so by its antioxidant nature (Mohur and Cook, 1957) by blocking isoniazid- and rifampicin-induced lipid peroxidation.

In conclusion, the results of the present study indicate that the co-treatment with chitosan may prevent antitubercular drugs-induced hepatotoxicity in rats. The overall hepatoprotective effect of chitosan is probably due to a counteraction of free radicals by its antioxidant nature and/or to its ability to inhibit lipid accumulation by its antilipidemic property.

References

- Alexander, M.R., Louie, S.G., Guemsey, B.G., 1982. Isoniazid associated hepatitis. *Clin. Pharmacol.* 1, 148–153.
- Allan, G.G., Altman, L.C., Bensinger, R.E., Ghosh, D.K., Hirabayashi, Y., Neogi, A.N., Neogi, S., 1984. Chitin, Chitosan and Related Enzymes. Academic Press Inc., Orland, pp. 119–133.
- Amr, A., Alaa, A.H., 2005. Oxidative stress mediates drug-induced hepatotoxicity in rats: a possible role of DNA fragmentation. *Toxicology* 208, 367–375.
- Anandan, R., Suseela Mathew, Viswanathan Nair, P.G., 2004. Anticarcinogenic effects of chitin and chitosan on mucosal antioxidant defense system in HCl-ethanol induced ulcer in rats. *J. Pharm. Pharmacol.* 56, 265–269.
- Brown, M.S., Goldstein, J.L., 1986. A receptor mediated pathway for cholesterol homeostasis. *Science* 232, 34–47.
- Burstein, M., Scholnick, H.R., 1972. Precipitation of chylomicron and very low density lipoprotein from human serum with sodium lauryl sulphate. *Life Sci.* 11, 177–184.
- Doroshov, J., Hochstein, P., 1982. Redox cycling and the mechanism of action of antibiotics in neoplastic diseases. In: *Pathology of Oxygen*. Academic Press, New York, pp. 245–250.
- Eidelman, R.A., Lamas, G.S., Hennerkens, C.H., 2002. The new national education program guidelines. *Arch. Intern. Med.* 162, 2033–2036.
- Filipovic-Grcic, J., Skalko-Basnet, N., Jalsenjak, I., 2001. Mucoadhesive chitosan-coated liposomes: characteristics and stability. *J. Microencapsul.* 18, 3–12.
- Fiske, C.H., Subbarow, Y., 1925. The colorimetric determination of phosphorus. *J. Biol. Chem.* 66, 375–400.
- Folch, J., Lees, M., Stanely, G.H.S., 1957. A simple method for isolation and purification of total lipids from animal tissues. *J. Biol. Chem.* 226, 497–509.
- Gibson, J.C., Lee, W.H., Stephan, Z.F., 1995. The ansamycins: a novel class of hypolipidemic agents with a high affinity for lipoproteins. *Atherosclerosis* 112, 47–57.
- Han, L.K., Kimura, Y., Okuda, H., 1999. Reduction in fat storage during chitin-chitosan treatment in mice fed a high-fat diet. *Int. J. Obes. Relat. Metab. Disord.* 23, 174–179.
- Horn, W.T., Menahan, L.A., 1981. A sensitive method for determination of free fatty acids in plasma. *J. Lipid Res.* 122, 377–381.
- Imogen, M., Julia, W., Sarah, F., Roger, W., 1995. Anti-tuberculous therapy and acute liver failure. *The Lancet* 345, 555–556.
- Karthikeyan, S., 2005. Isoniazid and rifampicin treatment on phospholipids and their subfractions in liver tissue of rabbits. *Drug Chem. Toxicol.* 28, 273–280.
- King, J., 1965a. In: Van, D. (Ed.), *Lactate Dehydrogenase in Practical Clinical Enzymology*. Nostrand Co., London, pp. 83–93.
- King, J., 1965b. In: Van, D. (Ed.), *Lactate Dehydrogenase in Practical Clinical Enzymology*. Nostrand Co., London, pp. 191–208.
- Kissler, H.J., Hauffen, J., Hennig, R., Gepp, H., Schwillie, P.O., 2005. Glucose and lipid metabolism after liver transplantation in inbred rats: consequences of hepatic denervation. *Metabolism* 54, 881–890.
- Krisana, S., Khatcharin, S., Tawun, R., Nongnuj, M., Werasak, U., Supot, H., 2004. Solvation structure of glucosamine in aqueous solution as studied by Monte Carlo simulation using ab initio fitted potential. *Chem. Phys. Lett.* 395, 233–238.
- Kruger, F.A., Leighty, E.G., Weissler, A.M., 1967. Catecholamine stimulation of myocardial lipolysis and fatty acid reesterification. *J. Clin. Invest.* 46, 1080–1081.
- Madhavan, P., 1992. Chitin and chitosan and their novel applications. In: *Popular Science Lecture Series—April*. Central Institute of Fisheries Technology, Cochin, India, pp. 10–11.
- Mindie, H.N., Gabriel, G., 2002. Does isoniazid cause more serious hepatotoxicity in Hepatitis B virus carriers? *Am. J. Gastroenterol.* 97, 1092–1093.
- Mohur, A., Cook, I.J.Y., 1957. Simple methods for measuring serum levels of glutamic-oxalo acetic and glutamic-pyruvic transaminase in routine laboratories. *J. Clin. Pathol.* 10, 394–399.
- Mori, T., Okumura, M., Matsuura, M., Ueno, K., Tokura, S., Okamoto, Y., Minami, S., Fujinaga, T., 1997. Effects of chitin and its derivatives on the proliferation and cytokine production of fibroblasts in vitro. *Biomaterials* 18, 947–951.
- Nelson, S.D., Mitchell, J.R., Timbrell, J.A., Snodgrass, W.R., Corcoran III, G.B., 1976. Isoniazid and isopronisid: activation of metabolites to toxic intermediate in man rat. *Science* 192, 901–903.

- Nishimura, K., Nishimura, S., Nishi, N., Saiki, I., Tokura, S., Azuma, I., 1984. Immunological activity of chitin and its derivatives. *Vaccine* 2, 93–99.
- O'Brien, R.J., Long, M.W., Floy, S.C., Lyle Jr., M.A., Snider, D.E., 1983. Hepatotoxicity from isoniazid and rifampicin among children treated for tuberculosis. *Pediatrics* 72, 491–493.
- Ohkawa, H., Onishi, N., Yagi, K., 1979. Assay for lipid peroxides in animal tissue by thiobarbituric acid reaction. *Anal. Biochem.* 95, 351–358.
- Okamoto, Y., Minami, S., Matsushashi, A., Sashiwa, H., Saimoto, H., Shigemasa, Y., Tanigawa, T., Tanaka, Y., Tokura, S., 1993. Polymeric *N*-acetyl-D-glucosamine (chitin) induces histionic activation in dogs. *J. Vet. Med. Sci.* 55, 739–742.
- Parekh, A.C., Jung, D.H., 1970. Cholesterol determination with ferric acetate–uranil acetate and sulphuric acid–ferrous sulphate reagents. *Anal. Chem.* 42, 1423–1427.
- Pepicelli, O., Fedele, E., Berardi, M., Raiteri, M., Levi, G., Greco, A., Ajmone-Cat, M.A., Minghetti, L., 2005. Cyclo-oxygenase-1 and -2 differently contribute to prostaglandin E2 synthesis and lipid peroxidation after in vivo activation of *N*-methyl-D-aspartate receptors in rat hippocampus. *J. Neurochem.* 93, 1561–1567.
- Ramachandran, P.C., 1980. Chemotherapy of tubercular meningitis with isoniazid plus rifampicin. Kinterim findings in a trial in children. *Ind. J. Tuberc.* 27, 54–56.
- Rice, E.W., 1970. Triglycerides in serum. In: Roedrick, P., McDonald, R.P. (Eds.), *Standard Methods in Clinical Chemistry*. Academic Press, New York, pp. 215–218.
- Sapelli, P.L., Baldassarre, V., Muzzarelli, R.A.A., Emanulli, M., 1986. *Chitin and Nature and Technology*. Plenum Press, New York, pp. 507–512.
- Saraswathy, S.D., Shyamala Devi, C.S., 2001. Modulating effect of Liv.100, an ayurvedic formulation on antituberculosis drug-induced alterations in rat liver microsomes. *Phytother. Res.* 15, 501–505.
- Sini, T.K., Santhosh, S., Mathew, P.T., 2005. Study of the influence of processing parameters on the production of carboxymethylchitin. *Polymer* 46, 3128–3131.
- Skakun, N.P., Slivka, Y., 1992. The correction of hepatotoxicity of antitubercular preparations with tocopherol acetate and riboxin. *Eksp. Klin. Farmakol.* 55, 52–54.
- Snider, D.E., Long, M.W., Cross, F.S., Farer, L.S., 1984. Six months isoniazid–rifampicin therapy for pulmonary tuberculosis. Report of a United States Health Service Cooperative trial. *Am. Rev. Respir. Dis.* 129, 573–578.
- Steele, M.A., Burk, R.F., Des Prez, R.M., 1991. Toxic hepatitis with isoniazid and rifampicin—a meta analysis. *Chest* 99, 465–468.
- Stenberg, D., 1976. Interconvertible enzymes in adipose tissue regulated by cyclic AMP-dependent protein kinase. *Adv. Cyclic Nucleotide Res.* 7, 157–198.
- Tasduq, S.A., Peerzada, K., Koul, S., Bhat, R., Johri, R.K., 2005. Biochemical manifestations of anti-tuberculosis drugs induced hepatotoxicity and the effect of silymarin. *Hepatol. Res.* 31, 132–135.
- Tokura, S., Ueno, K., Miyazaki, S., Nishi, N., 1997. Molecular weight dependent antimicrobial activity by chitosan. *Macromol. Symp.* 120, 1–9.
- Westphal, J.F., Vetter, D., Brogard, J.M., 1994. Hepatic side effects of antibiotics. *J. Antimicrob. Chemother.* 33, 387–401.
- Xing, R., Liu, S., Guo, Z., Yu, H., Wang, P., Li, C., Li, Z., Li, P., 2005. Relevance of molecular weight of chitosan and its derivatives and their antioxidant activities in vitro. *Bioorg. Med. Chem.* 13, 1573–1577.
- Yao, H.T., Chiang, M.T., 2002. Plasma lipoprotein cholesterol in rats fed a diet enriched in chitosan and cholesterol. *J. Nutr. Sci. Vitaminol.* 48, 379–383.

Biochemical Studies on the Antiulcer Effect of Glucosamine on Antioxidant Defense Status in Experimentally Induced Peptic Ulcer in Rats

Sethumadhavan Santhosh,^{1,*} Rangasamy Anandan,² Theruvathil Karunakaran Sini¹, Paruthapara Thomas Mathew,¹ and Thoduvayil Kurumpan Thankappan¹

¹*Fish Processing Division, Central Institute of Fisheries Technology, Cochin-682 029, Kerala, India*

²*Biochemistry and Nutrition Division, Central Institute of Fisheries Technology, Cochin-682 029, Kerala, India*

Received 18 April, 2005; Accepted 30 September, 2005

Summary The present study examined the antiulcer effect of glucosamine on mucosal antioxidant defense system in ibuprofen-induced peptic ulcer in male albino rats. The number of lesions in the gastric mucosa, volume of gastric juice, acid output, pepsin activity, lipid peroxides, reduced glutathione content and the activities of glutathione dependent antioxidant enzymes (glutathione peroxidase and glutathione-S-transferase) and antiperoxidative enzymes (catalase and superoxide dismutase) were determined. Prior oral administration of glucosamine significantly prevented the ibuprofen-induced increases in the number of lesions in the gastric mucosa, volume of gastric juice and acidity. It also maintained the activity of pepsin at near normal level. Oral pretreatment of glucosamine exerted a significant antioxidant effect by preventing ibuprofen-induced lipid peroxidation and by maintaining the level of reduced glutathione and the activities of mucosal antioxidant enzymes at near normalcy. The results of the present investigation indicate that the antiulcer activity of glucosamine is related to its ability to neutralize the hydrochloric acid secreted into the stomach and to its antioxidant capability to inhibit ibuprofen-induced lipid peroxidation.

Key Words: glucosamine, ibuprofen, ulcer, acid out put, antioxidant status

Introduction

Peptic ulcer is a non-malignant ulcer of the stomach or duodenum. A major causative factor (90% of gastric and 75% of duodenal ulcers) is chronic inflammation due to *Helicobacter pylori*, a spiral-shaped bacterium that lives in the acid environment of the stomach [1]. These ulcers can also be caused or worsened by drugs such as ibuprofen and other NSAIDs [2]. The lesion of peptic ulcer is a disruption

in the mucosal layer of the stomach or duodenum [3]. Peptic ulcer represents the imbalance between defensive factors that protect the mucosa and offensive or aggressive factors that disrupt this important barrier [4]. Some mucosal protective factors include the water-insoluble mucous gel layer, local production of bicarbonate, regulation of gastric acid secretion, and adequate mucosal blood flow [5]. Aggressive factors include the acid-pepsin environment, infection with *H. pylori*, and mucosal ischemia [6].

Ulcer induced by ibuprofen shows many metabolic and morphologic aberrations in the gastric mucosa of experimental animals similar to those observed in human peptic ulcer [7]. It is induced by a multiple

* to whom correspondence should be addressed.

†: 0091-4923-243558

mail: santhoshsethu@rediffmail.com

step mechanism. In particular, the peroxidation of mucosal endogenous lipids has been shown to be a major factor in the cytotoxic action of ibuprofen [8]. Ibuprofen-induced oxidative damage is generally attributed to the formation of the highly reactive hydroxyl radical ($\cdot\text{OH}$), stimulator of lipid peroxidation and source for destruction and damage to the mucosal cell membrane [9]. Alterations in the level of reduced glutathione (GSH) and the activities of glutathione-dependent antioxidant enzymes [glutathione peroxidase (GPX) and glutathione-S-transferase (GST)] and antiperoxidative enzymes [catalase (CAT) and superoxide dismutase (SOD)] have been reported in experimentally induced peptic ulcer condition in rats [10].

Traditional medicines have been the starting point for the discovery of many important modern drugs. This fact has led to the chemical and pharmacological investigations and general biological screening programs for natural products all over the world. Prawn shell, a main waste material of seafood industry, can be efficiently utilized by converting it into various useful by-products like chitin, chitosan, glucosamine *etc.* [11]. In traditional medicine, the shrimp and cuttlefish exoskeleton powder has been used to cure arthritis, diabetes, stomach disorders, epilepsy and various liver disorders. Earlier report [12] from our laboratory indicated that the supplementation of chitin and chitosan, along with feed, prevented HCl-ethanol mixture induced peptic ulcer in experimental rats. Glucosamine is the basic unit of chitin and chitosan and it is an essential component required for glycoprotein synthesis in living beings [13]. The beneficial actions of chitin and chitosan may be ascribable to their basic unit glucosamine. Hence, it is thought to be important to study the preventive effect of glucosamine on experimentally induced ulcer condition to derive a conclusion regarding the exact biochemical mechanism involved in the protective effects of shrimp and cuttlefish exoskeleton powder in alleviating the ulcerative disorders. In the present study, we assessed the antiulcer effects of glucosamine on mucosal antioxidant system in ibuprofen-induced ulcer in rats.

Materials and Methods

Drugs and chemicals

Ibuprofen was obtained from Cipla pharmaceuticals, Mumbai, India. Bovine serum albumin, tetraethoxypropane, epinephrine and D-galactosamine

were obtained from M/s Sigma Chemical Company, St. Louis, MO, USA. All other chemicals used were of analytical grade.

Animals

Male Wister rats (weighing 120–150 g) were allowed to undergo standard pelleted diet and water and housed under standard environmental conditions. Animals were deprived of food for 24 h prior to ulcer induction.

Experimental protocol

The experimental animals were divided into four groups of six rats each. Rats in Group I (normal control) received only the standard diet. In Group II, normal rats were treated with glucosamine (100 mg/kg/day, p.o for 20 days). In Group III, ulcer was induced by oral administration of ibuprofen (50 mg/kg, p.o) twice in a day at an interval of 12 h. Group IV was pretreated with glucosamine (100 mg/kg/day, p.o for 20 days) before induction of ulcer as described for Group III. At the end of the experiment, all four groups underwent surgery according to Takeuchi *et al.* [14] and the gastric juice was collected for 4 h. Rats were then killed with over dose of chloroform and the stomach was removed after the esophagus had been clamped. The gastric juice was centrifuged and the volume was noted. The total acidity was determined by titrating with 0.02 N NaOH with phenolphthalein used as indicator. The mucosal tissue scrapped was used for the determination of pepsin [15], lipid peroxides (LPO) [16], GSH [17], GPX [18], GST [19], CAT [20] and SOD [21].

The experiment was carried out as per the guidelines of Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA), New Delhi, India and approved by the Institutional Animal Ethics Committee (IAEC).

Statistical analysis

The values were expressed as mean \pm SD for six animals. The statistical significance was performed by using one way ANOVA followed by Tukey's test using SPSS10 Windows Software.

Results

Table 1 shows the number of lesions present in the gastric mucosa, volume of gastric juice, acid output and peptic activity in normal and experimental

Table 1. Number of lesions, volume of gastric juice, acid output and pepsin activity of the gastric mucosa of normal and experimental groups of rats.

	Group I Control	Group II Glucosamine (A)	Group III Ulcer (B)	Group IV (A+B)
Number of lesions	—	—	8.78±1.42 ^{a,b}	1.25 ^{a,b,c}
Volume of gastric juice (ml/4 h)	1.45±0.05	1.37±0.04	2.76±0.09 ^{a,b}	1.55±0.05 ^c
Acid output (μEq/4 h)	156±12.3	138±14.7	293±18.2 ^{a,b}	172±15.5 ^c
Pepsin (μmol tyrosine liberated/4 h)	723±58.1	698±63.5	576±41.9 ^{a,b}	682±54.7 ^c

Mean±SD for six animals in each group. ^a*p*<0.001 significantly different compared with Group I control animals; ^b*p*<0.001 significantly different compared with Group II glucosamine administered rats; ^c*p*<0.001 significantly different compared with Group III ulcer induced animals.

Table 2. Levels of lipid peroxides and reduced glutathione and the activities of glutathione peroxidase, glutathione-S-transferase, catalase and superoxide dismutase in the gastric mucosa of normal and experimental groups of rats.

	Group I Control	Group II Glucosamine (A)	Group III Ulcer (B)	Group IV (A+B)
LPO	1.02±0.02	0.96±0.01	2.45±0.05 ^{a,b}	1.18±0.01 ^c
GSH	5.02±0.36	4.89±0.25	2.12±0.12 ^{a,b}	4.73±0.18 ^c
GPX	179±14.2	195±16.8	108±7.3 ^{a,b}	172±15.5 ^c
GST	5.46±0.38	5.34±0.33	3.18±0.25 ^{a,b}	4.95±0.34 ^c
CAT	4.08±0.21	3.95±0.26	1.25±0.09 ^{a,b}	3.82±0.18 ^c
SOD	5.87±0.27	6.02±0.32	2.25±0.18 ^{a,b}	5.18±0.22 ^c

Mean±SD for six animals in each group. Values are expressed as: LPO, nmol/mg protein; GSH, nmol/g wet tissue; GPX, nmol GSH oxidized/min/mg protein; GST, μmol 1-chloro-2,4 dinitrobenzene conjugate formed/min/mg protein; CAT, μmol of H₂O₂ consumed/min/mg protein; SOD, one unit of the superoxide dismutase activity is the amount of protein required to give 50% inhibition of epinephrine autoxidation. ^a*p*<0.001 significantly different compared with Group I control animals; ^b*p*<0.001 significantly different compared with Group II glucosamine administered rats; ^c*p*<0.001 significantly different compared with Group III ulcer induced animals.

groups of rats. Oral administration of ibuprofen caused significant (*p*<0.001) increase in the number of lesions in the gastric mucosa of Group III rats as compared with that of Group I control animals. Significant (*p*<0.001) rise in the volume of gastric juice and acid output was observed in ulcer-induced animals as compared with Group I rats. Also a significant (*p*<0.001) reduction in the peptic activity was noted. Oral pretreatment with glucosamine significantly (*p*<0.001) reduced the number of ibuprofen-induced lesions and maintained the levels of gastric juice volume, acid output and peptic activity at near normal level.

Table 2 shows the levels of LPO and GSH and the activities of GPX, GST, CAT and SOD in the gastric mucosa of normal and experimental groups of rats. Significant (*p*<0.001) increase in the level of lipid peroxides was noticed in the gastric mucosa of Group III ulcer induced rats as compared with Group I animals. This was paralleled by the significant decline in the level of GSH and the activities of GPX, GST, CAT and SOD. Prior oral administra-

tion of glucosamine significantly (*p*<0.001) prevented all this ibuprofen-induced adverse effects and maintained the rats at near normal status.

Discussion

Significant (*p*<0.001) increase in the number of lesions present on the gastric mucosa is indicative of the severity of ibuprofen-induced ulcer (Table 1). Increased acid secretion into the stomach is the major causative factor in the development of human peptic ulcer disease [22]. The significant (*p*<0.001) rise noted in the volume of gastric juice and acidity with a concomitant decline in peptic activity in the Group III rats may be due to a consequence of increased permeability of the ulcerated gastric mucosa. This concurs with an earlier reported study [23], which indicated that the acid suppression therapy using proton pump inhibitors or histamine H₂-receptor antagonists is an effective newer approach for reducing the risk of occurrence of peptic ulcer.

In the present study, the oral pre-treatment with

glucosamine resulted in the significant ($p < 0.001$) reduction in number of lesions in the gastric mucosa, volume of gastric juice and acid output in Group IV animals as compared with Group III ulcer induced rats. Also the peptic activity was maintained at a level comparable to that of control animals. It probably did so by neutralizing the excessively secreted hydrochloric acid in the stomach. Mutoh *et al.* [24] have postulated that the adaptive cytoprotection in cultured gastric mucus producing cells is mediated by glucosamine rich mucus released in response to a mild irritant.

Lipid peroxidation reaction, a type of oxidative degeneration of polyunsaturated fatty acids, has been linked with altered membrane structure and enzyme inactivation. The level of lipid peroxidation was found to be significantly ($p < 0.001$) higher in the gastric mucosa of Group III ulcer-induced rats as compared with that of normal controls (Table 2). Significant increases in the levels of LPO in the ulcer-induced gastric mucosa have already been reported [25, 26]. Lipid peroxidation worsens the mucosal injury. The exact sources of peroxy radicals in peptic ulcer are not yet thoroughly understood. Xanthine oxidase has been considered as one of the major source of free radicals [27]. Another source may be neutrophils, since administration of non-steroidal anti-inflammatory drugs has already been reported to cause neutrophil infiltration into the gastric mucosa [28]. Jimenez *et al.* [29] proposed that the active oxygen species, derived both from xanthine oxidase and activated neutrophils, could play a role in the pathogenesis of gastric injury induced by ibuprofen. The metabolism of arachidonic acid *via* the lipoxygenase and cyclooxygenase pathways may also result in the formation of reactive oxygen species and other free radicals in the ulcerated gastric mucosa [30].

Prior administration of glucosamine resulted in a significant ($p < 0.001$) reduction in the level of LPO towards near normalcy as compared with Group III ulcer-induced rats, establishing its cytoprotective effect. The unpaired electron present in the ibuprofen-generated free radicals might have been trapped and subsequently dismutated by glucosamine by its antioxidant nature [31]. Reports by Chen *et al.* [32] showed that the dimeric form of glucosamine exerted free radical scavenging activity in a non-enzymatic system comprising of phenazine methosulfate and NADH.

GSH has a direct antioxidant function. It func-

tions by reaction with superoxide radicals, peroxy radicals and singlet oxygen, followed by the formation of oxidized glutathione and other disulphides [33]. Depletion of GSH results in enhanced lipid peroxidation, and excessive lipid peroxidation can cause increased GSH consumption [11], as observed in the present study (Table 2). This is in accordance with a previously reported study [34], which indicated that the depletion of gastric GSH by diethyl maleate produced gastric ulceration. This reduction might have resulted from the oxidation of GSH to ibuprofen-induced generation of free radicals. The depletion of GSH further enhances the susceptibility of the gastric mucosal cells to oxygen metabolites and acid mediated cell damage. Hiraishi *et al.* [35] has reported that exogenous GSH protects cultured gastric mucosal cells from oxidant-induced damage.

In the present study, the oral pretreatment with glucosamine maintained the level of GSH, which protects against oxidative damage by the way of regulating the redox status of proteins in the cell membrane [11], at near normalcy as compared with that of Group III ulcer-induced rats. This indicates that normal maintenance of mucosal GSH levels by glucosamine in ulcer rats may be the result, but not cause, of the cytoprotection by glucosamine.

Activities of glutathione-dependent antioxidant enzymes (GPX and GST) and antiperoxidative enzymes (CAT and SOD) were significantly ($p < 0.001$) lower in the gastric mucosa of Group III ulcer-induced rats as compared with that of Group I control animals (Table 2). GPX, an antioxidant enzyme, offers protection to the mucosal membrane from peroxidative damage [5]. A significant decrease in the activity of GPX makes mucosal membrane susceptible to ibuprofen-induced peroxidative damage, which leads to changes in the gastric mucosal composition and function. GST, another scavenging enzyme, binds to many different lipophilic compounds [33]; so it would be expected to bind ibuprofen and act for GSH conjugation reactions. The significant ($p < 0.001$) decrease in GST activity noted in this study might have been due to the decreased availability of GSH.

Reduction noticed in the activities of CAT and SOD in experimentally induced ulcerated mucosa is in line with earlier reported studies [5, 36], indicating that decline in the activities of these antiperoxidative enzymes leads to the formation of peroxy radicals, which are harmful to the gastric mucosa. In the present study, the prior oral administration of

glucosamine significantly ($p < 0.001$) prevented the ibuprofen-induced alterations in the activities of glutathione-dependent antioxidant enzymes (GPX and GST) and antiperoxidative enzymes (SOD and CAT) in the gastric mucosa of Group IV rats. It probably did so by the counteraction of ibuprofen-induced free radicals [37] due to its antioxidant nature.

The results of the present study indicate that the overall antiulcerogenic effect of glucosamine is probably related to its ability to neutralize the excess acid secreted into the stomach and to maintain near to the normal status the activities of the mucosal antioxidant enzymes and the level of GSH, which protect mucosa against oxidative damage by decreasing the lipid peroxidation and strengthening the mucosal barrier, and which are the first line of defense against exogenous and endogenous ulcerogenic agents.

References

- [1] Kuipers, E.J. and Malfertheiner, P.: *Helicobacter pylori* and nonmalignant diseases. *Helicobacter*, **9**, 29–34, 2004.
- [2] Shostak, N.A., Riabkova, A.A., Savel'ev, V.S., and Maliarova, L.P.: Gastrointestinal hemorrhages as complications of gastropathies associated with intake of nonsteroidal anti-inflammatory drugs. *Ter Arkh.*, **75**, 70–73, 2003.
- [3] Kivilaakso, E., Kiviluoto, T., Mustonen, H., and Paimela, H.: Gastric mucosal acid-base balance. *J. Intern. Med. Suppl.*, **732**, 63–68, 1990.
- [4] Gyires, K.: Gastric mucosal protection: from prostaglandins to gene-therapy. *Curr. Med. Chem.*, **12**, 203–215, 2005.
- [5] Prabha, T., Dora Babu, M., Priyambada, S., Agrawal, V.K., and Goel, R.K.: Evaluation of *Pongamia pinnata* root extract on gastric ulcers and mucosal offensive and defensive factors in rats. *Indian J. Exp. Biol.*, **41**, 304–310, 2003.
- [6] Sivri, B.: Trends in peptic ulcer pharmacotherapy. *Fundam. Clin. Pharmacol.*, **18**, 23–31, 2004.
- [7] Lloyd-Still, J.D., Beno, D.W., Uhing, M.R., Jiyamapa-Serna, V.A., and Kimura, R.E.: Ursodeoxycholic acid ameliorates ibuprofen-induced enteropathy in the rat. *J. Pediatr. Gastroenterol. Nutr.*, **32**, 270–273, 2001.
- [8] Wolfe, M.M.: Risk factors associated with the development of gastroduodenal ulcers due to the use of NSAIDs. *Int. J. Clin. Pract.*, **135**, 32–37, 2003.
- [9] Hariganesh, K. and Prathiba, J.: Effect of dimethylglycine on gastric ulcers in rats. *J. Pharm. Pharmacol.*, **52**, 1519–1522, 2000.
- [10] Sudheer Kumar, M., Sridhar Reddy, B., Kiran Babu, S., Bhilegaonkar, P.M., Shirwaikar, A., and Unnikrishnan, M.K.: Antiinflammatory and antiulcer activities of phytic acid in rats. *Indian J. Exp. Biol.*, **42**, 179–185, 2004.
- [11] Sini, T.K., Santhosh, S., and Mathew, P.T.: Study of the influence of processing parameters on the production of carboxymethylchitin. *Polymer*, **46**, 3128–3131, 2005.
- [12] Anandan, R., Suseela, M., and Nair, P.G.V.: Antiulcerogenic effects of chitin and chitosan on mucosal antioxidant defense system in HCl-ethanol induced ulcer in rats. *J. Pharm. Pharmacol.*, **56**, 265–269, 2004.
- [13] Wu, T., Zivanovic, S., Draughon, F.A., and Sams, C.E.: Chitin and chitosan-value-added products from mushroom waste. *J. Agric. Food. Chem.*, **52**, 7905–7910, 2004.
- [14] Takeuchi, K., Okabe, S., and Takagi, K.: A new model of stress ulcers in the rat with pylorus ligation and its pathogenesis. *Am. J. Dig. Dis.*, **21**, 782–788, 1976.
- [15] Anson, M.L.: The estimation of pepsin, trypsin, papain and cathepsin with haemoglobin. *J. Gen. Physiol.*, **22**, 79–89, 1938.
- [16] Ohkawa, H., Ohishi, N., and Yagi, K.: Assay for lipid peroxides in animal tissue by thiobabutaric acid reaction. *Anal. Biochem.*, **95**, 351–358, 1979.
- [17] Ellman, G.L.: Tissue sulfhydryl groups. *Arch. Biochem. Biophys.*, **82**, 70–77, 1959.
- [18] Paglia, D.E. and Valentine, W.N.: Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. *J. Lab. Clin. Med.*, **70**, 158–169, 1967.
- [19] Habig, W.H., Pabst, M.J., and Jackoby, W.B.C.: Glutathione-S-transferases: The first enzymatic step in enzymatic stepping mercapturic acid formation. *J. Biol. Chem.*, **249**, 7130–7139, 1974.
- [20] Takahara, S., Hamilton, B.H., Nell, J.V., Kobra, T.Y., Ogawa, Y., and Nishimura, E.T.: Hypocatalasemia: A new genetic carried state. *J. Clin. Invest.*, **29**, 610–619, 1960.
- [21] Misra, H.P. and Fridovich, T.: The role of superoxide ion in the auto-oxidation of epinephrine and a simple assay for superoxide dismutase. *J. Biol. Chem.*, **247**, 3170–3175, 1972.
- [22] Dembinski, A., Warzecha, Z., Ceranowicz, P., Brzozowski, T., Dembinski, M., Konturek, S.J., and Pawlik, W.W.: Role of capsaicin-sensitive nerves and histamine H1, H2, and H3 receptors in the gastroprotective effect of histamine against stress ulcers in rats. *Eur. J. Pharmacol.*, **508**, 211–221, 2005.
- [23] Welage, L.S.: Overview of pharmacologic agents for acid suppression in critically ill patients. *Am. J. Health. Syst. Pharm.*, **62**, S4–S10, 2005.
- [24] Mutoh, H., Ota, S., Hiraishi, H., Ivey, K.J., Terano, A., and Sugimoto, T.: Adaptive cytoprotection in cultured rat gastric mucus producing cells. Role of mucus and prostaglandin synthesis. *Dig. Dis. Sci.*, **40**, 872–878, 1995.
- [25] Yoshikawa, T., Yoshida, N., Miyagawa, H., Takemura,

- T., Tanigawa, T., Sugino, S., and Kondo, M.: Role of lipid peroxidation in gastric mucosal lesions induced by burn shock in rats. *J. Clin. Biochem. Nutr.*, **2**, 163–170, 1987.
- [26] Yoshikawa, T., Miyagawa, H., Yoshida, N., Sugino, S., and Kondo, M.: Increase in lipid peroxidation in rat gastric mucosal lesions induced by water-immersion restraint stress. *J. Clin. Biochem. Nutr.*, **1**, 271–277, 1986.
- [27] Yoshikawa, T., Yoshida, N., Naito, Y., Takemura, T., Miyagawa, H., Tanigawa, T., and Kondo, M.: Role of oxygen radicals in the pathogenesis of gastric mucosal lesions induced by water-immersion restraint stress and burn in rats. *J. Clin. Biochem. Nutr.*, **8**, 227–234, 1990.
- [28] Amagase, K., Yokota, M., Tsukimi, Y., and Okabe, S.: Characterization of “unhealed gastric ulcers” produced with chronic exposure of acetic acid ulcers to indomethacin in rats. *J. Physiol. Pharmacol.*, **54**, 349–360, 2003.
- [29] Jimenez, M.D., Martin, M.J., Alarcon de la Lastra, C., Bruseghini, L., Esteras, A., Herreras, J.M., and Motilva, V.: Role of L-arginine in ibuprofen-induced oxidative stress and neutrophil infiltration in gastric mucosa. *Free Radical Res.*, **38**, 903–911, 2004.
- [30] Argentieri, D.C., Ritchie, D.M., Ferro, M.P., Kirchner, T., Wachter, M.P., Anderson, D.W., Rosenthale, M.E., and Capetola, R.J.: Tepoxalin: a dual cyclooxygenase/5-lipoxygenase inhibitor of arachidonic acid metabolism with potent anti-inflammatory activity and a favorable gastrointestinal profile. *J. Pharmacol. Exp. Ther.*, **271**, 1399–1408, 1994.
- [31] Sal'nikova, S.I., Drogovoz, S.M., and Zupanets, I.A.: The liver-protective properties of D-glucosamine. *Farmako. Toksikol.*, **53**, 33–35, 1990.
- [32] Chen, A.S., Taguchi, T., Sakai, K., Kikuchi, K., Wang, M.W., and Miwa, I.: Antioxidant activities of chitobiose and chitotriose. *Biol. Pharm. Bull.*, **26**, 1326–1330, 2003.
- [33] Anandan, R., Rekha, R.D., and Devaki, T.: Protective effect of *Picrorhiza kurroa* on mitochondrial glutathione antioxidant system in D-galactosamine-induced hepatitis in rats. *Curr. Sci.*, **76**, 1543–1545, 1999.
- [34] Lu, S.C., Kuhlenkamp, J., Robert, A., and Kaplowitz, N.: Role of glutathione status in protection against ethanol-induced gastric lesions. *Pharmacology*, **38**, 57–60, 1989.
- [35] Hiraishi, H., Terano, A., Ota, S., Mutoh, H., Sugimoto, T., Harada, T., Razandi, M., and Ivey, K.J.: Protection of cultured rat gastric cells against oxidant-induced damage by exogenous glutathione. *Gastroenterology*, **106**, 1199–1207, 1994.
- [36] Raghavendran, H.R., Sathivel, A., and Devaki, T.: Efficacy of brown seaweed hot water extract against HCl-ethanol induced gastric mucosal injury in rats. *Arch. Pharm. Res.*, **27**, 449–453, 2004.
- [37] Balasubramanian, T., Somasundaram, M., and Felix, A.J.: Taurine prevents ibuprofen-induced gastric mucosal lesions and influences endogenous antioxidant status of stomach in rats. *Sci. World J.*, **4**, 1046–1054, 2004.



Study of the influence of processing parameters on the production of carboxymethylchitin

T.K. Sini, S. Santhosh*, P.T. Mathew

Fish Processing Division, Central Institute of Fisheries Technology, Matsyapuri (P.O.), Cochin 682 029, Kerala, India

Received 11 October 2004; received in revised form 2 February 2005; accepted 2 February 2005

Available online 5 March 2005

Abstract

Carboxymethylchitin was prepared at different reaction temperatures and from alkali chitin with different concentrations of alkali. Properties of the product were studied. Alkali chitin were prepared using freshly prepared sodium hydroxide of 45, 50, 55, 60 and 65% (w/w) concentration and carboxymethylated using monochloroacetic acid at controlled (35–40 °C) and uncontrolled (30–80 °C) temperature conditions. Molecular weight, viscosity, degree of deacetylation, etc. of the resultant product, i.e. carboxymethylchitin were determined. It was found that the reaction temperature has a profound influence on the property of the product than alkali concentration. A hygroscopic and completely water-soluble product was formed. Optimum conditions for the production of carboxymethylchitin were found to be 60% NaOH concentration and at 35–40 °C reaction temperature. At these conditions, it was obtained with a molecular weight of 4.11×10^6 Da, viscosity 1926 cP and degree of deacetylation 45.02%.

© 2005 Elsevier Ltd. All rights reserved.

Keywords: Chitin; Carboxymethylchitin; *Parapeneopsis styliifera*

1. Introduction

Aquatic organisms like shrimp, crab, etc. contribute a very good participation in our foods. Processing of these organisms gives head, shell, etc as waste materials. They will be easily spoiled and arises threats to the public health. Conversion of waste materials into useful products is one of the solutions of environmental pollution, nowadays. Shrimp shell contains 8–10% chitin [7], which is the most abundant biopolymer after cellulose [6]. So, shrimp shell can be converted to chitin, a nitrogen containing polysaccharide.

Polysaccharides and their derivatives hold a major part in our lives as medicines, cosmetics, textiles, paper, food and other branches of industry because of their unique nature in properties such as low toxicity, biocompatibility, hydrophobicity, etc. [12]. In the midst of these polysaccharides,

people show growing interest in chitin, since, it is widely distributed in nature [8,9]. The most commonly prepared derivative of chitin is chitosan by partial deacetylation of chitin [8]. The importance of chitin and chitosan has grown partly because they are renewable and biodegradable source of materials and partly because of the recent increased applications in biology, biotechnology and medicine. Carboxymethylchitin is another value shot derivative of chitin. The conversion of chitin into carboxymethylchitin came into practice by carboxymethylation reaction [2,9,13].

Carboxymethylchitin has successfully proven its use in the field of cosmetics as moisturizer, smoothener, cell activator and a cleaner for face skin conditioning. Carboxymethylchitin is widely used in food products also. Another innovative use of Carboxymethylchitin is in wound dressing, due to its hydrophobicity.

When, molecular characteristics of carboxymethylchitin are concerned the information available is very poor [3,5]. Here, comes the importance of the present work. This effort puts light on the preparation of carboxymethylchitin at different alkali concentrations and temperatures and the study of its properties.

* Corresponding author.

E-mail address: santhoshsethu@rediffmail.com (S. Santhosh).




Fig. 1. Structure of carboxymethylated monomer.

2. Experimental

Raw material—*Parapanaeopsis stylifera* procured from processing plant near Cochin, Kerala, India.

All chemicals were of analytical grade and were obtained from BDH (India).

Dialysis membrane obtained from Sigma chemicals (UK), which can retain material of molecular weight 12,000 or greater. Capacity—640 ml/feet.

Freeze drier-Heto drywinner(Denmark).

2.1. Preparation of chitin [10]

The fresh shell waste was washed in water and heated to boiling with 0.5% aqueous NaOH in the ratio of 2:3 for 30 min. Alkali was drained off and the shell was washed free of alkali. The residue was demineralised by immersing in 1.25 N HCl. at room temperature for 1 h with intermittent stirring. The residue was filtered, washed repeatedly with water to make it acid free. Then the product, chitin obtained was dried and pulverized.

2.2. Preparation of carboxymethylchitin

Carboxymethylchitin was obtained by the treatment of alkali chitin with monochloroacetic acid.

Fresh alkali chitin was prepared with different concentrations of alkali viz. 45, 50, 55, 60 and 65% (w/w) NaOH.

2.2.1. Alkali chitin

About 10 g of chitin was mixed well with 40 ml respective alkali prepared and kept in ice for 1 h. Then, it was kept overnight at -20°C .

Table 1
Yield of carboxymethylchitin prepared at different alkali concentrations and temperature

Percentage of NaOH for the preparation of alkali chitin	Yield in percentage	
	Uncontrolled temperature	Controlled temperature
45	36.7 ± 1.5	37.4 ± 1.4
50	49.8 ± 2.12	44.7 ± 1.2
55	56.5 ± 1.7	52.0 ± 1.6
60	69.6 ± 2.2	65.8 ± 1.85

2.2.2. Carboxymethylchitin

To the alkali chitin prepared, poured 200 ml isopropanol and monochloroacetic acid added, at intervals, with continuous stirring. Addition of monochloroacetic acid was stopped, when the whole mixture became neutral. When monochloroacetic acid was added to the mixture, reaction temperature went on increasing. For each concentration of alkali used, one set of carboxymethylation was allowed to take place at room temperature and in another set of experiment, reaction temperature maintained at $35\text{--}40^{\circ}\text{C}$ by using ice bath.

Resulting viscous solution was dissolved in 1 l of distilled water and then precipitated using 5 l of acetone. The precipitated mass was carboxymethylchitin and it was redissolved in distilled water. The solution was then dialysed for 24 h in running water, using a dialysis membrane, for the removal of salts and then freeze dried and stored.

Important quality parameters of carboxymethylchitin like molecular weight, degree of deacetylation, viscosity, yield, etc. were estimated. Molecular weight was determined by using Schott Gerate AVS 410 equipment [14]. By applying Schrodinger equation, degree of deacetylation was measured [11] using Spectronic Genysis spectrophotometer. Fungilab viscometer was used for viscosity determination.

3. Results and discussion

The carboxymethylchitin (Fig. 1) was prepared with alkali chitin using different concentrations of NaOH viz. 45, 50, 55, 60 and 65% (w/w). When 45, 50, 55 and 60% NaOH were used for alkali chitin preparation and when it was subjected to carboxymethylation, temperature was raised from 30 to 80°C as more and more monochloroacetic acid was added to it and finally a neutral viscous milky solution was obtained, which was water soluble. This water-soluble product was carboxymethylchitin. Carboxymethylation was a highly exothermic reaction and so only the temperature was raised. It was found that when alkali chitin prepared with NaOH, having concentration above 60%, was used for carboxymethylation, the whole mass turned to a colloidal elastic material as the reaction proceeded, which did not show characteristic reactions of carboxymethylchitin. It implied that at higher alkali concentration of alkali chitin, carboxymethylation may not be possible.

In another set of experiment, with the same as above concentrations of alkali chitin (45, 50, 55 and 60%), reaction temperature was maintained at $35\text{--}40^{\circ}\text{C}$ and the reaction was slow compared to the other set, where, temperature remained uncontrolled. It was also found that carboxymethylation reaction was not possible at low temperatures below 35°C .

Yields of carboxymethylchitin obtained at different experimental conditions were given in Table 1. Maximum

Table 2
Viscosity of 1% solution of carboxymethylchitin in water

Percentage of NaOH for the preparation of alkali chitin	Viscosity of 1% solution (cP)	
	Uncontrolled temperature	Controlled temperature
45	36 ± 2	2114 ± 15
50	30 ± 1	2010 ± 10
55	28 ± 1	1928 ± 7
60	22 ± 2	1926 ± 12

yield was obtained when 60% NaOH was used for the preparation of alkali chitin, which was used for carboxymethylation. Under controlled temperature, carboxymethylation was very slow and the reaction would not occur completely. As a result, we got comparatively less yield at controlled temperature than uncontrolled temperature condition.


Viscosity of the samples prepared at controlled and uncontrolled temperatures using different concentrations of alkali were given in Table 2. It is well clear from the table that there was remarkable difference in viscosity of the samples prepared at controlled and uncontrolled temperatures. Thus, it was found that the temperature of the reaction had a great influence on the viscosity of the product as was in the case of the production of chitosan [8]. There was a reduction in the viscosity of the final product with increasing alkali concentration for the preparation of alkali chitin. Carboxymethylchitin with good viscosity was obtained at controlled temperature, since, the operating temperature was around 35 °C. But, at uncontrolled temperature condition, temperature was increased to a large extent and as a result thermal degradation occurred [1, 4] and viscosity decreased.

Molecular weight of the samples prepared was given in Table 3. Considerable difference was noticed in the case of molecular weight when the samples were prepared under controlled and uncontrolled conditions. It was found that degradation of the compound had taken place at high temperatures. Due to degradation of the product, depolymerisation of chains occurred and as a result molecular weight was decreased. The effect of alkali concentration on the molecular weight of the product under the same temperature conditions was negligible.

The degree of deacetylation of the product prepared

Table 3
Molecular weight of carboxymethylchitin prepared from alkali chitin of different alkali concentrations

Percentage of NaOH for the preparation of alkali chitin	Molecular weight ($\times 10^6$ Da)	
	Controlled temperature	Uncontrolled temperature
45	4.86 ± 0.55	1.33 ± 0.21
50	4.92 ± 0.23	1.01 ± 0.10
55	4.21 ± 0.33	1.21 ± 0.18
60	4.11 ± 0.42	1.29 ± 0.32



Graph-I. Degree of deacetylation of carboxymethylchitin.

under different conditions were shown in Graph-I. Degree of deacetylation up to 75% was reported in products obtained by carboxymethylation at uncontrolled temperature condition. But in the controlled condition, degree of deacetylation was in the range of 40–45% depending on the alkali concentration.

Thus, a 65.8% yield of good quality carboxymethylchitin with high viscosity and high molecular weight was obtained when alkali chitin prepared at 60% NaOH was used for carboxymethylation and the reaction temperature was controlled at 35–40 °C. It is a hygroscopic product, which contains 10–12% moisture. From the experiment, it is well understood that the reaction temperature and alkali concentration had a great influence on its properties like viscosity, molecular weight and degree of deacetylation. Product prepared at controlled temperature condition had better properties than that obtained at uncontrolled conditions.

Acknowledgements

We acknowledge and thank Dr K. Devadasan, Director, Central Institute of Fisheries Technology, for providing us facilities to carry out this work. We thank Dr K.G. Ramachandran Nair, HOD, Fish Processing division, for his support and suggestions.

References

- [1] Barry JH, James NH. *Polym Int* 2000;49:943–8.
- [2] Danilov SN, Plisko EA. *J Gen Chem (Russia)* 1961;31:469.
- [3] Hackman RH, Goldberg M. *Carbohydr Res* 1974;38:35.
- [4] Home HK, Foros H, Petterser H, Dornish M, Smidsrod O. *Carbohydr Polym* 2001;46:287–94.
- [5] Inouc Y, Kaneko M, Tokura S. *Rep Prog Polym Phys Jpn* 1982;25:759.
- [6] Moret A, Rubio J. *Miner Eng* 2003;16:716.

C9101

- [7] Legarreta GI, Zakaria Z, Hall GM. *Advances in chitin science* (Vol I), France, 1996;399–406.
- [8] Muzarelli RAA. *Chitin*. New York: Pergamon Press; 1977.
- [9] Plisko EA, Nudiga LA, Danilov SN. *Uspechi khimii*(Progr Chem) 1977;46:1470.
- [10] Madhavan P, Ramachandran NKG. *Fish. Technol* 1974;11:50–3.
- [11] Muzarelli RAA, Rocchetto R. *Annl Chem Acta* 1973;64:371–9.
- [12] Tombs MP, Harding SE. *An Introduction to polysaccharide biotechnology*. London: Taylor and Francis; 1997.
- [13] Vichoreva GA, Gladyshev DY, Bazi MR, Kuznersova SY, Galbraich LS, Lavut EE, et al. *Acta Polym* 1991;42:330.
- [14] Rutherford F, Austin PR. *Proceedings of First Int Conf Chitin/Chitosan*, MIT, Cambridge 1978;182–92.

