

ОБЪЕДИНЕННЫЙ
ИНСТИТУТ
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ

Дубна

E1-2006-97

V. A. Bednyakov¹, J. A. Budagov¹, A. V. Gladyshev¹,
D. I. Kazakov¹, G. D. Khoriauli^{1,2}, J. I. Khubua^{1,3}

THE SEARCH FOR GLUINOS WITH ATLAS AT LHC

Submitted to «Particles and Nuclei, Letters»

¹ Joint Institute for Nuclear Research, Dubna

² Tbilisi State University, Tbilisi, Georgia

³ Institute of High Energy Physics and Informatization, Tbilisi, Georgia

2006

The minimal supersymmetric version of the Standard Model with universal soft supersymmetry breaking terms induced by supergravity (mSUGRA) [1] has a minimal set of free parameters:

$$m_0, m_{1/2}, \text{sign } \mu, A_0, \text{tg } \beta = v_2/v_1,$$

where m_0 and $m_{1/2}$ are respectively common masses for scalar and spinor superpartners at the unification scale, μ is the Higgs mixing parameter, A_0 is the trilinear soft supersymmetry breaking parameter and $\text{tg } \beta$ is the ratio of vacuum expectation values of two Higgs fields. The allowed values of these parameters are limited by both theoretical and experimental constraints as well as by the recent astrophysical data [2].

In supersymmetric models with conserved R parity, superpartners of ordinary particles may be created only in pairs, which leads to the existence of the stable lightest supersymmetric particle (LSP). Usually, it is the lightest neutralino – neutral, massive and weakly interactive particle. The mass spectrum of superpartners at some scale in mSUGRA can be calculated by solving renormalization group equations with parameters mentioned above being initial conditions at the unification scale.

In the present analysis, the region with large scalar masses m_0 and small fermion masses $m_{1/2}$ was studied. In this case, LSP properties are consistent with the observed relic density, i.e. they can serve as cold dark matter particles, and, on the other hand, can naturally explain the excess of diffuse galactic gamma rays observed by EGRET [3]. The set of the parameters used in the analysis (let us call it the EGRET point) is the following (Fig. 1):

$$m_0 = 1400 \text{ GeV}, m_{1/2} = 180 \text{ GeV}, \text{sign } \mu = +1, A_0 = 0, \text{tg } \beta = 50.$$

This region has not been intensively studied yet, though it is within the reach of Tevatron and forthcoming LHC. It appears to be very interesting phenomenologically, because of the mass splitting between light gauginos and heavy squarks and sleptons. The cross sections for chargino and neutralino production in this case are relatively large not being suppressed by masses and being comparable with squark and gluino production. The latter being enhanced by strong interactions remains suppressed by the heaviness of squarks. This means that in the EGRET region leptonic channels are not suppressed and so might give clear leptonic signature for supersymmetry in the upcoming LHC experiments [4].

The detection of a SUSY-like signal at LHC corresponding to the EGRET point of mSUGRA would be a strong indication both for supersymmetry and for solution of the dark matter problem. These parameters (the EGRET point) were used as input for the ISAJET code [5] which calculated the superparticle spectrum. Later, the PYTHIA generator [6] used the spectrum for event

generation. It is worth mentioning that the whole generation process was performed within the ATLAS software ATHENA [7]. Superparticle spectrum and cross sections of different processes depend on the chosen parameters of mSUGRA. The common property of SUSY-like processes in models with conserved R parity is undetectable LSP in the final state which is rather heavy and, thus, takes away quite high (and undetectable) transverse momentum. The condition for choosing a certain SUSY channel was not only the large cross section but also a peculiar signature which would tell it from different Standard Model background events. In the EGRET point the process of pair gluino production and their subsequent decay appeared to be the most interesting:

$$\begin{array}{c}
 gg \rightarrow \tilde{g}\tilde{g} \\
 \downarrow \\
 \bar{b} + b + \tilde{\chi}_2^0 \\
 \downarrow \\
 \mu^- + \mu^+ + \tilde{\chi}_1^0
 \end{array}
 \quad (1)$$

Here we assume that both gluinos decay in the same way. So, in the final state the gluino pair gives four b quarks (b jets), four muons and a pair of the lightest stable neutralinos $\tilde{\chi}_1^0$ giving the high missing transverse momentum. There are B hadrons in these jets, and, in general, the event could have four secondary vertices, which allows one to reduce the background even at the trigger level (Fig. 2).

Figure 3 shows the event inside the ATLAS pixel detector in cylindrical first layer ($R \approx 4$ cm). The cross section of the hard $gg \rightarrow \tilde{g}\tilde{g}$ process at $\sqrt{s} = 14$ TeV is quite large. It varies between 5.6–14.2 pb (PYTHIA) at the chosen values of SUSY parameters (the EGRET point). The cross section depends on the parton distribution function (PDF) for the proton and its dependence on Q^2 . We chose the Q^2 independent PDF model GRV94D (DIS) [9] and put $Q_{\min}^2 = 0.5$ GeV. In this case, the cross section is ≈ 13 pb.

The transverse momentum P_t distribution for one gluino decay products (two b quarks forming two b jets as well as two muons and the lightest neutralino) is shown in Fig. 4. One can see that neutralino takes away quite high transverse momentum. It is worth noticing that in the real experiment one can measure only the total missing (undetectable) energy of two neutralinos \cancel{E}_t . Figure 5 shows the total transverse momentum of two neutralinos. It is clear, that the careful reconstruction will allow one to detect such a high loss in the total measure transverse energy.

The b tagging of all b jets and the careful reconstruction of their energy are extremely important for the study of the considered processes. B hadrons with b quarks inside live rather a long time and can move rather far away of the creation point (the cross point of initial proton beams). As a result, it allows one to observe a secondary vertex of B -hadron decay at a certain distance from the primary beams collision initial vertex. This secondary vertex allows one to tag hadronic jets from b quarks. Figure 6 shows the distribution of the free path of B hadrons provided all four B hadrons have free paths more than $100 \mu\text{m}$ simultaneously. One can see that 94% of events satisfy this condition.




Fig. 1. *a*) MSSM parameter space. Light blue line corresponds to the region consistent with WMAP data. One can distinguish between different regions marked by digits. 5 — is the EGRET region. *b*) The light shaded (blue) area is the enlarged region preferred by EGRET data for $\tan \beta = 50$, $\mu > 0$ and $A_0 = 0$. The excluded regions where the stau would be the LSP, or the electroweak symmetry breaking fails, or the Higgs boson is too light, are indicated by dots




Fig. 2. Topology of the «half of the event» under study with two b jets, two muons and two secondary vertices

All above-presented distributions were performed for 150 events, which is expected for ATLAS detector after a year of running with total LHC luminosity $10^{34} \text{ cm}^{-2}\text{s}^{-1}$ for the chosen values of SUSY parameters, parton distribution functions and their Q^2 dependence. Other PYTHIA parameters were set to their default values. The process (1) has a very peculiar signature (four b jets, four muons and high transverse momentum); however, there is a mimic background process with only Standard Model particles (top-quarks and gluons) with alike




Fig. 3. Generation of the process 1 inside the cylindrical (Layer 1) pixel detector in the XY plane (transversely to the beam). One can see four muon tracks (green lines), two tracks from neutralino (light blue lines), four jets (dark blue lines) and one long-living B meson (red line). Visualisation has been performed with the help of ATLAS GeoModel HitDisplay [8]

signature:

$$\begin{aligned}
 gg(q\bar{q}) &\rightarrow t \bar{t} \quad t \bar{t} \\
 &\quad \downarrow \quad \downarrow \\
 &\quad \bar{b} + W^- (\rightarrow \mu^- + \nu_\mu) \\
 &\quad b + W^+ (\rightarrow \mu^+ + \bar{\nu}_\mu).
 \end{aligned} \tag{2}$$

In this process (2) we again have four b quarks (four b jets), four muons and the missing transverse momentum, but from two neutrinos. Now this background process is under study; however, we can make some qualitative conclusions. The cross section of the processes $gg(q\bar{q}) \rightarrow t\bar{t}t\bar{t}$ varies (depends on the choice of Q^2 scale) in the range of 2.6–9.4 (0.5–1.5) fb (Acer MC generator [10]). So, at the total LHC luminosity, $10^{34} \text{ cm}^{-2} \cdot \text{s}^{-1}$, the expected rate of $gg, q\bar{q} \rightarrow t\bar{t}t\bar{t}(W^+bW^-bW^+bW^-b)$ is up to 3500 events per




Fig. 4. Transverse momentum distributions for the products of one of the gluinos decay. a) The total P_t for the $b\bar{b}$ pair; b) P_t for the lightest stable neutralino $\tilde{\chi}_1^0$ from the heavier neutralino decay; c) total P_t for the muon pair




Fig. 5. Total missing transverse momentum P_t of two neutralinos. Event selection is made assuming that the total P_t of gluino pair is less than 10 GeV. Since in our study the initial state radiation was not taken into account, all events satisfy this condition




Fig. 6. The free path of B hadrons created in the process (1) before their decay. The criterion for the event selection is that the free path is not less than $100 \mu\text{m}$ for each of four B hadrons in the same event. 94% of events satisfy this condition

year. We require that all of four W bosons decay into muon-neutrino pair ($BR(W^- \rightarrow \mu^- \nu_\mu \approx 0.1082)$) to give the same signature as one for the SUSY process 1. Then the SM background (process 2) rate is < 0.5 events per year. So, the preliminary estimation of the SM background gives its negligible contribution to the total signal. Moreover, due to the negligible (0 in our analysis) neutrino mass the missing transverse momentum is smaller than in the process (1). That is, there is a possibility to distinguish between the background (2) and a SUSY-like process (1) by choosing events with the large missing transverse momentum (larger than 50 GeV). Now we perform the full analysis (event simulation and reconstruction) of these processes using ATHENA.

To conclude, the prospects for ATLAS observation of a SUSY-like signal from two gluinos are under investigation. The region of the mSUGRA relevant to SUSY dark matter interpretation of the EGRET observation is considered. The cross section of the two gluinos production via gluon-gluon fusion, $gg \rightarrow \tilde{g}\tilde{g}$, is rather high in the region. The clear signature of the process is four b jets, four muons and a rather large missing transverse energy (due to two lightest SUSY particles, the neutralinos). This large transverse missing energy carried away by two neutralinos allows the relevant Standard Model background to be reduced significantly. The generation and reconstruction processes are performed within ATHENA ATLAS software.

Acknowledgements. Financial support of the Russian Foundation for Basic Research (grants # 05-02-17603, # 06-02-04003) and the grant from the President of Russian Federation for support of leading scientific schools (NSh-5362.2006.2) is kindly acknowledged.

References

1. Nilles H. P. // Phys. Rep. 1984. V. 110. P. 1;
 Haber H. E., Kane G. L. // Phys. Rep. 1985. V. 117. P. 75;
 Haber H. E. Introductory Low-Energy Supersymmetry. Lectures Given at TASI

1992. SCIPP 92/33. 1993; hep-ph/9306207;
Kazakov D.I. Beyond the Standard Model: in Search of Supersymmetry. Lectures at the European School of High-Energy Physics. 2000; hep-ph/0012288;
Kazakov D.I. Beyond the Standard Model. Lectures at the European School of High-Energy Physics. 2004; hep-ph/0411064.
2. *Bennett C.L. et al.* // *Astrophys. J. Suppl.* 2003. V. 148. P. 1;
Spergel D.N. et al. // *Ibid.* P. 175.
 3. *Hartman R.C. et al. (EGRET Collab.)* // *Astrophys. J. Suppl.* 1999. V. 123. P. 79;
EGRET Public Data Archive. <ftp://coss.c.gsfc.nasa.gov/compton/data/egret/>
 4. *Bogachev D. Yu. et al.* Light Superpartners at Hadron Colliders // *Intern. J. Mod. Phys. A* (in press).
 5. *Paige F.E. et al.* ISAJET 7.69: A Monte Carlo Event Generator for pp , Anti- pp , and e^+e^- Reactions. hep-ph/0312045.
 6. *Sjostrand T. et al.* Pythia 6.3 Physics and Manual. FERMILAB-PUB-03-457, LU-TP-03-38; hep-ph/0308153.
 7. Athena Core Software.
<http://atlas-computing.web.cern.ch/atlas-computing/packages/athenaCore/athenaCore.php>
 8. The HEPVis/v-atlas.
<http://boudreau.home.cern.ch/boudreau/v-atlas-hepvis.htm>
 9. *Gluck M., Reya E., Schienbein I.* // *Phys. Rev. D.* 1999. V. 60. P. 054019;
Erratum // *Phys. Rev. D.* 2000. V. 62. P. 019902.
 10. *Kersevan B.P., Richter-Was E.* The Monte Carlo Event Generator AcerMC Version 2.0 with Interfaces to PYTHIA 6.2 and HERWIG 6.5. TPJU-6-2004; hep-ph/0405247.

Received on August 24, 2006.

Бедняков В. А. и др.

E1-2006-97

Поиск глюино на LHC с помощью детектора ATLAS

Исследованы перспективы наблюдения сигнала, подобного суперсимметрии, от двух глюино в реакции $pp \rightarrow \tilde{g}\tilde{g}$, в определенной области пространства параметров mSUGRA (суперсимметричного расширения стандартной модели с мягким нарушением суперсимметрии за счет эффектов супергравитации) с помощью детектора ATLAS. В данной области сечение рождения пары глюино в результате глюонного слияния $gg \rightarrow \tilde{g}\tilde{g}$ является довольно большим и достигает 13 пб. Для отбора событий используется отчетливая сигнатура процесса (4 адронные струи, 4 мюона и до 4 вторичных вершин); конечными продуктами распада каждого глюино являются $b\bar{b}$ - и $\mu^+\mu^-$ -пары и легчайшая суперсимметричная частица — нейтралино. Довольно большой недостающий поперечный импульс, уносимый парой нейтралино, является важным отличительным признаком события и позволяет значительно уменьшить количество соответствующих стандартной модели фоновых событий. Генерация и реконструкция событий выполнены при помощи принятого для ATLAS программного обеспечения ATHENA.

Работа выполнена в Лаборатории теоретической физики им. Н. Н. Боголюбова ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна, 2006

Bednyakov V. A. et al.

E1-2006-97

The Search for Gluinos with ATLAS at LHC

Prospects for ATLAS observation of a SUSY-like signal from two gluinos $pp \rightarrow \tilde{g}\tilde{g}$ are investigated within a certain region of the mSUGRA parameter space, where the cross section of the two gluinos production via gluon-gluon fusion, $gg \rightarrow \tilde{g}\tilde{g}$, is estimated at a rather high level of 13 pb. The event selection trigger uses a very clear signature of the process (4 jets + 4 muons + up to 4 secondary vertices topology), when final decay products of each gluino are b -anti- b and muon-anti-muon pairs and the lightest SUSY particle, the neutralino. Rather high transverse missing energy carried away by two neutralinos is an essential signature of the event and allows the relevant Standard Model background to be reduced significantly. The generation and reconstruction processes are performed by means of the ATLAS common software framework ATHENA.

The investigation has been performed at the Bogoliubov Laboratory of Theoretical Physics, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna, 2006